



# VOLUME - I & II

#### Based on the updated New Textbook

### Salient Features

- Prepared as per the updated New Textbook
- Exhaustive Additional MCQs, VSA, SA, LA questions with answers are given in each chapter.
- All the objective type (1 Mark) questions, are given with 4 options.
  - (i) Choose the correct option
  - (ii) Match the following
  - (iii) Fill in the blanks
  - (iv) Choose the odd one out
- (v) Choose the correct pair
- (vi) Choose the incorrect pair
- (vii) Assertion-Reason
- (viii) Choose the correct or incorrect statements
- Govt. Model Question Paper (2018) [Govt. MQP-2018], First Mid-Term Test (2018) [First Mid-2018], Quarterly Exam 2018 [QY-2018], Half Yearly Exam 2018 [HY-2018], March 2019 & 2020 [Mar.-2019 & 2020], June 2019 [Jun.-2019] Quarterly Exam 2019 [QY-2019], Half Yearly Exam 2019 [HY. 2019] and September 2020 [Sep. 2020] are incorporated in the appropriate sections.
- Govt. Suppl. Exam September 2020 Question Paper is given.



Chennai

Lab Manua

2021-22 Edition

© Reserved with Publishers

ISBN : 978-93-5330-061-6

Code No : SG 264

#### Author :

Mr.J.M.Joseph *M.sc.,B.Ed.,* 

Chennai.

#### **Reviewed by :**

Mr. P. George Paul *M.Sc.,M.Ed., M.Phil, PGDCA* Vice-Principal Annai Veilankanni's Matric Hr. Sec. School 81, VGP Salai, Saidapet, Chennai - 600 015

#### **Head Office:**

1620, 'J' Block, 16th Main Road, Anna Nagar,

Chennai	- 600	040.

- Phones : 044-4862 9977, 044-486 27755
- Mobile : 81242 01000/ 81243 01000
- e-mail : orders@surabooks.com
- website : www.surabooks.com

#### **For More Information - Contact**

Queries	:	enquiry@surabooks.com
For Order	:	orders@surabooks.com
Contact	:	80562 94222 / 80562 15222
Whatsapp	:	8124201000 / 9840926027
<b>Online Site</b>	:	www.surabooks.com
For Free Study Materia	als Vi	sit <b>http://tnkalvi.in</b>

#### Also available for Std. - XI

#### Guides :

- Sura's Tamil
- Sura's English
- Sura's Mathematics (EM/TM)
- Sura's Physics (EM/TM)
- Sura's Chemistry (EM/TM)
- Sura's Bio Botany (EM/ TM)
- Sura's Bio Zoology (EM/ TM)
- Sura's Botany Long Version (EM/ TM)

- Sura's Zoology Long Version (EM/ TM)
- Sura's Computer Science (EM/TM)
- Sura's Computer Applications (EM/TM)
- Sura's Computer Technology (EM/TM)
- Sura's Commerce (EM/TM)
- Sura's Economics (EM/TM)
- Sura's Accountancy (EM/TM)
- Sura's Business Maths (EM)

Also available 1 Mark Q & A (EM/TM),2,3 Marks (EM/TM) and 5 Marks Q & A (EM / TM) for all Subjects.

## NOTE FROM PUBLISHER

It gives me great pride and pleasure in bringing to you **Sura's Physics** guide for **11<sup>th</sup> Standard**. It is prepared as per the New Syllabus and New Textbooks Vol. I & II. A deep understanding of the text and exercises is rudimentary to have an insight into the subject. The students have to carefully understand the topics and exercises.

Our guide encompasses all the requirements of the students to comprehend the text and the evaluation of the textbook.

It will be a teaching companion to teachers and a learning companion to students.

- Concept Map, Must know Definitions are given in all units.
- Exhaustive Additional MCQs, VSA, SA, LA, HOTS questions with answers are given in each units.
- These features will help students practice and learn effectively all the sections of the textbooks.

In order to learn effectively, I advise students to learn the subject sectionwise and practice the exercises given.

Though these salient features are available in our Guide, I cannot negate the indispensable role of the teachers in assisting the student to understand the subject thoroughly.

I sincerely believe this guide satisfies the needs of the students and bolsters the teaching methodologies of the teachers.

I pray the almighty to bless the students for consummate success in their examinations.

Subash Raj, B.E., M.S.

- Publisher

**Sura Publications** 

All the Best

### **C**ONTENTS

### **VOLUME - I**

UNIT I	Nature of Physical	World and Measurement	
UNIT II	Kinematics		
UNIT III	Laws of motion		
UNIT IV	Work, Energy and	Power	
UNIT V	Motion of System	of Particles and Rigid Bodies	

### **VOLUME - II**

UNIT VI	: Gravitation	205 - 235
UNIT VII	: Properties of Matter	236 - 270
UNIT VIII	: Heat and Thermodynamics	271 - 321
UNIT IX	: Kinetic Theory of Gases	322 - 349
UNIT X	: Oscillations	350 - 387
UNIT XI	: Waves	
	Govt. Supply. Exam September Question Paper	425 - 426

### **TO ORDER WITH US**

#### SCHOOLS and TEACHERS

We are grateful for your support and patronage to **'SURA PUBLICATIONS'** Kindly prepare your order in your School letterhead and send it to us. For Orders contact: 80562 94222 / 80562 15222

#### **DIRECT DEPOSIT**

A/c Name	: Sura Publications	A/c Name : Sura Publications
Our A/c No.	: 36550290536	Our A/c No. : <b>21000210001240</b>
Bank Name	: STATE BANK OF INDIA	Bank Name : UCO BANK
Bank Branch	: PADI	Bank Branch : Anna Nagar West
IFSC	: SBIN0005083	IFSC : UCBA0002100
A/c Name	: Sura Publications	A/c Name : Sura Publications
A/c Name Our A/c No.	: Sura Publications : 6502699356	A/c Name : <b>Sura Publications</b> Our A/c No. : <b>1154135000017684</b>
A/c Name Our A/c No. Bank Name	: Sura Publications : 6502699356 : INDIAN BANK	A/c Name: Sura PublicationsOur A/c No.: 1154135000017684Bank Name: KVB BANK
A/c Name Our A/c No. Bank Name Bank Branch	: Sura Publications : 6502699356 : INDIAN BANK : ASIAD COLONY	A/c Name:Sura PublicationsOur A/c No.:1154135000017684Bank Name:KVB BANKBank Branch:Anna Nagar

After Deposit, please send challan and order to our address. email : orders@surabooks.com / Whatsapp : 81242 01000.

#### DEMAND DRAFT / CHEQUE

Please send Demand Draft / cheque in favour of **`SURA PUBLICATIONS'** payable at **Chennai**.

The Demand Draft / cheque should be sent with your order in School letterhead.

#### **STUDENTS**

Order via Money Order (M/O) to

### **SURA PUBLICATIONS**

1620, 'J' Block, 16th Main Road, Anna Nagar, Chennai - 600 040. Phones : 044-4862 9977, 044-486 27755 Mobile : 80562 94222 / 80562 15222 email : orders@surabooks.com Website : www.surabooks.com





(vi)





Sura's ■ XI Std - Physics IN Volume - I IN Unit 01 IN Nature of Physical World and Measurement

### FORMULAE TO REMEMBER

(1) Distance travelled by light in one year in vaccuum = [velocity of light  $\times$  1 year in seconds]

$$= 3 \times 10^{8} \times 365.25 \times 24 \times 60 \times 60$$
  
= 9.467 \times 10^{15} m

(2)  $\pi$  radian = 180°

(3) 1 radian = 
$$\frac{180^{\circ}}{\pi} = \frac{180^{\circ} \times 7}{22} = 57.27^{\circ}$$

(4) Also 1° (degree of arc) = 60' (minute of arc) and 1' (minute of arc) = 60'' (seconds of arc)

**Relations between radian, degree and minutes:** 

(5) 
$$1^{\circ} = \frac{\pi}{180}$$
 rad  $= 1.745 \times 10^{-2}$  rad

(6) 
$$1' = \frac{1^{\circ}}{60} = \frac{1.745 \times 10^{-2}}{60} = 2.908 \times 10^{-4} \text{ rad}$$
  
 $\approx 2.91 \times 10^{-4} \text{ rad}$ 

(7) 
$$1'' = \frac{1^{\circ}}{3600} = \frac{1.745 \times 10^{-2}}{3600} = 4.847 \times 10^{-6} \text{ rad}$$
  
 $\approx 4.85 \times 10^{-6} \text{ rad}$ 

- (8) Derived unit: Example : unit of speed =  $\frac{\text{unit of distance}}{\text{unit of time}} = \frac{\text{m}}{\text{s}} = \text{ms}^{-1}$
- (9) Absolute error  $a_m = \frac{a_1 + a_2 + a_3 + \dots + a_n}{n}$  or  $a_m = \frac{1}{n} \sum_{i=1}^{i=n} a_i$ ;  $a_m \to$  true value of measured quantity,

 $n \rightarrow$  number of values

(10) Mean Absolute error  $\Delta a_{\rm m} = \frac{1}{n} \sum_{i=1}^{n} |\Delta a_i|$ ;  $\Delta a_{\rm m} \rightarrow$  Mean absolute error,  $n \rightarrow$  number of values

(11) Relative error (or) Fractional error  $\Delta a = \frac{\Delta a_m}{a_m}$ ;  $a_m \rightarrow$  Mean value

(12) Percentage error, 
$$\Delta a = \frac{\Delta a_m}{a_m} \times 100\%$$

### SOME COMMON PRACTICAL UNITS

(1)	1 Fermi, 1 fm	=	10 <sup>-15</sup> m
(2)	1 Angstrom, 1Å	=	$10^{-10} \mathrm{m}$
(3)	1 nanometer, 1nm	=	10 <sup>-9</sup> m
(4)	1 micron (or) micro meter, 1µm	=	10 <sup>-6</sup> m
(5)	1 Light year	=	$9.467  imes 10^{15} \mathrm{m}$
(6)	1 Astronomical unit, 1 AU	=	$1.496 \times 10^{11} \text{ m}$
(7)	1 Parallactic second, 1 parsec	=	$3.08 \times 10^{16} \mathrm{m} = 3.26 \mathrm{light} \mathrm{year}$
(8)	1 CSL	=	1.4 times, the mass of the sun
(9)	1 shake	=	10 <sup>-8</sup> s (or) 10 nanoseconds

Prefixes for Powers of Ten						
Multiple	Prefix	Symbol	Sub multiple	Prefix	Symbol	
10 <sup>1</sup>	deca	da	10-1	deci	d	
10 <sup>2</sup>	hecto	h	10 <sup>-2</sup>	centi	с	
10 <sup>3</sup>	kilo	k	10 <sup>-3</sup>	milli	m	
10 <sup>6</sup>	mega	М	10 <sup>-6</sup>	micro	μ	
10 <sup>9</sup>	giga	G	10 <sup>-9</sup>	nano	n	
10 <sup>12</sup>	tera	Т	10 <sup>-12</sup>	pico	р	
10 <sup>15</sup>	peta	Р	10 <sup>-15</sup>	femto	f	
10 <sup>18</sup>	exa	Е	10 <sup>-18</sup>	atto	a	
10 <sup>21</sup>	zetta	Z	10 <sup>-21</sup>	zepto	Z	
10 <sup>24</sup>	yotta	Y	10 <sup>-24</sup>	yocto	у	

### IMPORTANT TERMS & DEFINITIONS

Science	:	Science is the systematic organization of knowledge gained through observation, experimentation and logical reasoning.
Physics	:	Physics is most basic science which deals with study of nature and natural phenomena.
Unification	:	Attempting to explain diverse physical phenomena with a few concepts and laws.
Reductionism	:	An attempt to explain a macroscopic system in terms of its microscopic constituents.

Sura's Nature of Physical World and Measurement

Technology	:	The application of the principles of physics, i.e. knowledge for practical purposes in various fields to invent and produce useful products or to solve problems.
Classical mechanics	:	The study of forces acting on bodies whether at rest or in motion
Thermodynamics	:	The study of the relationship between heat and other forms of energy
Optics	:	The study of light
Electricity and magnetism	:	The study of electricity and magnetism and their mutual relationship
Acoustics	:	The study of the production and propagation of sound waves
Astrophysics	:	The branch of physics which deals with the study of the physics of astronomical bodies
Relativity	:	One of the branches of theoretical physics which deals with the relationship between space, time and energy particularly with objects moving in different ways .
Quantum mechanics	:	The study of the discrete nature of phenomena at the atomic and subatomic levels
Atomic physics	:	The branch of physics which deals with the structure and properties of the atom
Nuclear physics	:	The branch of physics which deals with the structure, properties and reaction of the nuclei of atoms.
Condensed matter physics	:	The study of the properties of condensed materials (solids, liquids and those intermediate between them and dense gas). It branches into various sub-divisions including developing fields such as nano science, photonics etc. It covers the basics of materials science, which aims at developing new material with better properties for promising applications.
High energy physics	:	The study of the nature of the particles.
Range of time scales	:	Astronomical scales to microscopic scales, $10^{18}$ s to $10^{-22}$ s.
Range of masses	:	From heavenly bodies to electron, $10^{55}$ kg (mass of known observable universe) to $10^{-31}$ kg (mass of an electron) [the actual mass of an electron is $9.11 \times 10^{-31}$ Kg].
Measurement	:	The comparison of any physical quantity with its standard unit is known as measurement.
Physical Quantities	:	Quantities that can be measured, and in terms of which, laws of physics are described are called physical quantities.
Fundamental Quantities	:	Fundamental or base quantities are quantities which cannot be expressed in terms of any other physical quantities. These are length, mass, time, electric current, temperature, luminous intensity and amount of substance.
Derived Quantities	:	Quantities that can be expressed in terms of fundamental quantities are called derived quantities. Eg. area, volume, velocity, acceleration, force.
Unit of the quantity	:	An arbitrarily chosen standard of measurement of a quantity, which is accepted internationally is called unit of the quantity.

4

Fundamental or base units	:	The units in which the fundamental quantities are measured are called fundamental or base units.
Derived Unit	:	The units of measurement of all other physical quantities, which can be obtained by a suitable multiplication or division of powers of fundamental units, are called derived units.
System of units	:	A complete set of units which is used to measure all kinds of fundamental and derived quantities is called a system of units.
Radian (rad)	:	One radian is the angle subtended at the centre of a circle by an arc equal in length to the radius of the circle.
Steradian (sr)	:	One steradian is the solid angle subtended at the centre of a sphere, by that surface of the sphere, which is equal in area, to the square of radius of the sphere
macrocosm	:	Large objects like the galaxy, stars, Sun, Earth, Moon etc., and their distances constitute a macrocosm. It refers to a large world, in which both objects and distances are large.
Microcosm	:	Objects like molecules, atoms, proton, neutron, electron, bacteria etc., and their distances constitute microcosm, which means a small world in which both objects and distances are small-sized.
Parallax	:	The shift in the position of an object (say, a pen) when viewed with two eyes, keeping one eye closed at a time is known as Parallax.
Accuracy	:	Accuracy refers to how far we are from the true value.
Precision	:	Precision refers to how well we measure.
Systematic errors	:	Systematic errors are reproducible inaccuracies that are consistently in the same direction.
Least count error	:	Least count is the smallest value that can be measured by the measuring instrument, and the error due to this measurement is least count error.
Astronomical unit	:	It is the mean distance of the centre of sun from the centre of earth $1AU = 1.496 \times 10^{11}$ m
Light year	:	It is the distance travelled by light in vacuum in one year. 1 light year = $9.467 \times 10^{15}$ m.
Significant figures	:	The digits which tell us the number of units we are reasonably sure of having counted in making a measurement are called significant figures.
Error measurement	:	The uncertainity in the measurement of a physical quantity is called error. Error = True value – Measured value
Dimensions	:	Dimensions of a physical quantity are the powers to which the fundamental quantities must be raised.
Dimensional constant	:	Physical quantities which possess dimensions and have constant values are called dimensional constants.
Dimensionless constant	:	Quantities which have constant values and also have no dimensions are called dimensionless constants.
Principle of homogeneity of Dimensions	:	The principle of homogeneity of dimensions states that the dimensions of all the terms in a physical expression should be the same.

6

Sura's NI Std - Physics Wolume - I W Unit 01 W Nature of Physical World and Measurement

EVALUATION The dimensional formula of Planck's constant Ι. **MULTIPLE CHOICE QUESTIONS:** 8 h is [AMU, Main, JEE, NEET] 1. One of the combinations from the fundamental (a)  $[ML^2T^{-1}]$ (b)  $[ML^2T^{-3}]$ physical constants is  $\frac{hc}{G}$ . The unit of this (c)  $[MLT^{-1}]$ (d)  $[ML^{3}T^{-3}]$ expression is [Ans. (a)  $[ML^2T^{-1}]$ ] (a)  $kg^2$ (b)  $m^{3}$ (c)  $s^{-1}$ (d) m 9 The velocity of a particle v at an instant t is given by  $v = at + bt^2$ . The dimensions of b is [Ans. (a)  $kg^2$ ] (a) [L] (b)  $[LT^{-1}]$ 2. If the error in the measurement of radius is (c)  $[LT^{-2}]$ (d)  $[LT^{-3}]$ [Ans. (d) [LT<sup>-3</sup>]] 2%, then the error in the determination of **10.** The dimensional formula for gravitational volume of the sphere will be [Sep. - 2020] constant G is [Related to AIPMT 2004] (c) 4% (a) 8% (b) 2% (d) 6% (b)  $[M^{-1}L^3T^{-2}]$ (a)  $[ML^{3}T^{-2}]$ [Ans. (d) 6%] (c)  $[M^{-1}L^{-3}T^{-2}]$ (d)  $[ML^{-3}T^2]$ If the length and time period of an oscillating 3. [Ans. (b)  $[M^{-1}L^{3}T^{-2}]$ ] pendulum have errors of 1% and 3% **11.** The density of a material in CGS system of units respectively then the error in measurement of is 4 g cm<sup>-3</sup>. In a system of units in which unit of acceleration due to gravity is length is 10 cm and unit of mass is 100 g, then [Related to AMPMT 2008] [HY-2018] the value of density of material will be (a) 4% (b) 5% (c) 6% (d) 7% (a) 0.04 (b) 0.4 (c) 40 (d) 400 [Ans. (d) 7%] [Ans. (c) 40] **12.** If the force is proportional to square of The length of a body is measured as 3.51 m, 4. velocity, then the dimension of proportionality if the accuracy is 0.01m, then the percentage constant is [JEE-2000] [QY-2019] error in the measurement is (March 2020) (a)  $[MLT^0]$ (b)  $[MLT^{-1}]$ (a) 351% (b) 1% (c)  $[ML^{-2}T]$ (d)  $[ML^{-1}T^0]$ (c) 0.28% (d) 0.035% [Ans. (d)  $[ML^{-1}T^0]$ ] [Ans. (c) 0.28%] **13.** The dimension of  $(\mu_0 \varepsilon_0)^{-\frac{1}{2}}$  is [HY-2019] 5. Which of the following has the highest number [Main AIPMT 2011] of significant figures? (a) length (b) time (a)  $0.007 \text{ m}^2$ (b)  $2.64 \times 10^{24}$  kg (c) velocity (d) force (c)  $0.0006032 \text{ m}^2$ (d) 6.3200 J [Ans. (c) velocity] [Ans. (d) 6.3200 J] 14. Planck's constant (h), speed of light in vacuum (c) and Newton's gravitational constant (G) If  $\pi = 3.14$ , then the value of  $\pi^2$  is 6. are taken as three fundamental constants. [OY. - 2018; Jun.-2019] Which of the following combinations of these (a) 9.8596 (b) 9.860 has the dimension of length? **INEET 2016** (c) 9.86 (d) 9.9 [Ans. (c) 9.86] (phase II)] 7. Which of the following pairs of physical (a)  $\frac{\sqrt{hG}}{c^{\frac{3}{2}}}$  (b)  $\frac{\sqrt{hG}}{c^{\frac{5}{2}}}$  (c)  $\sqrt{\frac{hc}{G}}$  (d)  $\sqrt{\frac{Gc}{h^{\frac{3}{2}}}}$ quantities have same dimension? [Mar. - 2019] (a) force and power (b) torque and energy [Ans. (a)  $\frac{\sqrt{hG}}{\underline{3}}$ ] (c) torque and power (d) force and torque

#### Ph:9600175757 / 8124201000

#### orders@surabooks.com

[Ans. (b) torque and energy]

Sura's NI Std - Physics IN Unit 01 Nature of Physical World and Measurement

**15.** A length-scale (l) depends on the permittivity  $(\varepsilon)$  of a dielectric material, Boltzmann constant  $(k_B)$ , the absolute temperature (T), the number per unit volume (n) of certain charged particles, and the charge (q) carried by each of the particles. Which of the following expression for l is dimensionally correct? [JEE (advanced) 2016]

(a) 
$$l = \sqrt{\frac{nq^2}{\varepsilon k_B T}}$$
  
(b)  $l = \sqrt{\frac{\varepsilon k_B T}{nq^2}}$   
(c)  $l = \sqrt{\frac{q^2}{\varepsilon n^2 k_B T}}$   
(d)  $l = \sqrt{\frac{q^2}{\varepsilon n k_B T}}$   
[Ans. (b)  $l = \sqrt{\frac{\varepsilon k_B T}{nq^2}}$ ]

#### **II. SHORT ANSWER QUESTIONS.**

#### 1. Briefly explain the types of physical quantities.

- *Ans.* (i) Physical quantities are classified into two types. There are fundamental and derived quantities.
  - (ii) Fundamental or base quantities are quantities which cannot be expressed in terms of any other physical quantities. These are length, mass, time, electric current, temperature, luminous intensity and amount of substance.
  - (iii) Quantities that can be expressed in terms of fundamental quantities are called derived quantities. For example, area, volume, velocity, acceleration, force.

### 2. How will you measure the diameter of the Moon using parallax method? [HY-2018 & 19; QY-2019]

#### Ans. O - observation point on earth.

(i) In diagram, O is the observation point on the earth and d is the A diameter of moon. An astronomical telescope held at O is focussed on moon, the image is observed into moon of a circular disc.



(ii)  $\angle AOB = \theta$ 

S - average distance between moon and the surface of earth.

(iii) As 'S' is very large compared to the diameter, *d* of the moon, the diameter of the moon is considered as a circular arc of radius, S.

 $d = \mathbf{S} \times \mathbf{\theta}.$ 

Hence *d* can be calculated, when 'S' is known and  $\theta$  is measured.

## **3.** Write the rules for determining significant figures.

- Ans. (i) All non-zero digits are significant
  - (ii) All zeros between two non-zero digits are significant
  - (iii) All zeros to the right of a non-zero digit but to the left of a decimal point are significant.
  - (iv) a) The number without a decimal point, the terminal or trailing zero(s) are not significant.
    - b) All zeros are significant if they come from a measurement
  - (v) If the number is less than 1, the zero (s) on the right of the decimal point but to left of the first non-zero digit are not significant.
  - (vi) All zeros to the right of a decimal point and to the right of non-zero digit are significant.
  - (vii) The number of significant figures does not depend on the system of units used.

## 4. What are the limitations of dimensional analysis? [Govt. MQP-2018; HY-2018; Jun.-2019]

#### Ans. Limitations of Dimensional analysis:

- (i) This method gives no information about the dimensionless constants in the formula like 1, 2, ..... $\pi$ ,e, etc.
- (ii) This method cannot decide whether the given quantity is a vector or a scalar.
- (iii) This method is not suitable to derive relations involving trigonometric, exponential and logarithmic functions.
- (iv) It cannot be applied to an equation involving more than three physical quantities.
- (v) It can only check on whether a physical relation is dimensionally correct but not the correctness of the relation. For example using dimensional analysis,  $s = ut + 1/3 at^2$  is dimensionally correct whereas the correct relation is  $s = ut + 1/2 at^2$ .
- 5. Define precision and accuracy. Explain with one example.
- *Ans.* **Precision:** The closeness of two or more measurements to each other is known as precision.

8

Sura's Nature of Physical World and Measurement

**Accuracy:** The closeness of a measured value to the actual value of the object being measured is called accuracy.

**Example:** Suppose a man's true height is exactly 5'9''. When it is measured with a yardstick, the value is 5'0''. Hence measurement is not accurate. When height is measured with a laser yardstick, the value is 5'9'' then measurement is accurate.

If the height is measured consistently as 5'0'' with a yardstick, then measurements are precise.

#### **III.** LONG ANSWER QUESTIONS

- 1. (i) Explain the use of screw gauge and vernier caliper in measuring smaller distances.
  - (ii) Write a note on triangulation method and radar method to measure larger distances. [Govt.MQP-2018]

#### Ans. Measurement of small distances:

- (i) The screw gauge is an instrument used for measuring accurately the dimensions of objects up to a maximum of about 50 mm.
  - (a) It is used for measuring external dimensions. i.e. diameter.
  - (b) The least count of the screw gauge is 0.01 mm.

A vernier caliper is a versatile instrument for measuring the dimensions of an object namely diameter of a hole, or a depth of a hole. i.e. internal & external dimensions. The least count of vernier caliper is 0.1 mm

(ii) For measuring larger distances such as the height of a tree, distance of the Moon or a planet from the Earth, triangulation method, parallax method and radar method are used.

### Triangulation method for the height of an accessible object : (March 2020)

(a) Let AB = h be the height of the tree or tower to be measured. Let C be the point of observation at distance *x* from B. Place a range finder c at C and measure the angle of elevation,



 $\angle ACB = \theta \text{ as shown in Figure.}$ (b) From right-angled triangle ABC,  $\tan \theta = \frac{AB}{BC} = \frac{A}{2}$ (or) height  $h = x \tan \theta$ .

#### orders@surabooks.com

(c) Knowing the distance *x*, the height h can be determined.

#### **RADAR method**

- (i) The word RADAR stands for radio detection and ranging.
- (ii) A radar can be used to measure accurately the distance of a nearby planet such as Mars. In this method, radio waves are sent from transmitters which, after reflection from the planet, are detected by the receiver.
- (iii) By measuring, the time interval (*t*) between the instants the radio waves are sent and received, the distance of the planet can be determined as

 $Distance(d) = Speed of radio waves \times time$ 

taken 
$$d = \frac{v \times t}{2}$$

- (iv) where v is the speed of the radio wave. As the time taken (t) is for the distance covered during the forward and backward path of the radio waves, it is divided by 2 to get the actual distance of the object.
- (v) This method can also be used to determine the height, at which an aeroplane flies from the ground.

#### 2. Explain in detail the various types of errors.

[Mar., QY-2019]

#### Ans. Types of errors :

- (a) Systematic error (b) Random error
- (c) Gross error
- (a) Systematic errors : They are reproducible inaccuracies that are consistently in the same direction.

#### It is classified as follows :

- (1) **Instrumental errors :** It arises when an instrument is not calibrated properly at the time of manufacturing. It can be corrected by choosing accurate instruments.
- (2) Imperfections in experimental technique or procedure: It is due to the limitation in the experimental arrangement. To overcome this, necessary and proper correction is to be applied.
- (3) **Personal errors :** These errors are due to individuals performing the experiment, may be due to incorrect initial setting up to the experiment or carelessness of the individual making the observation due to improper precautions.

Sura's XI Std - Physics I Unit 01 I Nature of Physical World and Measurement

- (4) Errors due to external causes : The change in the external conditions during an experiment can cause error in measurement. For example, changes in temperature, humidity, or pressure during measurements may affect the result of the measurement.
- (5) Least count error : Least count is the smallest value that can be measured by the measuring instrument, and the error due to this measurement is least count error.

#### (b) Random error :

- (1) It arises due to random and unpredictable variations in experimental conditions like pressure, temperature, voltage supply, etc.
- (2) It also arises due to personal errors by the observer. It is sometimes called 'chance errors'.
- (3) It can be minimised by repeating the observations a large number of times and taking the arithmetic mean of all the observations.

#### (c) Gross error :

- (1) The error caused due to the shear carelessness of an observer is called gross error.
- (2) It can be minimized only when an observer is careful and mentally alert.
- **3.** What do you mean by propagation of errors? Explain the propagation of errors in addition and multiplication. (March 2020)
- **Ans.** A number of measured quantities may be involved in the final calculation of an experiment. Different types of instruments might have been used for taking readings. Then we may have to look at the errors in measuring various quantities, collectively. The error in the final result depends on
  - (i) The errors in the individual measurements
  - (ii) On the nature of mathematical operations performed to get the final result. So we should know the rules to combine the errors.

These types of combination of errors are known as Propagation of errors.

(i) Error in the sum of two quantities :

Let  $\Delta A$  and  $\Delta B$  be the absolute errors in the two quantities A and B respectively. Then, Measured value of  $A = A \pm \Delta A$  Measured value of  $B = B \pm \Delta B$ Consider the sum, Z = A + BThe error  $\Delta Z$  in Z is then given by  $Z \pm \Delta Z = (A \pm \Delta A) + (B \pm \Delta B)$  $= (A + B) \pm (\Delta A + \Delta B)$  $= Z \pm (\Delta A + \Delta B)$ (or)  $\Delta Z = \Delta A + \Delta B$ 

The maximum possible error in the sum of two quantities is equal to the sum of the absolute errors in the individual quantities.

(ii) Error in the difference of two quantities : Let  $\Delta A$  and  $\Delta B$  be the absolute errors in the two quantities, A and B, respectively. Then,

Measured value of  $A = A \pm \Delta A$ 

Measured value of  $B = B \pm \Delta B$ 

Consider the difference, Z = A - B

The error 
$$\Delta Z$$
 in Z is then given by

$$Z \pm \Delta Z = (A \pm \Delta A) - (B \pm \Delta B)$$
$$= (A - B) \pm \Delta A + \Delta B$$

$$= Z \pm \Delta A + \Delta B$$

(or)  $\Delta Z = \Delta A + \Delta B$ 

The maximum error in difference of two quantities is equal to the sum of the absolute errors in the individual quantities.

#### (iii) Error in the product of two quantities

Let  $\Delta A$  and  $\Delta B$  be the absolute errors in the two quantities, A and B, respectively. Consider the product Z = AB

=  $(AB) \pm (A \Delta B) \pm (B \Delta A) \pm (\Delta A \cdot \Delta B)$ Dividing L.H.S by Z and R.H.S by AB, we get,

 $1 \pm \frac{\Delta Z}{Z} = 1 \pm \frac{\Delta B}{B} \pm \frac{\Delta A}{A} \pm \frac{\Delta A}{A} \cdot \frac{\Delta B}{B}$ As  $\Delta A / A$ ,  $\Delta B / B$  are both small quantities, their product term  $\frac{\Delta A}{A} \cdot \frac{\Delta B}{B}$  can be neglected.

The maximum fractional error in Z is

$$\frac{\Delta Z}{Z} = \pm \left(\frac{\Delta A}{A} + \frac{\Delta B}{B}\right)$$

The maximum fractional error in the product of two quantities is equal to the sum of the fractional errors in the individual quantities.

Sura's XI Std - Physics W Volume - I W Unit 01 W Nature of Physical World and Measurement

- 4. Write short notes on the following.
  - (a) Unit
  - (b) Rounding off
  - (c) Dimensionless quantities
- Ans. (a) Unit : An arbitrarily chosen standard of measurement of a quantity, which is accepted internationally is called unit of the quantity. The units in which the fundamental quantities are measured are called fundamental or base units and the units of measurement of all other physical quantities, which can be obtained by a suitable multiplication or division of powers of fundamental units are called derived units.
  - (b) Rounding off : The result given by a calculator has too many figures. In no case should the result have more significant figures than the figures involved in the data used for calculation. The result of calculation with numbers containing more than one uncertain digit should be rounded off.
  - (c) Dimensionless quantities :
    - (i) Physical quantities which have no dimensions, but have variable values are called dimensionless variables. Examples are Gravitational constant, Planck's constant etc.
    - (ii) Quantities which have constant values and also have no dimensions are called dimensionless constants. Examples are π, e, numbers, etc.
- 5. Explain the principle of homogeneity of dimensions. What are its uses? Give example.

[HY-2018 & 19]

#### Ans. Principle of homogeneity of dimensions :

It states that the dimensions of all the terms in a physical expression should be the same.

```
For example
```

 $v^2 = u^2 + 2as$ 

 $[LT^{-1}]^2 = [LT^{-2}]^2 + 2 [LT^{-2}] [L]$  $[L^2T^{-2}] = [L^2T^{-2}] + 2[L^2T^{-2}]$ 

#### This method is used to

- (i) Convert a physical quantity from one system of units to another.
- (ii) Check the dimensional correctness of a given physical equation.
- (iii) Establish relations among various physical quantities.

To convert a physical quantity from one system of units to another

- (i) This is based on the fact that the product of the numerical values (n) and its corresponding unit (u) is a constant. i.e, n[u] = constant (or)  $n_1[u_1] = n_2[u_2]$ .
- (ii) Consider a physical quantity which has dimension 'a' in mass, 'b' in length and 'c' in time. If the fundamental units in one system are  $M_1$ ,  $L_1$  and  $T_1$  and the other system are  $M_2$ ,  $L_2$  and  $T_2$  respectively, then we can write,  $n_1 [M_1^{a} L_1^{b} T_1^{c}] = n_2 [M_2^{a} L_2^{b} T_2^{c}]$
- (iii) We have thus converted the numerical value of physical quantity from one system of units into the other system.

#### **IV. NUMERICAL PROBLEMS.**

1. In a submarine equipped with sonar, the time delay between the generation of a pulse and its echo after reflection from an enemy submarine is observed to be 80 s. If the speed of sound in water is 1460 ms<sup>-1</sup>. What is the distance of enemy submarine?

**Ans.** The speed of sound in water  $v = 1460 \text{ ms}^{-1}$ 

Time taken by the pulse for to and fro :

$$t = \frac{T}{2} = \frac{80s}{2} = 40s$$
Formula :  $v = \frac{d}{t}$ 

$$\therefore d = v \times \frac{T}{2} = 1460 \times 40$$

= 58400 m or 58.40 km.

2. The radius of the circle is 3.12 m. Calculate the area of the circle with regard to significant figures. [QY-2019]

**Ans.** Radius of the circle r = 3.12 m

Area of the circle A = ?

A =  $\pi r^2 = 3.14 \times 3.12 \times 3.12 = 30.566016$ 

According to the rule of significant fig,

 $A = 30.6 \text{ m}^2$  [Given data has three sig. fig.]

Sura's IXI Std - Physics IN Unit 01 IN Nature of Physical World and Measurement

- Assuming that the frequency  $\gamma$  of a vibrating 3. string may depend upon (i) applied force (F) (ii) length (l) (iii) mass per unit length (m), prove
  - that  $\gamma \propto \frac{1}{I} \sqrt{\frac{F}{m}}$  using dimensional analysis. (related to JIPMER 2001)

**Ans.** 
$$\gamma \propto l^a F^b m^c$$

$$\gamma = K l^a F^b m^c$$

K - dimensionless constant of proportionality a, b, c - powers of l, F, m Dimensional Formula of  $F = [MLT^{-2}]$ 

Dimensional Formula of linear density

$$m = \frac{\text{mass}}{\text{length}} = \frac{\lfloor \mathbf{M} \rfloor}{\lfloor \mathbf{L} \rfloor} = [\mathbf{M}^1 \ \mathbf{L}^{-1}]$$

writing dimensions of  $\gamma = K l^a F^b m^c$ 

 $[M^{0}L^{0}T^{-1}] = [L]^{a} [M^{1}L^{1}T^{-2}]^{b} [ML^{-1}]^{c}$ =  $L^{a}M^{b}L^{b}T^{-2b}M^{c}L = [M]^{b+c}[L]^{a+b-c}[T]^{-2b}$  $M^{0}L^{0}T^{-1} = [M]^{b+c} [L]^{a+b-c} [T]^{-2b}$ 

Applying the principle of homogeneity of dimension

$$b + c = 0 \qquad \dots(1)$$

$$a + b - c = 0 \qquad \dots(2)$$

$$-2b = -1 \text{ or } \boxed{b = +\frac{1}{2}}$$
From  $\bigcirc c = -b = -\frac{1}{2} \qquad \therefore \boxed{c = -\frac{1}{2}}$ 
From  $\oslash a + \frac{1}{2} - (-\frac{1}{2}) = 0 \qquad a + \frac{1}{2} + \frac{1}{2} = 0$ 

$$\therefore \boxed{a = -1}$$
Substituting the values  $a, b, c$  in  $\because K = 1$ 

$$\gamma = K \ l^{-1} F^{\frac{1}{2}} \ m^{-\frac{1}{2}}$$

$$\gamma = \frac{1}{2} \sqrt{\frac{F}{2}}$$

Ans. Distance of Jupiter from the earth,

orders@surabooks.com

 $D = 824.7 \times 10^6 \text{ km}$  $D = 824.7 \times 10^9 \text{ m}$ Angular diameter  $\theta = 35.72$ "  $= 35.72 \times (4.85 \times 10^{-6} \text{ rad})$  $[::1'' = 4.85 \times 10^{-6} \text{ rad}]$ 

#### Diameter of Jupiter, d = ?

$$\theta = \frac{d}{D}$$
  

$$\therefore d = \theta.D$$
  

$$= (824.7 \times 10^{9}) \times 35.72 \times 4.85 \times 10^{-6}$$
  

$$= 142872.677 \times 10^{3} \text{ m}$$
  

$$= 142872.677 \times 10^{3} \times 10^{-3} \text{ km}$$
  

$$d = 1.428 \times 10^{5} \text{ km}.$$

The measurement value of length of a simple **5**. pendulum is 20 cm known with 2 mm accuracy. The time for 50 oscillations was measured to be 40 s within 1s resolution. Calculate the percentage accuracy in the determination of acceleration due to gravity 'g' from the above measurement.

Ans. The errors in both l & T are least count errors.

$$T = 2\pi \sqrt{\frac{l}{g}} \Rightarrow T^2 = 4\pi^2 \cdot \frac{l}{g}$$
$$\therefore g = -4\pi^2 \frac{l}{T^2}$$

The errors in both *l* & T are least count errors  $\frac{\Delta g}{g} = \frac{\Delta l}{l} + \frac{2\Delta T}{T}$ Length of simple pendulum l = 20 cm

accuracy  $\Delta l = 2$ mm = 0.2 cm

Time for 50 oscillations T = 40 sresolution  $\Delta T$ = 1 s $\therefore \frac{\Delta g}{g} = \left(\frac{0.2}{20}\right) + 2\left(\frac{1}{40}\right) = \frac{0.2}{20} + \frac{2}{40} = \frac{1.2}{20}$ 

Percentage error

to

$$\frac{\Delta g}{g} \times 100 = \frac{1.2}{20} \times 100 = \pm 6\%$$

% accuracy in g = 6%.

**Government Exam Question & Answers** 

I. **MULTIPLE CHOICE OUESTIONS :** — 1 Mark —

A substance whose mass is 4.27 g occupies 1. 1. 3 cm<sup>3</sup>. The number of significant figure in density is [Govt. M.O.P - 2018] (a) 1 (b) 2 (c) 3 (d) 4 [Ans. (d) 4]

#### Ph:9600175757 / 8124201000

11

12

2.

3.

4.

Sura's NI Std - Physics Wolume - I W Unit 01 W Nature of Physical World and Measurement

**Triple point of water is :** 3. *[OY. - 2018]* (a) 273.16 k (b) 237.16 c (c) 273.16 c (d) 0 k [Ans. (a) 273.16k] Mass, temperature, electric current are [QY. - 2018] (a) fundamental quantities (b) scalar quantities (c) vector quantities (d) both a and b [Ans. (d) both a and b] The significant figure of the number 0.003401 is: [QY. - 2019] 4. (a) 6 (b) 3 (c) 5 (d) 4 [Ans. (d) 4] The amplitude and time period of a simple **5**. 5. pendulum bob are 0.05 m and 2 s respectively. Then the maximum velocity of the bob is : [Mar. - 2019] a)  $0.157 \text{ ms}^{-1}$ b)  $0.257 \text{ ms}^{-1}$ c)  $0.10 \text{ ms}^{-1}$ d)  $0.025 \text{ ms}^{-1}$ [Ans. (a) 0.157 ms<sup>-1</sup>] **II. VERY SHORT ANSWER OUESTIONS : —2 Marks** — **6**. Check the dimensional correctness for the 1. given equations. [First Mid-2018] (b)  $s = ut + \frac{1}{2}at$ (a) V = u + atAns. (a) v = u + at Apply dimensional formula on both sides  $[LT^{-1}] = [LT^{-1}] + [LT^{-2}] [T]$  $[LT^{-1}] = [LT^{-1}] + [LT^{-1}]$ 1. (Quantities of same dimension only can be added) Since dimensions on both sides are same, the given equation is dimensionally correct. (b)  $[L] = [LT^{-1}] [T] + [LT^{-2}] [T^2]$  $= [LT^{-1+1}] + [LT^{-2+2}]$ [L] $[L] = [LT^0] + [LT^0]$ ::[L] = [L] + [L]Since dimensions on both sides are same, the given equation is dimensionally correct. Round off to required significant figures [First Mid-2018] (a) 3.1 + 1.780 + 2.046(b) 12.637 - 2.42(c)  $1.21 \times 36.72$ (d)  $36.72 \div 1.2$ 

### orders@surabooks.com

(b) 10.22

(c) 44.4

(d) 31

**Ans.** (a) 6.96

#### What are random errors? How to minimise it? [First Mid-2018]

- Ans. (i) Random errors may arise due to random and unpredictable variations in experimental conditions like pressure, temperature, voltage supply, etc. Random errors are sometimes called "chance error".
  - (ii) Random errors can be evaluated through statistical analysis and can be reduced by averaging over a large number of observations.
- Write down the number of significant figures in the following: (i) 0.007 (ii) 400.[Govt. MOP-2018]
- Ans. (i) One (ii) One
- What are the advantages of SI system?

[**OY-2018**]

[QY-2018]

- **Ans.** (i) It is a rational system, in which only one unit is used for one physical quantity.
  - (ii) It is a coherent system, which means all the derived units can be easily obtained form basic and supplementary units.
  - (iii) It is a metric system which means that multiples and submultiples can be expressed as powers of 10.
  - What is fractional error?
- Ans. Ratio of mean absolute error to the mean value (or) relative error or fractional error = mean absolute error / mean value.

#### **III. SHORT ANSWER QUESTIONS :**

**—3 Marks** —

- Write a note on radar method to measure larger distances. [First Mid-2018]
- The word RADAR stands for radio detection Ans. (i) and ranging.
  - (ii) A radar can be used to measure accurately the distance of a nearby planet such as Mars. In this method, radio waves are sent from transmitters which, after reflection from the planet, are detected by the receiver.
  - (iii) By measuring, the time interval (t) between the instants the radio waves are sent and received, the distance of the planet can be determined as

 $Distance(d) = Speed of radio waves \times time$ taken  $d = \frac{v \times t}{2}$ 

where *v* is the speed of the radio wave. As the time taken (t) is for the distance covered

Sura's XI Std - Physics I Vnit 01 Nature of Physical World and Measurement

during the forward and backward path of the radio waves, it is divided by 2 to get the actual distance of the object.

- (iv) This method can also be used to determine the height, at which an aeroplane flies from the ground.
- 2. The voltage across a wire is  $(100 \pm 5)$  v and the current passing through it is  $(10 \pm 0.2)$  A. Find the resistance of the wire. [First Mid-2018]
- Ans.

Voltage V =  $(100 \pm 5)$ V Current I =  $(10 \pm 0.2)$ A

Resistance R = ?

Then resistance R is given by Ohm's law

$$R = \frac{V}{I} = \frac{100}{10} = 10 \Omega$$
$$\frac{\Delta R}{R} = \left(\frac{\Delta V}{V} + \frac{\Delta I}{I}\right)$$

$$\Delta \mathbf{R} = \left(\frac{\Delta \mathbf{V}}{\mathbf{V}} + \frac{\Delta \mathbf{I}}{\mathbf{I}}\right) \mathbf{R} = \left(\frac{5}{100} + \frac{0.2}{10}\right) \times 10$$
  
= (0.05 + 0.02) × 10 = 0.07 × 10 = 0.7

The resistance  $R = (10 \pm 0.7)\Omega$ .

**3.** Check the correctness of the equation  $E = mc^2$  using dimensional analysis method.

[Govt. MQP-2018; Jun.-2019]

- Ans. Consider the equation,  $E = mc^2$ Apply dimensional formula on both sides  $ML^2T^{-2} = [M] \cdot [LT^{-1}]^2$  $ML^2T^{-2} = [M] \cdot [L^2T^{-2}]$ The equation is dimensionally correct.
- 4. Two resistances  $R_1 = (100 \pm 3)\Omega$  and  $R_2 = (150\pm 2)\Omega$ are connected in series. What is their equivalent resistance? [Govt. MOP-2018]
- Ans.  $R_1 = 100 \pm 3 \Omega$ ;  $R_2 = 150 \pm 2\Omega$ Equivalent resistance R = ?Equivalent resistance  $R = R_1 + R_2$  $= (100 \pm 3) + (150 \pm 2) = (100 + 150) \pm (3 + 2)$  $R = (250 \pm 5) \Omega$

5. Find the dimensional formula of hC/G. [QY-2018]Ans. The dimensional formula for

planck's constant 
$$h - [ML^2T^{-1}]$$
  
 $c - [LT^{-1}]$   
 $G - [M^{-1}L^2T^{-2}]$   
 $\frac{hc}{G} = \frac{[ML^2T^{-1}][LT^{-1}]}{[M^{-1}L^3T^{-2}]} = [M^2]$ 

#### IV. LONG ANSWER QUESTIONS :

—5 Marks —

1. The force F acting on a body moving in a circular path depends on mass of the body(m) velocity(v) and radius (r) of the circular path. Obtain the expression for the force by dimensional analysis method. (k = 1). [First Mid-2018; Mar.-2019]

**Ans.** 
$$\mathbf{F} \propto m^a v^b r^c$$
;  $\mathbf{F} = k m^a v^b r^c$ 

where k is a dimensionless constant of proportionality. Rewriting the above equation in terms of dimensions and taking k = 1, we have

$$\begin{split} [MLT^{-2}] &= [M]^a \ [LT^{-1}]^b \ [L]^c = [M^a \ L^b \ T^{-b} \ L^c] \\ [MLT^{-2}] &= [M]^a \ [L^{b+c}] \ [T^{-b}] \end{split}$$

Comparing the powers of M, L and T on both sides a = 1; b + c = 1 - b = -2; 2 + c = 1, b = 2

$$a = 1, b = 2$$
 and  $c = -1$ 

From the above equation we get  $F = m^a v^b r^c$ 

$$F = m^1 v^2 r^{-1}$$
 or  $F = \frac{mv^2}{r}$ 

2. Obtain an expression for the time period T of a simple pendulum. [The time period T depend upon (i) mass *m* of the bob, (ii) length *l* of the pendulum and (iii) acceleration due to gravity *g* at the place where pendulum is suspended. Assume the constant,  $k = 2\pi$ ].

Ans.

[Govt. MQP-2018; QY, HY-2019]

$$T \propto m^a l^b g^c$$
;  $T = k. m^a l^b g^c$ 

Here k is the dimensionless constant.

Rewriting the above equation with dimensions.

$$[T^1] = [M^a] [L^b] [LT^{-2}]^c$$

 $[\mathbf{M}^{0}\mathbf{L}^{0}\mathbf{T}^{1}] = [\mathbf{M}^{a} \mathbf{L}^{b+c} \mathbf{T}^{-2c}]$ 

Comparing the powers of M, L and T on both sides, a = 0, b + c = 0, -2c = 1

Solving for *a*, *b* and *c*, we get a = 0, b = 1/2, and c = -1/2

From the above equation  $T = k. m^0 l^{1/2} g^{-1/2}$ 

$$\mathbf{T} = k \left(\frac{l}{g}\right)^{1/2} = k \sqrt{\frac{l}{g}}$$

Experimentally  $k = 2\pi$ , hence T =  $2\pi \sqrt{\frac{l}{g}}$ 

#### orders@surabooks.com

14

Sura's XI Std - Physics W Volume - I W Unit 01 W Nature of Physical World and Measurement

**3**. In a series of successive measurements in an experiment, the readings of the period of oscillation of a simple pendulum were found to be 2.63 s, 2.56 s, 2.42 s, 2.71 s and 2.80 s. Calculate (i) the mean value of the period of oscillation (ii) the absolute error in each measurement (iii) the mean absolute error (iv) the relative error (v) the percentage error. Express the results in proper form. [Govt. MQP-2018]

#### Ans.

$$t_{1} = 2.63s, t_{2} = 2.56s, t_{3} = 2.42 s,$$

$$t_{4} = 2.71s, t_{5} = 2.80 s$$
(i)  $T_{m} = \frac{t_{1} + t_{2} + t_{3} + t_{4} + t_{5}}{5}$ 

$$= \frac{2.63 + 2.56 + 2.42 + 2.71 + 2.80}{5}$$
 $T_{m} = \frac{13.12}{5} = 2.624 s$ 
 $T_{m} = 2.62 s$  (Rounded off to 2<sup>nd</sup> decimal place)

- (ii) Absolute error  $\Delta T = T_m t$   $\Delta T_1 = 2.62 - 2.63 = -0.01$   $\Delta T_2 = 2.62 - 2.56 = +0.06s$   $\Delta T_3 = 2.62 - 2.42 = +0.20s$   $\Delta T_4 = 2.62 - 2.71 = -0.09s$  $\Delta T_5 = 2.62 - 2.80 = -0.18s$
- (iii) Mean absolute error =  $\frac{\sum |\Delta T_t|}{n}$

$$\Delta T_{\rm m} = + \frac{0.01 + 0.06 + 0.20 + 0.09 + 0.18}{5}$$
$$\Delta T_{\rm m} = \frac{0.54}{5} = 0.108 \text{s} = 0.11 \text{s}$$

(Rounded off to 2<sup>nd</sup> decimal place) (iv) Relative error:

$$S_{\rm T} = \frac{\Delta T_{\rm m}}{T_{\rm m}} = \frac{0.11}{2.62} = 0.0419 = 0.04$$

- (v) Percentage error in T =  $0.04 \times 100\% = 4\%$ (vi) Time period of simple pendulum = T =  $(2.62 \pm 0.11)$ s
- 4. What are the applications of dimensional analysis. Verify  $S = ut + 1/2 at^2$  by dimensional analysis. [Govt.MQP-2018; QY-2018; Sep.-2020]
- Ans. (i) To convert a physical quantity from one system of units to another.

#### orders@surabooks.com

This is based on the fact that the product of the numerical values (*n*) and its corresponding unit (u) is a constant. i.e, n[u] = constant (or)  $n_1[u_1] = n_2[u_2]$ .

Consider a physical quantity which has dimension 'a' in mass, 'b' in length and 'c' in time. If the fundamental units in one system are  $M_1$ ,  $L_1$  and  $T_1$  and the other system are  $M_2$ ,  $L_2$  and  $T_2$ , respectively, then we can write,  $n_1 [M_1^{\ a} L_1^{\ b} T_1^{\ c}] = n_2 [M_2^{\ a} L_2^{\ b} T_2^{\ c}].$ 

We have thus converted the numerical value of physical quantity from one system of units into the other system.

(ii) To check the dimensional correctness of a given physical equation.

Let us take the equation of motion

v = u + at

Apply dimensional formula on both sides

 $[LT^{-1}] = [LT^{-1}] + [LT^{-2}] [T]$ 

 $[LT^{-1}] = [LT^{-1}] + [LT^{-1}]$ 

(Quantities of same dimension only can be added)

We see that the dimensions of both sides are same. Hence the equation is dimensionally correct.

(iii) To establish the relation among various physical quantities.

If the physical quantity Q depends upon the quantities  $Q_1$ ,  $Q_2$  and  $Q_3$ , i.e., Q is proportional to  $Q_1$ ,  $Q_2$  and  $Q_3$ .

Then  $Q \propto Q_1^a Q_2^{\ b} Q_3^{\ c}$ ;  $Q = k Q_1^a Q_2^{\ b} Q_3^{\ c}$ where *k* is a dimensionless constant. When the dimensional formula of Q, Q<sub>1</sub>, Q<sub>2</sub> and Q<sub>3</sub> are substituted, then according to the principle of homogeneity, the powers of M, L, T are made equal on both sides of the equation. From this, we get the values of *a*, *b*, *c*.

$$s = ut + \frac{1}{2}at^2$$

Substituting dimensions

 $[L] = [LT^{-1}][T] + [LT^{-2}][T^2]$ 

[L] = [L] + [L]

The equation is dimensionally correct.

Sura's XI Std - Physics I Unit 01 I Nature of Physical World and Measurement

**5.** Express 76 cm of mercury pressure in terms of Nm<sup>-2</sup> using the method of dimensions. [Sep.-2020]

#### Ans. Solution :

In cgs system 76 cm of mercury pressure

 $= 76 \times 13.6 \times 980 \text{ dyne cm}^{-2}$ 

The dimensional formula of pressure P is  $[ML^{-1}T^{-2}]$ 

 $P_{1}[M_{1}^{a}L_{1}^{b}T_{1}^{c}] = P_{2}[M_{2}^{a}L_{2}^{b}T_{2}^{c}]$ 

We have  $P_2 = P$ 

$$\mathbf{P}_{1} \left[ \frac{\mathbf{M}_{1}}{\mathbf{M}_{2}} \right]^{a} \left[ \frac{\mathbf{L}_{1}}{\mathbf{L}_{2}} \right]^{b} \left[ \frac{\mathbf{T}_{1}}{\mathbf{T}_{2}} \right]^{b} \mathbf{T}_{1}$$

$$M_{1} = 1g, M_{2} = 1 \text{ kg}$$

$$L_{1} = 1 \text{ cm}, L_{2} = 1 \text{ m}$$

$$T_{1} = 1s, T_{2} = 1 \text{ s}$$
As  $a = 1, b = -1$ , and  $c = -2$   
Then
$$P_{2} = 76 \times 13.6 \times 980 \left[\frac{1g}{1\text{ kg}}\right]^{1} \left[\frac{1\text{ cm}}{1\text{ m}}\right]^{-1} \left[\frac{1\text{ s}}{1\text{ s}}\right]^{-1}$$

$$= 76 \times 13.6 \times 980 \left[\frac{10^{-3}\text{ kg}}{1\text{ kg}}\right]^{1} \left[\frac{10^{-2}\text{ m}}{1\text{ m}}\right]^{-1}$$

$$= 76 \times 13.6 \times 980 \times [10^{-3}] \times 10^{-3}$$
  
P<sub>2</sub> = 1.01 × 10<sup>5</sup> Nm<sup>-2</sup>.

### Additional Questions

I. MULTIPLE CHOICE QUESTIONS :

— 1 Mark =

#### A. CHOOSE THE BEST ANSWER :

#### **1.** The word scientia is meaning to

- (a) exact
- (c) control
- (d) implement

(b) to know

#### [Ans. (b) to know]

- 2. Astronomical Scale is dealt with the Physics.
  - (a) Mesoscopic (b) Microscopic
  - (c) Macrospic (d) None
    - [Ans. (c) Macrospic]
- **3.** Microscopic group of Physics dealt with the study of \_\_\_\_\_.
  - (a) classical physics (b) statistical mechanics

(c) fluid mechanics

(d) quantum physics

#### [Ans. (d) quantum physics]

### 4. Which deals with the study of materials of an intermediate length scale.

- (a) Mesoscopic physics
- (b) Macroscopic physics
- (c) Microscopic physics
- (d) All the above [Ans. (a) Macroscopic physics]

- 5. What is the range of astronomical time scales to microscopic scales?
  - (a)  $10^{15}$  s to  $10^{-15}$  s (b)  $10^{9}$  s to  $10^{-18}$  s (c)  $10^{18}$  to  $10^{-22}$  s (d)  $10^{11}$  s to  $10^{-16}$  s
    - (d)  $10^{-1}$  s to  $10^{-1}$  s

[Ans. (c) 10<sup>18</sup> to 10<sup>-22</sup>s]

#### 6. The law of electricity and magnetism is used to

- (a) Wireless communication
- (b) Nuclear reactor (c) Steam engine
- (d) Aeroplane[Ans. (a) Wireless communication]

#### 7. Match the following.

	(1)	Steam engine		(a)	Bernoulli's principle
	(2)	Nuclear reactor	•	(b)	Laws of thermodynamics
	(3)	Product ultra hi magnet	tion of gh ic fields	(c)	Controlled nuclear fission
		)		1	
	(4)	Aeropla	ane	( <b>d</b> )	Super conductivity
	( <b>4</b> )	Aeropla	(3)	( <b>d</b> ) (4)	Super conductivity
(a)	( <b>4</b> ) (1) ) b	Aeropla ) (2) c	(3) a	( <b>d</b> ) (4) d	Super conductivity
(a) (b)	( <b>4</b> ) (1) ) b ) d	Aeropla ) (2) c a	ane (3) a b	(d) (4) d c	Super conductivity
(a) (b) (c)	(4) (1) ) b ) d ) c	Aeropla ) (2) c a d	ane (3) a b a	(d) (4) d c b	Super conductivity

15

 $\frac{1s}{1s}$ 

16

Sura's NI Std - Physics Wolume - I W Unit 01 W Nature of Physical World and Measurement

- Match the following fundamental forces with || 17. The acceleration of 20 m/s<sup>2</sup> in km/h<sup>2</sup> is 8. respect to relative strengths.
  - Gravitational force 1 (1) (a)
    - (2) **Electromagnetic force (b)** Weak nuclear force (3) (c)
    - (4) Strong nuclear force
  - (1)(2) (3) (4)d (a) a h с
  - (b) b с d a
  - (c) c d а

(d) c

(a) 10g

- [Ans: (b) b c d a] b d a
- 9. How many gram make 1 deca gram?
  - (b) 100g (c) 1kg (d) 100kg [Ans. (a) 10g]

b

- **10.** 1 nano second is equivalent to
  - (a)  $10^{-6}$ s (b)  $10^{-3}$ s
  - (c)  $10^{-15}$ s (d)  $10^{-9}$ s[Ans. (d)  $10^{-9}$ s]

#### **11.** Which unit is used to measure size of a nucleus?

(a) Angstrom (b) Micron (c) Nano (d) Fermi

[Ans. (d) Fermi]

10-39

 $10^{-2}$ 

 $10^{-13}$ 

(**d**)

#### **12.** One parallactic second is,

(a) $3.08 \times 10^{16}$ m	(b) $1.49 \times 10^{11}$ m
(c) $9.46 \times 10^{15}$ m	(d) $1.66 \times 10^{-27}$ m
	[A

#### [Ans. (a) $3.08 \times 10^{16}$ m]

#### **13.** How many light years make 1 parsec?

(a) 3.26 (b) 6.67 (c) 1.5 (d) 9.4 [Ans. (a) 3.26]

#### 14. How many AU makes one metre?

		[Ans. (d) $6.684 \times 10^{-12}$ AU]
(c)	$3.08 \times 10^{16} \mathrm{AU}$	(d) $6.684 \times 10^{-12} \text{AU}$
(a)	$3.26 \times 10^{11} AU$	(b) $1.496 \times 10^{11} \text{AU}$

#### **15.** One lunar month is equal to

(b) 27.3 days (a) 29 days (c) 365 days (d) 30 days

[Ans. (a) 29 days]

#### **16**. What is the value of one light year in tera metre?

(a) $9.46 \times 10^{6}$ Tm	(b) $9.46 \times 10^9 \mathrm{Tm}$
(c) $9.46 \times 10^{2}$ Tm	(d) $9.46 \times 10^3  \text{Tm}$
	[Ans. (d) $9.46 \times 10^3$ Tm]

- (a)  $2.59 \times 10^5$  km/h<sup>2</sup> (b)  $1.29 \times 10^5 \text{ km/h}^2$ (c)  $2.0 \times 10^3$  km/h<sup>2</sup> (d)  $3.5 \times 10^5 \text{ km/h}^2$ [Ans. (a)  $2.59 \times 10^5$  km/h<sup>2</sup>]
- **18.** Which device is used for measuring the mass of atoms?
  - (a) Spectrograph (b) Fermi
    - (d) Microscope

#### [Ans. (a) Spectrograph]

#### **19.** Which of the following statement is wrong?

(a) one fermi =  $10^{15}$ m

(c) Telescope

- (b) All non-zero digits are significant.
- (c)  $1 \text{ AU} = 1.496 \times 10^{11} \text{m}$
- (d) Speed is a derived unit.

#### [Ans. (a) one fermi = $10^{15}$ m]

#### **20.** Which of the following statement is wrong?

- (a) Strain is a dimensionless quantity.
- (b) Fundamental quantity is also called the base quantity.
- (c) force = mass  $\times$  acceleration
- (d) 1 Solar year = 1500 days.
  - [Ans. (d) 1 Solar year = 1500 days]

#### **21.** Which of the following statement is true?

- (a) Velocity is a fundamental unit.
- (b) 1 Solar day = 24 hours. (c) 1 Shake =  $10^4$ s
- (d) mass is a derived unit. [Ans. (b) 1 Solar day = 24 hours]
- 22. The number of significant figures in 0.0006012 m is (a) 3 (b) 4
  - (d) 5 [Ans. (b) 4]

(d)  $LT^{-3}$ 

[Ans. (d) 3]

[Ans. (c)  $LT^{-2}$ ]

**23.** The number significant figures in of  $2.64 \times 10^4$  kg is  $(1 \cdot)$ (a) 2

24. The displacement of a particle moving along x-axis with respect to times is given by  $x = at + bt^2 - ct^3$ . The dimensions of b are (b)  $L^0T^{-3}$ 

(a) 
$$L^{0}T^{-3}$$
  
(c)  $LT^{-2}$ 

Hin

(c) 7

- $x = at + bt^{2} ct^{3}$ : L = aT + bT^{2} cT^{3}
- $\therefore$  b = LT<sup>-2</sup>

#### Ph:9600175757 / 8124201000

#### orders@surabooks.com

Sura's XI Std - Physics I Unit 01 I Nature of Physical World and Measurement

**25.** If E & B respectively, represent electric field  $\frac{\Delta P}{P} = \frac{\Delta M}{M} + \frac{\Delta V}{V} \Rightarrow \frac{\Delta P}{P}\% = \left(\frac{\Delta M}{M} + 3.\frac{\Delta L}{T}\right)\%$ and magnetics induction field, then the ratio E & B has the dimensions of  $\left(\frac{\Delta P}{P}\right)\% = (5\% + 3.3\%) = (5 + 9) = 14\%$ (a) angle (b) acceleration displacement (c) velocity (d) Hint: [Ans. (c) velocity] **30.** The fractional error  $\left(\frac{\Delta x}{r}\right)$ (b)  $\pm n \left( \frac{\Delta a}{a} \right)$ (a)  $\pm \left(\frac{\Delta a}{\Delta a}\right)$  $\vec{F} = a (\vec{E} + \vec{v} \times \vec{B})$ (c)  $\pm n \log_e \left(\frac{\Delta a}{a}\right)$  (d)  $\pm n \log_{10} \frac{\Delta a}{a}$  $\therefore$  The dimensions of E are the same as those of vB[Ans. (b)  $\pm n \left(\frac{\Delta a}{a}\right)$ ]  $\therefore$  The dimensions of E/B = dimension of v. **26.** If force  $|\mathbf{F}|$ , velocity |v| and time  $|\mathbf{T}|$  are taken **31.** How many light years make 1 par sec? as to fundamental units then the dimensions of (a) 3.26 LY (b) 6.67 LY mass are (c) 1.5 LY (d) 9.4 LY (a)  $Fv^{-1}T$ (b)  $Fv^{-1}T$ (d)  $FvT^{-2}$ (c)  $FvT^{-1}$ [Ans. (a) 3.26 LY] [Ans. (c)  $FvT^{-1}$ ] **32.** If  $\pi = 3.14$ , then the value of  $\pi^2$  is Hint: (a) 9.8596 (b) 9.860 (c) 9.86 (d) 9.9  $\mathbf{F} = \frac{mv}{t} \qquad [\mathbf{m}] = [\mathbf{F}\frac{mv}{t}\mathbf{T}]$  $\therefore a = \frac{v}{-}$ [Ans. (c) 9.86] 33. Which of the following pairs of physical **27.** The dimensions of K.E. is quantities have same dimension? (a)  $M^2L^2T^{-1}$ (b)  $M^{1}L^{1}T^{1}$ (a) Force and Power (d)  $M^2L^2T^{-2}$ (c)  $M^{1}L^{2}T^{-2}$ (b) Stress and Pressure [Ans. (c)  $M^{1}L^{2}T^{-2}$ ] (c) Momentum and Moment of force (d) Torque and impulse of force 28. The dimensions of universal gravitational [Ans. (b) Stress and Pressure] constant is 34. The Dimensional formula for Boltzmann (a)  $M^{-2}L^{3}T^{-2}$  $M^{-2}L^{2}T^{-1}$ (b) constant is (d)  $ML^2T^{-1}$ (c)  $M^{-1}L^{3}T^{-2}$ (a)  $[ML^2T^{-1}]$ (b)  $[ATmol^{-1}]$ [Ans. (c)  $M^{-1}L^{3}T^{-2}$ ] (c)  $[ML^2T^{-2}K^{-1}]$ (d) None of the above **29.** The density of a cube is measured by measuring [Ans. (c)  $[ML^2T^{-2}K^{-1}]$ ] its mass and length of its side. If the maximum 35. Specific gravity (Relative Density) is an error in the measurement of mass ansd length example for are 5% and 3% respectively, the maxmimum (a) Dimensional Variables error in the measurement of density is (b) Dimensionless Variables (c) Dimensional Constant (a) 9% (b) 8% (d) Dimension less Constant (c) 14% (d) 2% [Ans. (b) Dimensionless Variables] Hint: [Ans. (c) 14%] **36.** 8.250 can be Rounded off to (a) 8.3 (b) 8.2 Density  $\rho = \frac{M}{V} = \frac{M}{L^3} \Delta M = 5\%; \Delta L = 3\%$ (c) 8.25 (d) 8.26 [Ans. (b) 8.2]

#### orders@surabooks.com

Sura's NI Std - Physics W Volume - I W Unit 01 W Nature of Physical World and Measurement 18 **37.** If E and B respectively, represent electric field || **4**. The temperature at which Saturated vapur, pure and melting ice are all in equilibrium is and magnetic field of Induction, then the ratio of E and B has the dimensional formula of called (a)  $[LT^{-2}]$ (b)  $[MLT^{-2}]$ (a) sublimation (b) melting point (c)  $[LT^{-1}]$ (d)  $[MLT^{-1}]$ [Ans. (c) [LT<sup>-1</sup>]] (c) Triple point of water (d) heat capacity **38.** Which one has more significant figures [Ans. (c) Triple point of water] (a) 600800 (b) 5213.0 5. The expression for Solid Angle is (c)  $2.65 \times 10^{24}$ (d) 0.0006032 (b) surface area /  $(radius)^2$ (a) rod/s [Ans. (b) 5213.0] (d) surface area / radius (c)  $(radius)^2$ **39.** Angle of 1 Second of arc is [Ans. (b) Surface Area /  $(radius)^2$ ] (a)  $48.5 \times 10^{-6}$  rad (b)  $0.485 \times 10^{-5}$  rad J Kg<sup>-1</sup>K<sup>-1</sup> is the unit for (c)  $4.85 \times 10^{-6}$  rad (d)  $48500 \times 10^{-6}$  rad **6**. • [Ans. (c)  $4.85 \times 10^{-6}$  rad] (a) Heat capacity (b) Latent heat (c) Specific heat (d) Energy **40.** 1 Yotta = (b)  $10^{-24}$ (c)  $10^{-21}$  (d)  $10^{24}$ [Ans. (c) Specific heat] (a)  $10^{21}$ [Ans. (d)  $10^{24}$ ] 7. 1 degree = rad. **41.** If mass of an electron is  $9.11 \times 10^{-31}$  Kg, then (a)  $1.754 \times 10^{-2}$ (b)  $1.745 \times 10^2$ how many electrons would weight in 1 mg? (c)  $1.745 \times 10^{-2}$ (d)  $1.547 \times 10^{-2}$ (a)  $1.68 \times 10^{18}$ (b)  $1.097 \times 10^{24}$ [Ans. (c)  $1.745 \times 10^{-2}$ ] (c)  $1.45 \times 10^{22}$ (d)  $1.970 \times 10^{23}$ 8. means a large world in which both [Ans. (b)  $1.097 \times 10^{24}$ ] objects and distances are large - sized. **B.** FILL IN THE BLANKS : (a) Macrocosm (b) Microcosm (c) Astronomy (d) Universe An attempt to explain a Macroscopic system 1. [Ans. (a) Macrocosm] in terms of its Microscopic constituents is 9. The largest practical unit of mass is . (a) CSL (b) Par sec (a) unification (b) Reductionism (c) Ly (d) AU (c) Microphysics (d) Macrophysics [Ans. (a) ChandraSekhar Limit (CSL)] [Ans. (b) Reductionism] **10.** The error caused due to the shear carelessness 2. The range of masses from heavenly bodies to of an observer is called electro is (a)  $10^{52}$  kg to  $10^{-28}$  kg (b)  $10^{55}$  kg to  $10^{+28}$  kg (b) Gross Error (a) Absolute Error (c)  $10^{55}$  kg to  $10^{-31}$  kg (d)  $10^{-55}$  kg to  $10^{31}$  kg (c) Instrumental Error (d) Zero Error [Ans. (b) Gross Error] [Ans. (c)  $10^{55}$  kg to  $10^{-31}$  kg] **11.** Ouantities which have constant values and also The CGS, MKS and SI system of units are 3. have no dimensions are called . system of units. (a) Dimensionless Constants (a) metric (b) cubic (b) Dimensional variables (c) periodic (d) atomic (c) Dimensionless constants [Ans. (a) metric] (d) Derived quantities [Ans. (a) Dimensionless Constants]

12.	Dime	nsional formul	a for I	Magne	tic Induction is	3		Prefix (	Symbol)		Sub	-Multiple
		•					(1)	Zepto (z	)	(a) 1	0-18	-
	(a) M	$T^2A^{-1}$	(b)	$MT^2A$	A		(2)	Pico (p)	·	(b) 1	0-1	
	(c) M	$T^{-2}A^{-1}$	(d)	$MA^{-1}$	l		(3)	atto (a)		(c) 1	$10^{-12}$	
				[Ans	$(c) MT^{-2}A^{-1}$		(4)	deci (d)		( <b>d</b> ) 1	$10^{-21}$	
13.	Form	ula (or) expre	ssion	for sur	face energy is		(1	) (2)	(3) (4	4)		
		•					(a) b	d d	a c	2		
	(a) we	ork / length	(b)	force	/ length		(b) d	l b	a c	2		
	(c) we	ork / time	(d)	work	/ area		(c) c	d	a t	)		
			A	ans. (d	l) work / area]		(d) d	l c	a ł	)	[Ans	s:(d) d c a b]
14.	Relati	ve error is also	o calle	d as	•	4.		Dev	vices		Pı	rinciples
	(a) G	ross Error	(b)	Perce	entage Error		(1)	Steam er	ngine	<b>(a)</b> ]	Berno	llis Principle
	(c) Al	bsolute Error	(d)	Fract	ional Error		(2)	Nuclear	Reactor	<b>(b)</b>	Laws	of
		I	(Ans. (	(d) Fra	actional error]						therm	odynamics
<b>15</b> .	The	name Physi	cs w	as ir	ntroduced by		(3)	Product of ultra	lon high	(c)	Contro	olled chain
		in 350 B	B.C					magneti	c fields	'	cacin	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
	(a) Th	nalus	(b)	Ptole	my		(4)	Aeropla	ne	(d) \$	Super	
	(c) An	ristotle	(d)	Cope	rnicus						condu	ctivity
				[An	s. (c) Aristotle]		(1	) (2)	(3) (4	4)		
<b>C</b> .	Мате	CH THE FOLLO	OWING	<b>;</b> :			(a) t	) C	a c	1		
							(b) c	1 a	bc	2		
1.		Branch		Ma	ijor Focus		(c) c	e d	a t	)		
	(1)	Acoustics	(a) A	About s and ene	space, time ergy		(d) t	о с	d a	ı	[Ans	s:(d) b c d a]
	(2)	High energy	( <b>b</b> ) <i>A</i>	About s	sound	5.		Types o	f fundam	ental		Strengths
	(3)	Quantum	(c) <i>I</i>	About 1	nature of par-		(1)	Gravita	tional for	ce	(a)	1
	(4)	Relativity	ו (d) ג	acies	liscrete nature		(2)	Electro	magnetic	force	(b)	10 <sup>-39</sup>
	(.)	Relativity	(u) 1 (	of phen	omena at		(3)	Weak n	uclear for	ce	(c)	10 <sup>-2</sup>
			8	atomic	and		(4)	Strong 1	nuclear fo	rce	(d)	10 <sup>-13</sup>
	(1	(2) (3)	(4)	Sub-ato	inic levels		(1	(2)	(3) (4	1)		
	(a) (	(2) $(3)$	(+)				(a) a	d	b (	.,		
	$(\mathbf{u}) \mathbf{c}$	a d	h				(b) b	) C	d a	1		
	(c) b	c d	a				(c) c	d	a ł	)		
	(d) d	c b	a	[A	Ans:(c) b c d a]		(d) c	a	b c	1	[Ans	: (b) b c d a]
2.		Physical Qua	antity		Unit	6.		Clas	sification	of	1	Examples
	(1)	<b>Force Constan</b>	t	(a)	Ns			q	uantities			-
	(2)	Boltzmann Co	nstant	(b)	Nm		(1)	Dimens	ional		(a)	<b>Pi</b> (T)
	(3)	Impulse		(c) (d)	JK <sup>1</sup> Nm <sup>-1</sup>			variabl	es			
	(1	) $(2)$ $(3)$	(4)	( <b>u</b> )			(2)	Dimens	ion less		<b>(b)</b>	Planck's
	1) h (s)	(2) $(3)$	(+) b					variabl	e			Constant
	(h) h	a c	h				(3)	Dimens	ional con	istant	: (c)	Velocity
	(c) $h$	d a	C				(4)	Dimensi	on less co	nstan	t ( <b>d</b> )	Strain
	(d) d	b a	c	[A	ns:(a) d c a bl							

#### orders@surabooks.com

20				Sura	<b>i's ■</b> XI Std - Physics 🗰 Vo	olume -	I 🗰 Unit 01 🗰 Nature of Physical World and Measurement
	(1 (a) b	) (2) d	(3) a	(4) c		2.	<ul><li>(a) Absolute Error</li><li>(b) Relative Error</li><li>(c) Percentage Error</li><li>(d) Gross Error</li></ul>
	(b) b (c) c	c d	a b	d a			[Ans. (d) Gross Error]
	(d) a	с	d	b	[Ans: (c) c d b a]	3.	<ul> <li>(a) Solar clock</li> <li>(b) Electronic Oscillators</li> <li>(c) Radio active dating</li> <li>(d) Electronic balance</li> </ul>
7.		Physica	al quant	tity	Dimensional		[Ans. (d) Electronic balance]
	(1)	Surface	Tension	1	$\begin{array}{c c} Formula \\ \hline (a) & [ML^{-1}T^{-1}] \end{array}$	4.	(a) Energy (b) Work
	(2)	Heat Ca	pacity	(	(b) [ML <sup>2</sup> ]		(c) Torque (d) Force [Ans. (d) Force]
	(3)	Moment	t of Iner	rtia	(c) $[MT^{-2}]$	5.	(a) Length (b) Mass
	(4)	Co-effici viscosity	ient of		(d) $[ML^2 I^{-2} K^{-1}]$		(c) Time (d) Volume [Ans. (d) Volume]
	(1	) (2)	(3)	(4)		6.	(a) f.p.s (b) c.g.s
	(a) c (b) $a$	d d	b	a b			(c) m.k.s (d) r.m.s [Ans. (d) r.m.s]
	(b) $a$ (c) $a$	c u	d	b		7.	(a) Optics (b) Acoustics
	(d) b	а	с	d	[Ans: (a) c d b a]		(c) Astrophysics (d) Nuclear Physics
8.		Er	rors		Cause		
	(1)	Systema Errors	atic	(a	) due to shear carelessness	8.	<ul> <li>(a) Force constant</li> <li>(b) Planck's constant</li> <li>(c) Boltzmann constant</li> <li>(d) Refractive Index</li> </ul>
	(2)	Randon	n Error	s (b	) Fractional Error	F	[Ans. (d) Keiracuve index]
	(3)	Relative	e Errors	(C)	) Reproducible		CHOOSE THE INCORRECT FAIR .
					inaccuracies	1.	(a) Work - Energy (b) Strange Dreasure
	(1	) (2)	(3)	(4)			(c) Force - Tension
	(a) the table $(b)$ of the table $(b)$ of table		a a	a b			(d) Surface Tension - Force
	(c) a	n c	d	b			[Ans. (d) Surface Tension - Force]
	(d) t	o a	d	c	[Ans:(b) d c a b]	2.	(a) Velocity - Angular velocity
9.		Num	bers		Significant figures		(b) Force - Torque
	(1)	40.00	0	(a)	6		(d) Frequency - Wavelength
	(2)	0.00345	0	(D) (C)	5		[Ans. (d) Frequency - Wavelength]
	(4)	307000.		( <b>d</b> )	4	3.	(a) Density - Relative Density
	(1	) (2)	(3)	(4)	·		(b) Strain - refractive Index
	(a) b	d	c	а			(c) $\pi$ - e (d) Planaly's Constant Stafar's constant
	$(\mathbf{D})$ $\mathbf{D}$	c d	a a	a b			(d) Planck's Constant - Steran's constant [Ans. (a) Density - Relative Density]
	(d) d	c	b	a	[Ans:(d) d c b a]		(a) Hoat Energy
D.	Сно	OSE THE	ODD O	ONE O	OUT:	4.	(a) Heat - Energy (b) Mass - Inertia
1.	(a) Sr	pecific or	avity	(h)	) Strain		(c) Charge - Current
	(c) ret	fractive ir	ndex	(d)	Planck's constant		(d) Moment of force - Torque
			<b>[A</b> ]	ns. (d	l) Planck's constant]		[Ans. (c) Unarge - Uurrent]

#### orders@surabooks.com

Sura's XI Std - Physics I Unit 01 Nature of Physical World and Measurement

**F.** CHOOSE THE CORRECT PAIR :

- 1. (a) 30.00 2009 (b) 0.00345 2.6 (c) 0.040500 - 20100m (d) 153 - 3072 [Ans. (a) 30.00 - 2009]
- (a) Telescope Microscope
  (b) Screw gauge Radar Method
  - (c) Parallax Method -Vernier Caliper
  - (d) Spring balance Common balance

[Ans. (d) Spring balance - Common balance]

- **3.** (a) Torque Nm
  - (b) Planck's constant J / s
  - (c) Specific heat J kg  $k^{-1}$
  - (d) Moment of Inertia  $kg/m^2$

[Ans. (a) Torque - Nm]

#### G. Assertion & Reason :

#### **Directions :**

- (a) Assertion and Reason are correct and Reason is correct explanation of Assertion
- (b) Assertion and Reason are true but Reason is the false explanation of the Assertion
- (c) Assertion is true but Reason is false
- (d) Assertion is false but Reason is true
- 1. Assertion: Attempting to explain diverse physical phenomena with a few concepts and laws is unification.
  - **Reason** : Attempting to explain a macroscopic system in terms of its microscopic constituents is reductionism.

[Ans. (b) Assertion and Reason are true but Reason is the false explanation of the Assertion]

2. Assertion: Study of light is called optics Reason : Properties of light is studied in optics. They are Reflection, Retraction etc.,

[Ans. (a) Assertion and Reason are correct and Reason is correct explanation of Assertion]

- **3.** Assertion: Quantities that can be expressed in terms of fundamental quantities are derived quantities.
  - **Reason** : Examples are Mass, Length, Time etc.,
  - [Ans. (c) Assertion is true but Reason is false]

 Assertion: In centimeter, the spelling meter is internationally accepted.
 Reason : 'Metre' is internationally used unit whereas 'Meter' is used by Americans.
 [Ans. (d) Assertion is false but Reason is true]
 Assertion: Mass due to rotational motion is moment of Inertia.
 Reason : Rotational mass explains about radius of Gyration.
 [Ans. (a) Assertion and Reason are correct and Reason is correct explanation of

Assertion]

- 6. Assertion: Very large distances such as distance of a planet or star can be measured by parallax method.
  - Reason : For measuring small masses of atomic / sub - atomic particles, mass spectrograph is used.
  - [Ans. (b) Assertion and Reason are true but Reason is the false explanation of the Assertion]
- 7. Assertion: The least value that can be measured using screw gauges, vernier calipers is called least count.
  - Reason : The magnitude of difference between the true value and the measured value is called relative error or fractional error.

#### [Ans. (c) Assertion is true but Reason is false]

- 8. Assertion: The rounding off of 27.653 upto 3 digits is 27.6
  - **Reason** : 10200 has three significant figures
  - [Ans. (d) Assertion is false but Reason is true]
- 9. Assertion: Dimensional analysis method is used to convert a physical quantity from one system of units to another

Reason : 
$$\frac{1}{2}$$
 mv<sup>2</sup> = mgh

[Ans. (b) Assertion and Reason are true but Reason is the false explanation of the Assertion]

#### orders@surabooks.com

22

Sura's Nature of Physical World and Measurement

H. CHOOSE THE CORRECT OR INCORRECT || 7. **STATEMENTS :** 1. (I) RADAR method is used for measurement (a) I only of length in the case of long distances (II) The uncertainty in a measurement is called error. Which statement is correct? 8. (b) II only (a) I only (c) Both are correct (d) None [Ans. (c) Both are correct] (I)  $G_{CGS} = 6.6 \times 10^{-8} \text{ dyne } \text{Cm}^2 \text{ g}^{-2}$ 2. (a) I only (II) T = 2  $\pi$   $\frac{g}{2}$ Which statement is correct? 9. (a) I only (b) II only (c) Both are correct (d) None [Ans. (a) I only] 3. (I) Expression for charge is current / time (a) I only (II) Expression for Faraday constant is Avagadro constant × elementary charge Which statement is correct? (a) I only (b) II only (c) Both are correct (d) None [Ans. (b) II only] 1. (I) Force constant and Faraday constant are 4. examples for Dimensional constant Ans. (i) (II) Radius of gyration does not depend on moment of Inertia. Which statement is incorrect? (a) I only (b) II only (c) Both are correct (d) None 2. [Ans. (b) II only] (I) The ratio of mean absolute error to the 5. method? mean value is called fractional error (II) Due to the wrong observations Recording, Random errors occur. Which statement is correct? (a) I only (b) II only (c) Both are correct (d) None [Ans. (a) I only] 3. 6. (I) Distance of moon from earth is 10<sup>11</sup>m (II) Mass of a cell is 10<sup>-10</sup> kg Which statement is incorrect? 'nature'. (a) I only (b) II only (c) Both are correct (d) None [Ans. (a) I only]

(I) Least count of screw gauge is 0.01mm (II) Least count of vernier caliper is 0.1mm Which one is correct? (b) II only (c) both are correct (d) none [Ans. (c) Both are correct] (I) Parallax angle,  $\theta = \frac{\text{Unknown distance}}{(x)}$ base (b) (II) Distance of the planet,  $d = Velocity of radio wave (V<sup>e</sup>) \times time taken (t)$ Which statement is incorrect? (b) II only (c) Both are correct (d) None [Ans. (a) I only] (I) Frequency and angular velocity has same dimensional formula (II) Torque is also called as rotational force Which one is correct? (b) II only (c) Both are correct (d) None [Ans. (c) Both are correct] **II. VERY SHORT ANSWER QUESTIONS :** =2 Marks — What is science? The word science comes from a Latin word 'scientia' meaning 'to know'. (ii) Science is a systematic attempt to understand natural phenomena in much knowledge gained through systematic observation and experiment.

What are the steps involved in scientific method? (or) What are the general features of scientific

- Ans. (i) Systematic observation
  - (ii) Controlled experimentation
  - (iii) Reasoning (qualitative and quantitative)
  - (iv) Modelling (Mathematical)
  - (v) Prediction and verification (theories)
- What is Physics?
- Ans. (i) Physics is a branch of science.
  - (ii) The word comes from a Greek word meaning
  - (iii) It deals with the study of nature and natural phenomena.

Sura's NI Std - Physics World and Measurement

#### 4. What is mechanics?

- Ans. (i) Mechanics is a branch of physics.
  - (ii) It is divided into statics and Dynamics.
  - (iii) Mechanics deals with the study of motion of particles, deformable bodies and general system of particles.

#### **5.** Write a note on scope of physics.

- Ans. (i) Physics has a large scope as it covers a various physical quantities. (Length, mass, time, energy).
  - (ii) It deals with the Macroscopic group (Mechanics, electrodynamics, thermodynamics and optics) and microscopic group (Quantum physics).
     Example:

Range of time scale (astronomical scale to microscopic group (Quantum physics).

#### 6. What is MKS system ?

- **Ans.** The mks system is based on metre, kilogram and second as the fundamental units of length, mass, and time respectively.
- 7. What is the aim of our Science Education?
- **Ans.** According to part IV Article 51A (h) of Indian Constitution "It shall be the duty of every citizen of India to develop scientific temper, humanism and spirit of inquiry and reform". This is the aim of our Science Education.

#### 8. Name three practical units to measure mass.

- **Ans.** (i) Pound 1b = 0.4536 kg
  - (ii) Quintal  $1q = 100^{-9} \text{ kg}$
  - (iii) Atomic mass unit (1amu) =  $1.66 \times 10^{-27}$  kg

#### 9. Define Solar Year.

- Ans. (i) It is the time taken by the earth to complete one revolution around the sun in its orbit.
  - (ii) 1 Solar year = 365.25 average Solar days
     = 366.25 Sedrial days.

#### **10.** What is Leap year?

**Ans.** The year which is divisible by 4 and in which the month of February has 29 days is called Leap year.

#### **11.** Name three practical units to measure Area.

Ans. <mark>(i</mark> )	Barn,	1 barn	=	$10^{-28} \text{ m}^2$
<b>(ii)</b>	Acre,	1 acre	=	$4047 \ m^2$
	TT t	1 1	_	1042

(iii) Hectare,  $1 \text{ hectare} = 10^4 \text{ m}^2$ .

#### orders@surabooks.com

## **12.** What is the importance of physical quantity? What are its types?

- Ans. (i) Physical quantities are important to understand the properties of materials.
  - (ii) It is classified into fundamental physical quantity and derived physical quantity.

#### **13**. Define unit of a physical quantity.

**Ans.** Unit of a physical quantity is defined as the established standard used for comparison of the given physical quantity. It is classified in to fundamental and derived unit.

## **14.** What are the things needed to express the measurement of a physical quantity?

- Ans. (i) The unit in which the quantity is measured
  - (ii) The numerical value or the magnitude of the quantity (n), the number of times that unit (u) is contained, in the given physical quantity. Q = nu.

## **15.** Name the prefixes for powers of ten with its symbol.

- **Ans.** (i)  $10^1$  = deca and its symbol is da
  - (ii)  $10^6$  = mega and its symbol is M
    - (iii)  $10^{12}$  = tera and its symbol is T
    - (iv)  $10^{-1}$  = deci and its symbol is d
    - (v)  $10^{-6}$  = micro and its symbol is  $\mu$
    - (vi)  $10^{-12}$  = pico and its symbol is p
- **16.** Name four units to measure extremely small distances.
- Ans. Units used to measure extremely small distances are
  - (i) 1 micron or micrometer,  $1 \mu m = 10^{-6} m$
  - (ii) 1 nanometer,  $1 \text{ nm} = 10^{-9} \text{m}$
  - (iii) 1 Angstrom unit  $1\text{\AA} = 10^{-10}\text{m}$
  - (iv) 1 Fermi, 1 Fm =  $10^{-15}$ m.

## **17.** Name three units to measure extremely large distances.

- Ans. Units used to measure extremely large distances are,
  - (i) Astronomical Unit :

It is the mean distances of the earth from the sun. 1 AU =  $1.496 \times 10^{11}$ m.

24

Sura's XI Std - Physics W Volume - I W Unit 01 W Nature of Physical World and Measurement

#### (ii) Light year :

It is the distance travelled by light in vacuum in one year. 1 ly =  $9.46 \times 10^{15}$  m

(iii) Parallactic second :

It is the distance at which an arc of length 1 astronomical unit subtends an angle of 1 second of arc. 1 par sec =  $3.084 \times 10^{16}$ m = 3.26 ly.

- **18.** What is an error? Name the three Errors in Measurement.
- Ans. The uncertainty in a measurement is called an error.

The three possible errors are

- (i) Systematic errors
- (ii) Random errors
- (iii) Gross errors

#### **19.** What is Absolute Error.

- **Ans.** The magnitude of difference between true value and the measured value of a quantity is called absolute error.
  - $\Delta a_{\rm n} = a_{\rm m} a_{\rm n}$

#### 20. What is Mean Absolute error?

**Ans.** The arithmetic mean of the magnitude of absolute errors in all the measurements is called the mean absolute error.

If  $a_{\rm m}$  is the true value and  $\Delta a_{\rm m}$  is the mean absolute error, then the magnitude of the quantity may lie between  $a_{\rm m} + \Delta a_{\rm m}$  and  $a_{\rm m} - \Delta a_{\rm m}$ .

#### **21**. What is Relative error?

**Ans.** The ratio of the mean absolute error to the mean value is called relative error. This is also called as fractional error.

Relative error = 
$$\frac{\text{Mean absolute error}}{\text{Mean value}} = \frac{\Delta a_{\text{m}}}{a_{\text{m}}}$$

#### 22. What is Percentage error?

Ans. The relative error expressed as a percentage is called Percentage error.

Percentage error = 
$$\frac{\Delta a_{\rm m}}{a_{\rm m}} \times 100\%$$

#### 23. What is significant figures?

**Ans.** The digits that are known reliably plus the first uncertain digit are known as significant figures or significant digits.

### 24. What is meant by the dimensions of a physical quantity?

**Ans.** The dimensions of a physical quantity are the powers to which the units of base quantities are raised to represent a derived unit of that quantity.

#### 25. What are Dimensional variables? Give example.

- **Ans.** Physical quantities, which possess dimensions and have variable values are called dimensional variables. Examples are length, velocity, and acceleration etc.
- 26. What is meant by Dimensionless variables? Give example.
- **Ans.** Physical quantities which have no dimensions, but have variable values are called dimensionless variables. Examples are specific gravity, strain, refractive index etc.

#### **27.** Define Dimensional Constant. Give example.

- **Ans.** Physical quantities which possess dimensions and have constant values are called dimensional constants. Examples are Gravitational constant, Planck's constant etc.
- 28. What is meant by Scientific method?
- **Ans.** The scientific method is a step-by-step approach in studying natural phenomena and establishing laws which govern these phenomena.
- 29. What do you mean by unification and reductionism?
- Ans. Unification: Attempting to explain diverse physical phenomena with a few concepts and laws is unification.
   Reductionism: An attempt to explain a macroscopic system in terms of its microscopic constituents is reductionism.

#### **30.** What is Classical mechanics?

*Ans.* The study of forces acting on bodies whether at rest or in motion.

#### **31**. What is Thermodynamics?

*Ans.* The study of the relationship between heat and other forms of energy.

#### **32**. What is the meaning of Acoustics?

**Ans.** The study of the production and propagation of sound waves.

#### **33**. What is Astrophysics?

*Ans.* The branch of physics which deals with the study of the physics of astronomical bodies.

#### orders@surabooks.com

Sura's NI Std - Physics World and Measurement

#### **34**. What is meant by Quantum mechanics?

- *Ans.* The study of the discrete nature of phenomena at the atomic and subatomic levels.
- **35.** Which branches of physics deal at the level of atom & nucleus?
- Ans. Atom : Atomic physics. Nucleus : Nuclear physics.

#### **36.** What are types of discoveries in physics?

- Ans. (i) Accidental discoveries and well-analysed research outcome in the laboratory based on intuitive thinking and prediction.
  - (ii) For example, magnetism was accidentally observed but the reason for this strange behavior of magnets was later analysed theoretically.
  - (iii) This analysis revealed the underlying phenomena of magnetism. With this knowledge, artificial magnets were prepared in the laboratories.

#### **37**. What is meant by Range of time scales?

**Ans.** Range of time scales: astronomical scales to microscopic scales,  $10^{18}$ s to  $10^{-22}$ s.

#### **38**. What is meant by Range of masses?

**Ans.** Range of masses: from heavenly bodies to electron,  $10^{55}$  kg (mass of known observable universe) to  $10^{-31}$  kg (mass of an electron) [the actual mass of an electron is  $9.11 \times 10^{-31}$  kg].

### **39.** How Physics is related to technology and define technology with respect to Physics.

**Ans.** Technology is the application of the principles of physics for practical purposes. The application of knowledge for practical purposes in various fields to invent and produce useful products or to solve problems is known as technology.

#### **40**. In what ways physics is in relation to astronomy?

**Ans.** Astronomical telescopes are used to study the motion of planets and other heavenly bodies in the sky. Radio telescopes have enabled the astronomers to observe distant points of the universe. Studies of the universe are done using physical principles.

#### 41. Define the SI unit of length.

*Ans.* Metre is the SI unit of length. One metre is the length of the path travelled by light in vacuum in

299,792,458 of a second.

- **42.** Define the SI unit of mass. (or) What is one kilogram in SI system of units?
- **Ans.** Kilogram is the SI unit of mass. One kilogram is the mass of the prototype cylinder of platinum iridium alloy (whose height is equal to its diameter), preserved at the International Bureau of Weights and Measures at Serves, near Paris, France.

### **43.** Define the SI unit of time. (or) What is one second in SI system of units?

- **Ans.** Second is the SI unit time. One second is the duration of 9,192,631,770 periods of radiation corresponding to the transition between the two hyperfine levels of the ground state of Cesium-133 atom.
- 44. Define the SI unit of electric current. (or) What is one second in SI system of units? (or) Define one ampere (S.I standard for current)
- **Ans.** "ampere" is the SI unit of electric current. One ampere is the constant current, which when maintained in each of the two straight parallel conductors of infinite length and negligible cross section, held one metre apart in vacuum shall produce a force per unit length of  $2 \times 10^{-7}$  N/m between them.
- **45.** What is the SI unit of temperature and define it? What is one kelvin in SI system of units?
- Ans. Kelvin is the SI unit of temperature. One degree

kelvin is the fraction of  $\left(\frac{1}{273.16}\right)$  of the thermodynamic

temperature of the triple point of the water.

- **46.** What is the SI unit of amount of substance? (or) What is one mole in SI system of units? (or) Define one mole (S.I standard for amount of substance)
- *Ans.* Mole is the SI unit of amount of substance. One mole is the amount of substance which contains as many elementary entities as there are atoms in 0.012 kg of pure carbon-12.

#### orders@surabooks.com

Sura's ■ XI Std - Physics III Volume - I IIII Unit 01 IIII Nature of Physical World and Measurement

# **47.** What is meant by one candela? and which base quantity is measured by this unit? (or) Define one candela (S.I standard for Luminous intensity)

**Ans.** One candela is the luminous intensity in a given direction, of a source that emits monochromatic radiation of frequency  $5.4 \times 10^{14}$  Hz and that has a radiant intensity of  $\frac{1}{683}$  watt/steradian in that direction.

#### **48.** What is meant by the triple point of water?

**Ans.** Triple point of water is the temperature at which saturated vapour, pure and melting ice are all in equilibrium. The triple point temperature of water is 273.16K.

#### **49.** What is meant by Parallax?

**Ans.** Parallax is the name given to the apparent change in the position of an object with respect to the background, when the object is seen from two different positions.

#### 50. What is 1 Light year?

**Ans.** 1 Light year is distance travelled by light in vacuum in one year. 1 Light Year =  $9.467 \times 10^{15}$  m.

#### **51.** Define a Astronomical Unit.

- **Ans.** 1 astronomical unit is the mean distance of the Earth from the Sun.  $1 \text{ AU} = 1.496 \times 10^{11} \text{ m.}$
- **52.** What is parsec? (or) Define one parsec (parallactic second)
- Ans. 1 parsec, Parallactic second is the distance at which an arc of length 1 AU subtends an angle of 1 second of arc 1 parsec =  $3.08 \times 10^{16}$  m = 3.26 light year.
- **53**. Why is the cylinder used in defining kilogram made up of platinum-iridium alloy?
- **Ans.** This is because the platinum-iridium alloy is least affected by environment and time.

#### 54. Write the largest and the smallest practical unit of mass and time respectively. (or) Define Chandrasekar Limit (CSL)

Ans. Chandrasekhar Limit (CSL) is the largest practical unit of mass.1 CSL = 1.4 times the mass of the Sun. The smallest practical unit of time is Shake. 1 Shake =  $10^{-8}$  s.

#### **55.** Define mass.

- Ans. Mass of a body is defined as the quantity of matter contained in a body. The SI unit of mass is kilogram (kg).
- **56.** Write the masses of tiny as well as huge matter?
- **Ans.** A tiny mass is of electron  $(9.11 \times 10^{-31} \text{kg})$ . The huge mass is of the known universe (=  $10^{55} \text{ kg}$ ).

### **57.** Write the methods to determine the masses of objects?

Ans. The mass of an object is determined in kilograms using a common balance. Spring balance, electronic balance are also used. For measuring larger masses like that of planets, stars etc., gravitational methods are used. For measurement of small masses of atomic/ subatomic particles etc., mass spectrograph is used.

#### 58. What is clock? Write the principle and its types.

- Ans. A clock is used to measure the time interval. An atomic standard of time, is based on the periodic vibration produced in a Cesium atom. Some of the clocks developed later are electric oscillators, electronic oscillators, solar clock, quartz crystal clock, atomic clock, decay of elementary particles, radioactive dating.
- **59.** Which units are used to measure large distance i.e. distance of planets and stars? Which method is used for measurement?
- Ans. AU The mean distance between earth and star  $1 \mbox{ AU}$  =  $1.496 \times 10^{11} \mbox{ m}$ 
  - Light year The distance travelled by light in vacuum in one year

1 light year =  $9.46 \times 10^{15}$  m

Parsec - Distance at which an arc of length of 1 AU subtends an angle of 1 sec at a point parsec =  $3.08 \times 10^{16}$  m.

Parallax method is used to measure large distance.

### **60.** Is it possible to have length and velocity both as fundamental quantities? Why?

**Ans.** No, length is fundamental quantity whereas velocity is the derived quantity.

Sura's XI Std - Physics I Unit 01 I Nature of Physical World and Measurement

- 61. Which of these unit is largest: AU, light year and parsec. Express the average distance of earth from the sun in (i) light year (ii) per sec.
- Ans. Parsec is the largest unit.

parsec > light year > AU

Average distance of earth from the sun is (d) (astronomical unit).

- (i) d =  $1 \text{ AU} = 1.496 \times 10^{11} \text{ m}$ =  $\frac{1.496 \times 10^{11}}{9.46 \times 10^{15}} = 1.58 \times 10^{-5}$  light year. (ii) d =  $\frac{1.496 \times 10^{11}}{3.08 \times 10^{16}}$  par sec. =  $4.86 \times 10^{-6}$  par sec.
- 62. The radius of gold nucleus is 41.3 Fermi. Express its volume in m<sup>3</sup>.

**Ans.** Radius of gold nucleus = 
$$41.3 \times 10^{-15}$$
 m

Volume (V) = 
$$\frac{4}{3}\pi r^3 = \frac{4}{3} \times 3.14 \times (41.3 \times 10^{-15})^3$$
  
V = 2.95 × 10<sup>-40</sup> m.

#### **63.** Describe the relation of Physics with mathematics.

Ans. (i) Physics is a quantitative science.

- (ii) Physics is closely related to mathematics as a tool for its development.
- **64.** What is the difference between Accuracy and Precision?

Ans.

S.No.	Accuracy	Precision
1.	Measurements	Measurements
	close to true value.	close to each
		other.
2.	All the accuracy	All the precised
	values are	values
	precised.	are not accurate.

#### **65.** Describe the Personal errors.

**Ans.** These errors occur due to individual performing experiment without initial setting up or careless observation without precautions.

#### **66.** Describe the errors due to external causes.

**Ans.** These errors are due to external conditions like change in temperature, humidity or pressure during an experiment.

#### 67. What is dimensional equation? Give an example.

- **Ans.** When the dimensional formula of a physical quantity is expressed in the form of an equation, such equation is known as the dimensional equation. Ex: acceleration =  $[M^0LT^2]$
- **68.** Define FPS system of units.
- Ans. It is the British Engineering system of units, which uses foot, pound and second as the basic units for measuring length, mass and time respectively.
- 69. Define CGS system of units.
- Ans. It is the Gaussian system of units, which uses centimeter, gram and second as the basic units for measuring length, mass and time respectively.
- 70. Define MKS system of units.
- Ans. It is the Metric system of units, which uses metre, kilogram and second as the basic units for measuring length, mass and time respectively.
- 71. Define one radian (S.I standard for plane angle)
- **Ans.** It is the angle subtended at the centre of a circle by an arc equal in length to the radius of the circle.  $180^{\circ}$

$$1 \text{ rad} = \frac{100}{\pi} = 57.27^{\circ}.$$

- 72. Define one steradian (S.I standard for solid angle)
- **Ans.** It is the solid angle subtended at the centre of a sphere, by that surface of the sphere, which is equal in area to the square of radius of the sphere.
- **III. SHORT ANSWER QUESTIONS :**

- **1.** Give any three applications of physics in our society.
- Ans. (i) Basic laws of electricity and magnetism led to the discovery of wireless communication technology which has shrunk the world with effective communication over large distances.
  - (ii) The launching of satellite into space has revolutionized the concept of communication.
  - (iii) Microelectronics, lasers, computers, superconductivity and nuclear energy have comprehensively changed the thinking and living style of human beings.

28

Sura's ■ XI Std - Physics III Volume - I IIII Unit 01 IIII Nature of Physical World and Measurement

## 2. What are fundamental quantities and derived quantities? (Sep. 2020)

#### Ans. Fundamental quantities

Fundamental quantities are quantities which cannot be expressed in terms of any other physical quantity.

**Example:** Quantities like length, mass, time, temperature are fundamental quantities.

#### **Derived quantities:**

Quantities that can be expressed in terms of fundamental quantity are called derived quantities. **Example:** Quantities like area, volume, velocity are derived quantities.

### 3. What are fundamental units and derived units?

#### Ans. Fundamental units:

The units in which the fundamental quantities are measured are called fundamental units. It is also known as base units.

#### **Derived units:**

The units in which the derived quantities are measured are called Derived units. **Example:** 

Unit of speed = 
$$\frac{\text{Unit of distance}}{\text{Unit of time}} = \frac{\text{m}}{\text{s}} = \text{ms}^{-1}$$

ms<sup>-1</sup> is a derived unit.

4. Name the SI unit for electric current and give a definition for it.

Ans. The SI unit for electric current is ampere (A) Definition:

One ampere is the constant current, which when maintained in each of the two straight parallel conductors of infinite length and negligible cross section, held one metre apart in vacuum shall produce a force per unit length of  $2 \times 10^{-7}$  N/m between them.

5. Name the SI unit for Luminous intensity and give a definition for it.

**Ans.** The SI unit for Luminous intensity is candela. Its symbol is cd.

#### **Definition:**

One candela is the luminous intensity in a given direction, of a source that emits monochromatic radiation of frequency  $5.4 \times 10^{14}$  Hz and that has

a radiant intensity of  $\frac{1}{638}$  watt/steradian in that direction.

6. Give the values for the following units with prefixes.

- (i) 1 Mega ohm (ii) 1 milliampere
- (iii) 1 deca gram (iv) 1 nano second
- (v) 1 micro volt (vi) 1 centimetre

#### Ans.

- (i) 1 Mega ohm (M $\Omega$ ) =  $10^6 \Omega$
- (ii) 1 milliampere (mA) =  $10^{-3}$ A
- (iii) 1 daca gram (da g) = 10g
- (iv) 1 nano second (ns)  $= 10^{-9}$ s
- (v) 1 microvolt ( $\mu$ V) = 10<sup>-6</sup>V
- (vi) 1 centimetre (cm)  $= 10^{-2}$ m

#### 7. What are the advantages of the SI system?

- *Ans.* (i) This system makes use of only one unit for one physical quantity, which means a rational system of units.
  - (ii) In this system, all the derived units can be easily obtained from basic and supplementary units, which means it is a coherent system of units.
  - (iii) It is a metric system which means that multiples and submultiples can be expressed as powers of 10.

## 8. Distinguish between fundamental and derived units.

#### Ans.

Fu	indamental Units	<b>Derived Units</b>		
1.	Using fundamental units fundamental quantities are measured	Using derived units, derived quantities are measured		
2.	These units cannot be expressed in terms of other fundamental units	These units can be expressed in terms of all fundamental units.		
3.	Examples : metre, kilogram, second (m, s, kg. A. mol)	Examples: metre/ second, newton/ meter, kilogram/ metre seconds. (m/s, kg/m <sup>3</sup> )		

#### 9. Give any three practical units of time.

#### Ans. (i) Solar year :

It is the time taken by the earth to complete one revolution around the sun in its orbit. 1 solar year = 365.25 average solar days.

#### orders@surabooks.com

Sura's XI Std - Physics I Unit 01 I Nature of Physical World and Measurement

#### (ii) Leap year:

The year which is divisible by 4 and in which the month of February has 29 days is called leap year.

#### (iii) Lunar month :

It is the time taken by the moon to complete one revolution around the earth in its orbit. 1 lunar month = 27.3 days.

**10.** What is Gross Error & How can it be minimised?

#### Ans. Gross Error

- (i) The error caused due to the shear carelessness of an observer is called gross error. For example
- (ii) Reading an instrument without setting it properly.
- (iii) Taking observations in a wrong manner without bothering about the sources of errors and the precautions.
- (iv) Recording wrong observations. These errors can be minimized only when an observer is careful and mentally alert.

#### **11.** Explain Random errors.

#### Ans. Random errors

- (i) Random errors may arise due to random and unpredictable variations in experimental conditions like pressure, temperature, voltage supply etc.
- (ii) Errors may also be due to personal errors by the observer who performs the experiment. Random errors are sometimes called "chance error".
- (iii) When different readings are obtained by a person every time he repeats the experiment, personal error occurs.
- (iv) For example, consider the case of the thickness of a wire measured using a screw gauge.
- (v) The readings taken may be different for different trials. In this case, a large number of measurements are made and then the arithmetic mean is taken.
- (vi) If n number of trial readings are taken in an experiment, and the readings are a<sub>1</sub>, a<sub>2</sub>, a<sub>3</sub>,....a<sub>n</sub>. The arithmetic mean is

$$a_{m} = \frac{a_{1} + a_{2} + a_{3} + \dots + a_{n}}{a_{m}} = \frac{1}{n} \sum_{i=1}^{i=n} a_{i}$$

Usually this arithmetic mean is taken as the best possible true value of the quantity.

#### **12**. Explain unification with example.

- Ans. (i) Attempting to explain diverse physical phenomena with a few concepts and laws is unification.
  - (ii) Newton's universal law of gravitation explains the motion of freely falling bodies towards the Earth, motion of planets around the Sun, motion of the Moon around the Earth, thus unifying the fundamental forces of nature.

#### **13**. Explain reductionism with example.

- *Ans.* (i) An attempt to explain a macroscopic system in terms of its microscopic constituents is reductionism.
  - (ii) For example, thermodynamics was developed to explain macroscopic properties like temperature, entropy, etc., of bulk systems.
  - (iii) The above properties have been interpreted in terms of the molecular constituents (microscopic) of the bulk system by kinetic theory and statistical mechanics.

#### 14. How are theoretical predictions useful?

- Ans. (i) Theoretical predictions aided with recent simulation and computation procedures are widely used to identify the most suited materials for robust applications.
  - (ii) The pharmaceutical industry uses this technique very effectively to design new drugs. Bio compatible materials for organ replacement are predicted using quantum prescriptions of physics before fabrication.
  - (iii) Thus, experiments and theory work, hand in hand, complimenting one another.

#### **15.** In what way physics is exciting us?

- *Ans.* A small number of basic concepts and laws can explain diverse physical phenomena.
  - (i) The most interesting part is the designing of useful devices based on the physical laws.
     For example i) use of robotics ii) journey to Moon and to nearby planets with controls from the ground iii) technological advances in health sciences etc.

#### orders@surabooks.com

Sura's ■ XI Std - Physics III Volume - I IIII Unit 01 IIII Nature of Physical World and Measurement

- (ii) Carrying out new challenging experiments to unfold the secrets of nature and in verifying or falsifying the existing theories.
- (iii) Probing and understanding the science behind natural phenomena like the eclipse, and why one feels the heat when there is a fire? (or) What causes the wind, etc.

#### **16.** Write the role of Physics in Technology.

- Ans. (i) Basic laws of electricity and magnetism led to the discovery of wireless communication technology which has shrunk the world with effective communication over large distances.
  - (ii) The launching of satellite into space has revolutionized the concept of communication.
  - (iii) Microelectronics, lasers, computers, superconductivity and nuclear energy have comprehensively changed the thinking and living style of human beings.

#### **17.** In what way Physics is in relation to Chemistry.

- Ans. (i) In physics, we study the structure of atom, radioactivity, X-ray diffraction etc.
  - (ii) Such studies have enabled researchers in chemistry to arrange elements in the periodic table on the basis of their atomic numbers.
  - (iii) This has further helped to know the nature of valence and chemical bonding and to understand the complex chemical structures. Inter-disciplinary branches like Physical chemistry and Quantum chemistry play important roles here.

#### **18.** What is the relation of Physics to Biology?

- Ans. (i) Biological studies are impossible without a microscope designed using physics principles.
  - (ii) The invention of the electron microscope has made it possible to see even the structure of a cell. X-ray and neutron diffraction techniques have helped us to understand the structure of nucleic acids, which help to control vital life processes. X-rays are used for diagnostic purposes.
  - (iii) Radio-isotopes are used in radiotherapy for the cure of cancer and other diseases. In recent years, biological processes are being studied from the physics point of view.

**19.** How is physics useful in geology and oceanography?(or) Describe the relation of Physics with Geology.

### Ans. (i) Diffraction techniques helps to study the crystal structure of various rocks.

- (ii) Radioactivity is used to estimate the age of rocks, fossils and the age of the Earth.
- (iii) Oceanographers seek to understand the physical and chemical processes of the oceans. They measure parameters such as temperature, salinity, current speed, gas fluxes, chemical components.

#### **20.** How can the systematic errors be minimised?

- Ans. (i) By choosing the instrument carefully.
  - (ii) Necessary correction is to be made.
  - (iii) High precision instrument is to be used.
  - (iv) Proper setting up of experiments is to be done.
  - (v) Taking proper precautions is a must, while making observations.
- **21.** What are systematic errors? (or) What are the Classifications of Systematic errors?
- Ans. Systematic error is reproducible inaccuracies that are consistently in the same direction. They can be classified as
  - (i) Instrumental errors,
  - (ii) Imperfection in experimental technique or procedure
  - (iii) Personal errors.
  - (iv) Errors due to external causes
  - (v) Least count error.
- **22.** Describe the relation of Physics with Psychology.
- *Ans.* (i) All the psychological interactions can be derived from a physical process.
  - (ii) The movements of neurotransmitters are governed by the physical properties of diffusion and molecular motion.
  - (iii) The function of our brain is related to our underlying dualism (wave -particle nature).

#### 23. Describe Instrumental errors. How is it minimised?

- *Ans.* (i) It is happened when an instrument is not calibrated properly at the time of manufacture.
  - (ii) For example, If a measurement is made with a meter scale whose end is worn out, result obtains error.

Sura's NI Std - Physics W Unit 01 Nature of Physical World and Measurement

(iii) These errors can be rectified by using the good quality instruments.

#### **24**. Write a note on parallax method.

- **Ans. (i)** Parallax is the name given to the object with respect to the background, when the object is seen from two different positions.
  - (ii) The distance between the two positions is called basis (b).
  - (iii) This method is used for measuring very large distance such as distance of a planet or star.

#### **IV. LONG ANSWER QUESTIONS :**

**—5** Marks —

- **1.** Discuss the relation of physics with other branches of science
- **Ans.** Physics is the most fundamental branch of science. It has played a key role in the development of all branches of sciences.

#### **Physics in relation to mathematics**

- Physics is a quantitative science. Mathematics provides the necessary signs and tools which the physicist use.
- It has played an important role in the development of theoretical physics.
- Had newton not invented calculus, he would not have been able to discover the universal law of gravitation.

#### Physics in relation to chemistry:

- In physics, we study the structure of atom, radio activity, X-ray diffraction, etc.
- Such studies have enabled chemists to arrange elements in the periodic table on the basis of their atomic numbers.
- This has further helped to know the nature of valency and chemical bonding and to understand the complex chmical structures.

#### **Physics in relation to biology:**

orders@surabooks.com

- The developments in life sciences a great deal to physics.
- Optical microscopes are extensively used in the study of biology.
- With the help of an electron microscope, one can study the structure of cell.
- The X-rays and neutron diffraction techniques have helped in understanding the structure of nucleic acids, which helped to control vital life process.
- Radio isotopes are used in radiation therapy for the cure of deadly diseases like cancer.

2. Explain propagation of errors in the difference of two quantities and also in the division of two quantities.

#### Ans. Errors in the difference of two quantities :

Let  $\Delta A$  and  $\Delta B$  be the absolute errors in the two quantities, A and B, respectively. Then, Measured value of  $A = A \pm \Delta A$ Measured value of  $B = B \pm \Delta B$ 

Consider the difference, Z = A - B

The error 
$$\Delta Z$$
 in Z is then given by  
 $Z \pm \Delta Z = (A \pm \Delta A) - (B \pm \Delta B)$   
 $= (A - B) \pm (\Delta A + \Delta B)$ 

$$= Z \pm (\Delta A + \Delta B)$$

(or)  $\Delta Z = \Delta A + \Delta B$ 

The maximum error in difference of two quantities is equal to the sum of the absolute errors in the individual quantities.

Error in the division or quotient of two quantities

Let  $\Delta A$  and  $\Delta B$  be the absolute errors in the two quantities A and B respectively.

Consider the quotient,  $Z = \frac{A}{B}$ 

The error  $\Delta Z$  in Z is given by

$$Z \pm Z = \frac{A \pm \Delta A}{B \pm \Delta B} = \frac{A \left(1 \pm \frac{\Delta A}{A}\right)}{B \left(1 \pm \frac{\Delta B}{B}\right)}$$
$$= \frac{A}{B} \left(1 \pm \frac{\Delta A}{A}\right) \left(1 \pm \frac{\Delta B}{B}\right)^{-1}$$
$$Z \pm \Delta Z = Z \left(1 \pm \frac{\Delta A}{A}\right) \left(1 \mp \frac{\Delta B}{B}\right)$$

[using  $(1+x)^n \approx 1 + nx$ , when x << 1] Dividing both sides by Z, we get,

or

$$1 \pm \frac{\Delta Z}{Z} = \left(1 \pm \frac{\Delta A}{A}\right) \left(1 \mp \frac{\Delta B}{B}\right)$$
$$= 1 \pm \frac{\Delta A}{A} \mp \frac{\Delta B}{B} \pm \frac{\Delta A}{A} \cdot \frac{\Delta B}{B}$$

As the terms  $\Delta A/A$  and  $\Delta B/B$  are small, their product term can be neglected.

The maximum fractional error in Z is given by

$$\frac{Z}{Z} = \left(\frac{\Delta A}{A} + \frac{\Delta B}{B}\right)$$

The maximum fractional error in the quotient of two quantities is equal to the sum of their individual fractional errors.

32

Sura's XI Std - Physics Wolume - I W Unit 01 W Nature of Physical World and Measurement

#### **3.** Write to causes of errors in measurement.

#### Ans.

(i)	Least count error	Associated with the
		poor resolution of
		the instrument
(ii)	Instrumental	Associated with the
	errors	faulty calibration
		or change in
		conditions.
(iii)	Random errors	Getting difficult
		results for the same
		measurement done
		repeatedly
(iv)	Personal errors	Associated with the
		individual performing
		the experiment ie.
		improper precautions,
		incorrect initial set up
		of experiment.
(v)	Systematic errors	Which tends to be in
		the same direction.

#### 4. Write the rules for rounding off. (or) Explain the rules framed for rounding off the numbers with the examples.

#### Ans.

	Rule	Example			
i)	If the digit to be dropped is smaller than 5, then the preceding digit should be left unchanged.	i) ii)	7.32 is rounded off to 7.3 8.94 is rounded off to 8.9		
ii)	If the digit to be dropped is greater than 5, then the preceding digit should be increased by 1	i) ii)	17.26 is rounded off to 17.3 11.89 is rounded off to 11.9		
iii)	If the digit to be dropped is 5 followed by digits other than zero, then the preceding digit should be raised by 1	i) ii)	7.352, on being rounded off to first decimal becomes 7.4 18.159 on being rounded off to first decimal, become 18.2.		

iv)	If the digit to be dropped is 5 or 5 followed by zeros, then the preceding digit is not changed if it is even	i) ii)	3.45 is rounded off to 3.4 8.250 is rounded off to 8.2.
v)	If the digit to be dropped is 5 or 5 followed by zeros, then the preceding digit is raised by 1 if it is odd	i) ii)	3.35 is rounded off to 3.4 8.350 is rounded off to 8.4

## 5. Explain the rules framed to count significant figures with the examples.

#### Ans.

	Rule	Example			
1.	All non-zero digits are significant	1342 has four significant figures			
2.	All zeros between two non-zero digits are significant	2008 has four significant figures 30700. has five significant figures 30700 has three significant figures			
3.	All zeros right to nonzero digit but left to decimal point are significant.				
4.	The terminal or trailing zeros in the number without decimal point are not significant.				
5.	All zeros are significant if the number given with measurement unit.	30700 m has five significant figures.			
6.	f a number is less than1, the zeros between decimal point and first non- zero digit are not significant but the zeros right to last non-zero digit are significant.	i) ii) iii)	0.00345 has three significant figures. 0.030400 has five significant figures. 40.00 has four significant figures.		
7.	The number of significant figures doesn't depend on the system of units used	1.53 cm, 0.0153 m, 0.0000153 km all have three significant figures.			

Sura's Nature of Physical World and Measurement





#### orders@surabooks.com

34

Sura's NI Std - Physics Wolume - I W Unit 01 W Nature of Physical World and Measurement

- **11**. The angle of an object is 18.2°. What is the angular diameter of the object in radians?
  - (b)  $3.64 \times 10^{-2}$  rad (a) 36.4 rad (d) 3.17 rad
  - (c)  $31.74 \times 10^{-2}$  rad

[Ans. (c)  $31.74 \times 10^{-2}$  rad]

$$\theta = 18.2^{\circ} = \frac{18.2}{60 \times 60} \times \frac{\pi}{180} = \frac{18.2}{60 \times 60} \times \frac{3.14}{180}$$

 $= 31.74 \times 10^{-2}$  rad

- **12.** If a circle with 10 m radius and angle 60° at centre, then what will be the length of arc?
  - (a) 5.24m (b) 6.21m
  - (c) 7.1 mm (d) 10m [Ans. (a) 5.24m]

Hint:

Radius r = 10m, angle 
$$\theta = \frac{\pi}{6}$$
 (60°)

Length of the arc (l)  $= r\theta = 10 \times \frac{\pi}{6} = 5.24$ m

- 13. The mass of an iron sheet is 0.250 kg and volume of the sheet is 1.5m<sup>3</sup>. Then what is the density of the iron sheet? Express the result in SI unit system.
  - (b)  $0.167 \text{ kg m}^{-3}$ (a)  $0.267 \text{kg m}^{-3}$ (c)  $0.255 \text{ kg m}^{-3}$ (d) 0.285 kg m<sup>-3</sup>

Hint:

#### [Ans. (b) 0.167 kg m<sup>-3</sup>]

[Ans. (c) kg ms $^{-1}$ ]

[Ans. (b)  $m^2$ ]

Mass of the iron sheet = 0.250 kgVolume of the iron sheet =  $1.5 \text{ m}^3$ 

Density = 
$$\frac{Mass}{Volume} = \frac{0.250}{1.5} = 0.167 \text{ kg m}^{-3}$$

(b)  $ms^{-2}$ 

(d) kg  $m^2 s^{-1}$ 

#### **14.** What is the SI unit of linear momentum?

- (a)  $ms^{-1}$
- (c) kg ms<sup>-1</sup>

Hint:

Linear momentum = mass  $\times$  velocity SI unit of mass = kg

SI unit of velocity =  $\frac{\text{distance}}{\text{time}} = \frac{\text{m}}{\text{s}} = \text{ms}^{-1}$ Linear momentum = kg ms<sup>-1</sup>

#### **15.** What is the SI unit of Area?

(a) m

(b)  $m^2$ (c)  $\text{Nm}^{-1}$  (d)  $\text{cm}^{-1}$ 

Hint:

#### Area = length $\times$ breadth SI unit of length = mSI unit of breadth $\Rightarrow$ (length) = m Area = length $\times$ breadth $= m \times m = m^2$ .

16. SI unit of the universal constant of gravitation (G) is,

(a)  $kg^{-2} m^{-2}$ (b) kg ms<sup>-1</sup> (c)  $Nm^2 kg^{-2}$ (d)  $Nm^{-1}$ 

Ans. (c) 
$$Nm^2 kg^{-2}$$
]

*Hint*: Universal constant of gravitation (G) is,

$$= \frac{Fr^{2}}{m_{1}m_{2}} \qquad [\because F = G. \frac{m_{1}m_{2}}{r^{2}}]$$
SI unit of force = N  
SI unit of distance (r) = m = r^{2} = m^{2}
SI unit of masses m<sub>1</sub> and m<sub>2</sub> = kg<sup>2</sup>  

$$\Rightarrow \frac{Fr^{2}}{m_{1}m_{2}} = \frac{Nm^{2}}{kg^{2}} = Nm^{2} kg^{-2}.$$

$$= 2 Marks =$$

1. Using a screw gauge the thickness of a wire was measured as 5mm. Calculate (i) the fractional error (ii) the percentage error.

Given data:

Thickness of the wire (t) = 5mmAccurancy  $\Delta t = 0.01$  mm

#### Solution:

- (i) Fractional error =  $\delta_t = \frac{\Delta t}{t} = \frac{0.01}{5} = 0.002$ .
- (ii) Percentage error  $= \delta_t = \frac{\Delta t}{t} \times 100\%$  $= 0.002 \times 100\%$ = 0.02%
- If a mass of a proton is  $1.67 \times 10^{-27}$ kg, how 2. many protons will be present in 1kg?

#### Given data:

Mass of a proton  $= 1.67 \times 10^{-27}$ kg  $1.67 \times 10^{-27}$ kg is mass of 1 proton.

#### Solution:

1 kg is the mass of  $\frac{1}{1.67 \times 10^{-27}}$  protons  $= 0.5988 \times 10^{27}$  protons  $= 5.988 \times 10^{26}$  protons  $5.988 \times 10^{26}$  protons will be present in 1kg.

#### orders@surabooks.com

Sura's XI Std - Physics I Unit 01 I Nature of Physical World and Measurement

**3.** Calculate angle of 1 second of arc.

#### Solution:

1 second of arc = 
$$1'' = \frac{1^{\circ}}{60 \times 60}$$
  
=  $\frac{1}{60 \times 60} \times \frac{\pi}{180} = \left(\frac{1}{3600} \times \frac{3.14}{180}\right) \left[\because 1^{\circ} = \frac{\pi}{180}\right]$   
1 second of arc =  $4.85 \times 10^{-6}$  rad.

4. The radius of a nucleus is  $1.5 \times 10^{-15}$ m of the order fermi. Find the volume of the nucleus.

#### Solution:

Radius of a nucleus 
$$r = 1.5 \times 10^{-15}$$
m.  
Volume of the nucleus  $V = \frac{4}{3} \pi r^3$ .

$$=\frac{4}{3}\times 3.14\times (1.5\times 10^{-15})^3 = \frac{4}{3}\times 3.14\times (4.5\times 10^{-45})$$

 $= 18.84 \times 10^{-45} m^3$ 

Volume of the nucleus is  $V = 18.84 \times 10^{-45} \text{m}^3$ 

5. A beam of metal has length, breadth and height as 4m, 3m and 5m respectively. Then what will be the volume of the metal beam? Express the result in SI unit system.

#### Solution:

- Volume = length × breadth × height SI unit of volume  $(v) = m \times m \times m = m^3$ SI unit of length (l) = 4mSI unit of breadth (b)= 3mSI unit of height (h) = 5mVolume  $(v) = l \times b \times h = 4 \times 3 \times 5 = 60m^3$
- 6. Find the SI unit of moment of inertia. 5.64kg mass of a object is moving uniformly. The radius of gyration is measured as 30cm of an object. Then what is the moment of Inertia?

#### Solution:

$$\begin{split} Mass &= 5.64 kg \\ radius of gyration &= 30 cm = 0.3m \\ SI unit of a mass is kg \\ SI unit of the radius (gyration) &= m^2 = 0.9 \\ Moment of inertia (I) &= mass \times radius of gyration \\ &= 5.64 kg \times 0.9m = 1.5228 \ kgm^2 \end{split}$$

 The ratio of stress and strain of a wire is 3:2. Find the co-efficient of elasticity. Express the result in SI unit system.

#### Solution:

The ratio of stress and strain of a wire is 3 : 2

SI unit of stress =  $\frac{\text{Force}}{\text{Area}} = \frac{\text{N}}{\text{m}^2} = \text{Nm}^{-2}$ Strain is dimensionless variable.

So, co-efficient of elasticity of a wire is =  $\frac{\text{Stress}}{\text{Strain}}$ 

$$= \frac{3 \text{ Nm}^{-2}}{2 \text{ (No unit)}} = 1.5 \text{ Nm}^{-2}.$$

- =3 Marks —
- **1.** In a following physical units, how many units are there in 1 metre?
  - (i) 1 Astronomical unit (AU =  $1.496 \times 10^{11}$  m
  - (ii) 1 light year =  $9.467 \times 10^{15}$ m
  - (iii) 1 micron ( $\mu$ ) = 10<sup>-6</sup>m
  - (iv) 1 parallactic second (parsec) =  $3.08 \times 10^{16}$ m

#### Given data:

1 AU =  $1.496 \times 10^{11}$ m 1 ly =  $9.467 \times 10^{15}$ m 1 mm =  $10^{-6}$ m 1 parsec =  $3.08 \times 10^{16}$ m

#### Solution:

(i)  $1.496 \times 10^{11}$ m is equivalent to 1 AU.

1 metre is equivalent to  $\frac{1}{1.496 \times 10^{11}}$ 

=  $0.6684 \times 10^{-11} = 6.68 \times 10^{-12}$  AU In one metre,  $6.68 \times 10^{-12}$  astronomical units are present

(ii)  $9.467 \times 10^{15}$ m is equivalent to 1 ly. 1 metre is equivalent to  $\frac{1}{9.467 \times 10^{15}}$ 

=  $0.1056\times 10^{-15}$  =  $1.05\times 10^{-16}$  ly In one metre,  $1.05\times 10^{-16}$  light years are present.

36

Sura's XI Std - Physics W Volume - I W Unit 01 W Nature of Physical World and Measurement

- (iii)  $10^{-6}$ m equivalent of 1µm 1 metre is equivalent to  $\frac{1}{10^{-6}} = 10^{6}$ µm In one metre  $10^{6}$  microns are present
- (iv)  $3.08 \times 10^{16}$  m equivalent to 1 parsec.

1 metre is equivalent to  $\frac{1}{3.08 \times 10^{16}}$ 

=  $0.324\times10^{-16}$  =  $3.24\times10^{-17}m$  In one metre  $3.24\times10^{-17}$  parsec are present.

## 2. How many parallactic second are there in one Astronomical unit?

#### Given data:

- 1 parallactic second =  $3.08 \times 10^{16}$  m
- 1 Astronomical unit =  $1.496 \times 10^{11}$  m

Solution:

$$\frac{1 \text{ AU}}{1 \text{ parsec}} = \frac{1.496 \times 10^{11}}{3.08 \times 10^{16}} = \frac{1.496 \times 10^{11} \times 10^{-16}}{3.08}$$

$$= \frac{1.496 \times 10^{-5}}{3.08} = 0.485 \times 10^{-5}$$
  
= 4.85 × 10<sup>-6</sup> par sec.

 $4.85\times10^{-6}$  parsec present in one astronomical unit.

3. If mass of an electron is  $9.11 \times 10^{-31}$  kg, how many electrons would weigh in 1 mg?

#### Given data:

Mass of an electron =  $9.11 \times 10^{-31}$ kg  $9.11 \times 10^{-31}$ kg is the mass of 1 electron

Solution:

 $\begin{array}{ll} 9.11\times 10^{-31} \text{kg} &= 9.11\times 10^{-31}\times 10^3\,\text{g} \\ &= 9.11\times 10^{-28}\,\text{g} \\ &= 9.11\times 10^{-25}\,\text{mg} \end{array}$ 

1mg is the mass of  $\frac{1}{9.11 \times 10^{-25}}$  mg electron

 $= 0.1097 \times 10^{25} electrons = 1.097 \times 10^{24} electrons$ 

=  $1.097 \times 10^{24}$  electrons would weigh in 1 mg.

4. The unit of length convenient on the atomic scale is known as angstrom and is denoted by Å. The size of a helium atom is about 30 pico meter. What is the total atomic volume in metre<sup>3</sup> of one mole of helium atom?

#### Solution:

1Å = 100 pm  $\Rightarrow$  30pm = 0.3Å pm  $\rightarrow$  picometer (10<sup>-12</sup>m) Å = Angstrom (10<sup>-10</sup>)m Radius of the helium atom (r) = 30pm = 0.3Å = 0.3 × 10<sup>-10</sup>m Volume of helium nucleus V=  $\frac{4}{3}$  × 3.14 × (0.3 × 10<sup>-10</sup>)m<sup>3</sup> = 1.256 × 10<sup>-30</sup>m<sup>3</sup> Number of atoms in 1 mole of helium atom = Avogadro's number(N) = 6.023 × 10<sup>23</sup> Atomic volume of 1 mole of helium atoms (v') V' = V × N

#### Solution:

The radius of the nucleus = 60.2 fermi =  $60.2 \times 10^{-15}$ m

Volume of the platinum nucleus

$$= \frac{4}{3} \pi r^{3} \text{m}^{3} = \frac{4}{3} \times 3.14 \times (60.2 \times 10^{-15})^{3} \text{ m}^{3}$$
$$= \frac{4}{3} \times 3.14 \times (218167 \times 10^{-45})$$
$$= \frac{4}{3} \times 3.14 \times (2.18167 \times 10^{-40})$$
$$= 9.133 \times 10^{-40} \text{m}^{3}$$

Volume of the platinum nucleus is  $9.133\times 10^{-40}m^3.$ 

fermi is used for measure very small distance so it is used to measure the nuclear size of an atom. It is denoted as 1 Fermi =  $10^{-15}$ m.

Sura's NI Std - Physics IN Unit 01 Nature of Physical World and Measurement

#### VALUE BASED QUESTIONS

- 1. (i) Monica was watching the night sky. She saw a star, moving towards her, with increase in brightness. After some few minutes when she watched closely, she found it was the light from a flight in the sky. She was surprised, but initially the flight looked stationary, after some time it was glowing brightly moving towards her. So she went and asked her father. Why this effect occured?
  - (ii) Is RADAR used in launching a missile from the ground to hit the target (i.e. fight air craft)?
- Ans. (i) Any moving object, which is perpendicular to our eye sight will look like a stationary thing for a while. But as the moving object changes its angle of vision, its movement will be known.
  - (ii) Yes, RADAR is used in launching missile from the ground.

#### **Creative Questions (HOTS)**

- 1. Why has 'second' been defined is terms of periods of radiations from cesium-133?
- Ans. Second has been defined is terms of periods of radiations because
  - (i) This period is accurately defined
  - (ii) This period is not affected by change of physical conditions like temperature. pressure and volume etc.
  - (iii) The unit is easily reproducible in any good laboratory.
- 2. How many Astronomical units are there in one light year?

#### Given data:

1 Astronomical unit =  $1.496 \times 10^{11}$ m

1 light year =  $9.46 \times 10^{15}$ m

#### Solution:

$$\frac{1 \text{ ly}}{1 \text{ AU}} = \frac{9.46 \times 10^{15}}{1.496 \times 10^{11}} = \frac{9.46 \times 10^{15} \times 10^{-11}}{1.496}$$
$$= \frac{9.46 \times 10^4}{1.496} = 6.32 \times 10^4 \text{AU}$$

In  $6.32 \times 10^4$ AU are present in one light year.

#### orders@surabooks.com

**3.** When the planet Jupiter is at a distance of 824.7 million kilometers from the earth, its angular diameter is measured to be 35.72 of arc. Calculate the diameter of Jupiter.

Distance,  $D = 824.7 \times 10^6 \text{ km}$ 

$$\theta = 35.72'' = \frac{35.72}{60 \times 60} \times \frac{\pi}{180}$$
 rad

Diameter, 
$$d = ?$$
:  $d = D \times \theta$ 

$$= 824.7 \times 10^{6} \times \frac{35.72}{60 \times 60} \times \frac{\pi}{180} \text{ km}$$
$$= 824.7 \times 10^{6} \times \frac{35.72}{3600} \times \frac{\pi}{180} \text{ km}$$
$$= 824.7 \times 10^{6} \times \frac{35.72 \times 3.14}{3600 \times 180} \text{ km}$$

Diameter of Jupiter =  $1.427 \times 10^5$  km

In a submarine fitted with a SONAR, the time delay between generation of a signal and reception of its echo from an enemy ship is 110.3 seconds. If speed of sound in water is 1450 ms<sup>-1</sup> then calculate the distance of the enemy ship from the submarine.

Solution:

**4**.

Speed of a sound in water 1450 ms<sup>-1</sup> Time delay T = 110.3s

Distance of the enemy ship,  $D = \frac{vT}{2}$ 

$$=\frac{1450\times110.3}{2}=\frac{159935}{2}=79967 \text{ m}$$

Distance of enemy ship = 79.96 km

5. In an ocean surveillance system of ship fitted with a (RADAR), the time delay between generation of a radio waves reflected from an enemy ship is observed to be 5.6s. Calculate the distance of the enemy ship from the surveillance ship.

#### Solution:

Time delay T = 5.6s

$$t = \frac{\mathrm{T}}{2} = \frac{5.6}{2} = 2.8$$

Speed of radiowaves = speed of light (v)

 $= 3 \times 10^8 m s^{-1}$ 

38

Sura's ■ XI Std - Physics III Volume - I IIII Unit 01 IIII Nature of Physical World and Measurement

Distance from the surveillance ship to enemy ship D =  $v \times t = 3 \times 10^8 \times 2.8 = 8.4 \times 10^8 \text{m}$ Distance (D) =  $8.4 \times 10^8 \text{ km}$ .

#### **CONCEPTUAL QUESTIONS**

- 1. Why is it convenient to express the distance of stars in terms of light year (or) parsec rather than in km?
- **Ans.** The distances of astronomical objects like stars, planets etc from the earth are huge. The distance on the earth are relatively small so it can be measured in km.

#### For Example :

The distance to be next nearest big galaxy Andromeda is 21,000,000,000,000,000,000 km. i.e.  $21 \times 10^{18}$  km.

This number is so large that it becomes hard to write and to interpret.

So astronomical units like light year, parsec A.U are used for large distances.

2. Show that a screw gauge of pitch 1 mm and 100 divisions is more precise than a vernier caliper with 20 divisions on the sliding scale.

Ans. Least count of screw gauge

 $=\frac{\text{Pitch}}{\text{No. of divisions}} = \frac{1}{100} = 0.01 \text{ mm} \text{ (or) } 0.001 \text{ cm}$ on the Head scale. Least count of vernier calipers

$$= 1MSD - 1VSD = (1 - \frac{19}{20}) MSD$$

$$=\frac{1}{20}=0.05$$
 cm.

So screw gauge is more precise than vernier.

- **3.** If humans were to settle on other planets, which of the fundamental quantities will be in trouble? Why?
- *Ans.* Time will be in trouble. Time becomes irrelevant. Because day and year based on spinning and revolution of the planet. So each planet has its own year length.

Eg. : Uranus and Neptune move too slow.

- 4. Having all units in atomic standards is more useful. Explain.
- Ans. All units in atomic standards are more useful because they never change with time.
- 5. Why dimensional methods are applicable only up to three quantities?

*Ans.* If a quantity depends on more than three factors than dimensional formula cannot be derived.

Because on equating the powers of M, L & T on either side of the dimensional equation, three equations can be obtained, from which only three unknown dimensions can be calculated.

 $\star \star \star \star \star$