

CHEMISTRY

11th Standard

VOLUME - I & II

Based on the Updated New Textbook

Salient Features

- Prepared as per the Updated New Textbook.
- Answers for all Textual Questions.
- Exhaustive Additional MCQs, VSA, SA & LA questions with answers are given in each chapter.
- All the **Objective Type (1 Mark)** questions, are given with 4 options.
 - (i) Choosing the correct option
- (ii) Matching

(iii) Filling the blanks


- (iv) Assertion & Reason
- (v) Choosing the correct Statement
- (vi) Choosing the Incorrect Statement

PH: 8124201000 / 8124301000

- Govt. Model Question Paper-2018 [Govt. MQP-'18], First Mid-Term Test 2018 [First Mid-'18], Quarterly Exam 2018, 2019, 2023 & 2024 [Qy-'18, '19, '23 & '24], Half Yearly Exam 2018, 2019 & 2023 [Hy '18, '19 & '23], Public Exam March 2019, May 2022, March 2023 & 2024 [Mar-'19, May- '22, Mar-'23 & '24], Instant Supplementary Exam June 2019, August 2022. July-2023 & 2024 [June-'19, Aug-'22, July-'23 & '24], Govt. Suppl. Exam. September 2020 & 2021 [Sep-2020 & Sep-'21] and Common Revision Test 2022 [CRT-'22] questions are incorporated in the appropriate sections.
- Half yearly Exam December 2024-25 Question Paper is given with answers.



For Orders Contact



All rights reserved © SURA Publications.

No part of this book may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, digitally, electronically, mechanically, photocopying, recorded or otherwise, without the written permission of the publishers. Strict action will be taken

Code No: SG 266

Authors:

Mrs. P. Thilagam PGT (Chemistry)

Chennai

Published by:

Mr. Subash Raj, B.E., M.S.

Our Guides for XI & XII Standard

- 🚓 சுராவின் தமிழ் உரைநூல்
- * Sura's Smart English
- Sura's Mathematics (EM/TM)
- ❖ Sura's Physics (EM/TM)
- Sura's Chemistry (EM/TM)
- Sura's Bio-Botany & Botany (EM/TM) (Short Version & Long Version)
- Sura's Bio-Zoology & Zoology (EM/TM) (Short Version & Long Version)
- ❖ Sura's Computer Science (EM/TM)
- Sura's Computer Applications (EM/TM)
- Sura's Commerce (EM/TM)
- ❖ Sura's Economics (EM/TM)
- Sura's Accountancy (EM/TM)
- Sura's Business Maths (EM)

NOTE FROM PUBLISHER

It gives me great pride and pleasure in bringing to you **Sura's Chemistry** guide Vol. I & II for **11**th **Standard**. A deep understanding of the text and exercises is rudimentary to have an insight into the subject. The students have to carefully understand the topics and exercises.

Sura's Chemistry 11th Standard Guide encompasses all the requirements of the students to comprehend the text and the evaluation of the textbook.

It will be a teaching companion to teachers and a learning companion to students.

- ▲ Chapter Snapshot, Concept Map, Formulae to Remember, Must know Definitions are given in almost each chapter.
- ▲ Exhaustive Additional MCQs, SA, LA questions with Answers are given in each chapter. Solved numerical problems are given wherever necessary.
- ▲ These features will help students practice and learn effectively all the sections of the textbook.

In order to learn effectively, I advise students to learn the subject sectionwise and practice the exercises given.

Though these salient features are available in our Sura's Chemistry Guide 11th Standard, I cannot negate the indispensable role of the teachers in assisting the student to understand the subject thoroughly.

I sincerely believe this guide satisfies the needs of the students and bolsters the teaching methodologies of the teachers.

I pray the almighty to bless the students for consummate success in their examinations.

Subash Raj, B.E., M.S.
- Publisher
Sura Publications

All the Best

Head Office:

Sura Publications

1620, 'J' Block, 16th Main Road, Anna Nagar, **Chennai - 600 040.**

Phone: 044 - 4862 9977, 044 - 4862 7755.

E-mail: orders@surabooks.com **Website**: www.surabooks.com

For Orders Contact

80562 94222

81242 01000

81243 01000

96001 75757

98409 26027

PH: 8124201000 / 8124301000

1000/00/0

CONTENTS

	VOLUME - I		
Units		Page No.	Month
1	Basic Concepts of Chemistry and Chemical Calculations	1 - 46	June
2	Quantum Mechanical Model of Atom	47 - 76	July
3	Periodic Classification of Elements	77 - 104	August
4	Hydrogen	105 - 126	September
5	Alkali and Alkaline Earth Metals	127 - 148	October
6	Gaseous State	149 - 178	June
7	Thermodynamics	179 - 212	July

VOLUME - II							
8	Physical and Chemical Equilibrium	213 - 242	August				
9	Solutions	243 - 272	October				
10	Chemical Bonding	273 - 306	November				
11	Fundamentals of Organic Chemistry	307 - 338	July				
12	Basic Concept of Organic Reactions	339 - 358	August				
13	Hydrocarbons	359 - 396	October				
14	Haloalkanes and Haloarenes	397 - 420	November				
15	Environmental Chemistry	421 - 436	December				
Half y answe	early Exam - December 2024-25 Question Paper with ers.	437 - 444					

I Mid Term Test	- June & July (1, 6, 2, 11, 7)	
Quarterly Examination	- June to September (1, 6, 2, 11, 7, 3, 8, 12, 4)	
II Mid Term Test	- October & November (5, 9, 13, 10, 14)	
Half Yearly Examination	- Full Portion	

TO ORDER WITH US

SCHOOLS and TEACHERS:

We are grateful for your support and patronage to 'SURA PUBLICATIONS'

A/c Name

A/c Name

IFSC

IFSC

IFSC

A/c Name

Kindly prepare your order in your School letterhead and send it to us.

For Orders contact: 81242 01000 / 81243 01000

DIRECT DEPOSIT

A/c Name : Sura Publications

Our A/c No. : **36550290536**

Bank Name : STATE BANK OF INDIA

Bank Branch: Padi

IFSC : SBIN0005083

A/c Name : Sura Publications

Our A/c No. : 6502699356 Bank Name : INDIAN BANK

Bank Branch: Asiad Colony : IDIB000A098 **IFSC**

: Sura Publications A/c Name Our A/c No. : 13240200032412

Bank Name : FEDERAL BANK

Bank Branch: Anna Nagar

IFSC : FDRL0001324

After Deposit, please send challan and order to our address.

email to: orders@surabooks.com/Whatsapp:81242 01000.



For Google Pay: 98409 26027



For PhonePe: 98409 26027

PH: 8124201000 / 8124301000

: Sura Publications

: Sura Publications

: Sura Publications

Our A/c No. : 21000210001240

: UCBA0002100

: KVBL0001154

Our A/c No. : 1154135000017684

Our A/c No. : 50200031530945

Bank Branch: Cenotaph Road, Teynampet

: HDFC0001216

Bank Name : UCO BANK Bank Branch: Anna Nagar West

Bank Name : KVB BANK

Bank Branch: Anna Nagar

Bank Name : HDFC BANK



DEMAND DRAFT / CHEQUE

Please send Demand Draft / cheque in favour of 'SURA PUBLICATIONS' payable at Chennai. The Demand Draft / cheque should be sent with your order in School letterhead.

STUDENTS:

Order via Money Order (M/O) to

SURA PUBLICATIONS

1620, 'J' Block, 16th Main Road, Anna Nagar, Chennai - 600 040.

Phone: 044-4862 9977, 044-4862 7755.

Mobile: 96001 75757 / 81242 01000 / 81243 01000. Email: orders@surabooks.com Website: www.surabooks.com





www.surabooks.com

CONTENTS

1.	Basic Concepts of Chemistry and Chemical Calculations	3 – 46
2.	Quantum Mechanical Model of Atom	47 – 76
3.	Periodic Classification Of Elements	77 – 104
4.	Hydrogen	105 – 126
5.	Alkali and Alkaline Earth Metals	127 – 148
6.	Gaseous State	149 – 178
7.	Thermodynamics	179 – 212


01

BASIC CONCEPTS OF CHEMISTRY AND CHEMICAL CALCULATIONS

CHAPTER SNAPSHOT

- **1.1** Chemistry the Centre of Life
- **1.2** Classification of Matter
 - **1.2.1** Physical Classification of Matter
 - 1.2.2 Chemical Classification
- 1.3 Atomic and Molecular Masses
 - **1.3.1** Atomic Masses
 - **1.3.2** Molecular Mass
- **1.4** Mole Concept
 - **1.4.1** Avogadro Number
 - **1.4.2** Molar Mass
 - **1.4.3** Molar Volume
- **1.5** Gram Equivalent Concept
 - **1.5.1** Equivalent Mass of Acids, Bases, Salts, Oxidising Agents and Reducing Agents.

- **1.6** Empirical Formula and Molecular Formula
 - 1.6.1 Determination of Empirical Formula from Elemental Analysis Data
 - 1.6.2 Calculation of Molecular Formula from Empirical Formula
- **1.7** Stoichiometry
 - **1.7.1** Stoichiometric Calculations
 - **1.7.2** Limiting Reagents
- 1.8 Redox Reactions
 - 1.8.1 Oxidation Number
 - **1.8.2** Types of Redox Reactions
 - **1.8.3** Balancing (the equation) of Redox Reactions



FORMULAE TO REMEMBER

- * Atomic mass = $\frac{\text{Mass of an atom}}{\left(\frac{1}{12}\right) \times \text{mass of carbon atom}} = \frac{12}{12} \text{ Mass of an atom}$
- * Molecular Mass = n × Vapour Density
- * Molar mass = $\frac{\text{Mass}}{\text{Mole}}$
- * Molecular Formula = n × Empirical Formula
- * Mass % of an element = $\frac{\text{Mass of that element in the compound}}{\text{Molar mass of the compound}} \times 100$
- * Equivalent Mass of Acid = $\frac{\text{Molar mass of the Acid}}{\text{Basicity of Acid}}$
- * Equivalent Mass of Base = $\frac{\text{Molar mass of the Base}}{\text{Acidity of Base}}$
- * Molarity = $\frac{\text{No. of moles of solute}}{\text{Volume of solution in litres}}$
- * Molality = $\frac{\text{No. of moles of solute}}{\text{Mass of solvent in Kg}}$
- * Normality = $\frac{\text{No. of gram equivalents of solute}}{\text{Volume of solution in litres}}$

Mole fraction of A =
$$\frac{\text{No. of moles of A}}{\text{Total no. of moles in solution}} = \frac{\text{nA}}{\text{nA} + \text{nB}}$$

Mole fraction of B =
$$\frac{nB}{nA + nB}$$

MUST KNOW DEFINITIONS

Matter: Matter is defined as anything that has mass and occupies space. All matter is composed of atoms.

Mixtures : Mixtures consist of more than one chemical entity present without any chemical interactions.

Pure substances: Pure substances are composed of simple atoms or molecules. They are further classified as elements and compounds.

: An element consists of only one type of atom.

Element can exist as monatomic or polyatomic units.

PH: 8124201000 / 8124301000

Element

,		
Compound	:	Compounds are made up of molecules which contain two or more atoms of different elements.
Relative atomic mass	:	The relative atomic mass is defined as the ratio of the average atomic mass to the unified atomic mass unit.
Relative molecular mass	:	Relative molecular mass is defined as the ratio of the mass of a molecule to the unified atomic mass unit. The relative molecular mass of any compound can be calculated by adding the relative atomic masses of its constituent atoms.
Mole	:	One mole is the amount of substance that contains as many elementary particles as the number of atoms in 12 g of carbon-12 isotope.
Avogadro Number	:	The total number of entities present in one mole of any substance is equal to 6.022×10^{23} . This number is called Avogadro number
Molar Mass	:	Molar mass is defined as the mass of one mole of a substance. The molar mass of a compound is equal to the sum of the relative atomic masses of its constituents expressed in $g \text{ mol}^{-1}$.
Molar Volume	:	The volume occupied by one mole of any substance in the gaseous state at a given temperature and pressure is called molar volume.
Gram equivalent mass	:	Gram equivalent mass is defined as the mass of an element, (compound or ion) that combines or displaces 1.008 g hydrogen or 8 g oxygen or 35.5 g chlorine.
Empirical formula	:	Empirical formula of a compound is the formula written with the simplest ratio of the number of different atoms present in one molecule of the compound as subscript to the atomic symbol.
Molecular formula	:	Molecular formula of a compound is the formula written with the actual number of different atoms present in one molecule as a subscript to the atomic symbol.
Stoichiometry	:	Stoichiometry is the quantitative relationship between reactants and products in a balanced chemical equation in moles. The quantity of reactants and products can be expressed in moles or in terms of mass unit or as volume.
Limiting reagent	:	When a reaction is carried out using non-stoichiometric quantities of the reactants, the product yield will be determined by the reactant that is completely consumed. It limits the further reaction from taking place and is called as the limiting reagent.
Oxidation Number		It is defined as the imaginary charge left on the atom when all other atoms of the compound have been removed in their usual oxidation states that are assigned according to set of rules.
Combination reactions	:	Redox reactions in which two substances combine to form a single compound are called combination reaction.
Decomposition reaction	:	Redox reactions in which a compound breaks down into two or more components are called decomposition reactions. These reactions are opposite to combination reactions.
Displacement reactions	:	Redox reactions in which an ion (or an atom) in a compound is replaced by an ion (or atom) of another element are called displacement reactions.
Disproportionation reaction	:	In some redox reactions, the same compound can undergo both oxidation and reduction. In such reactions, the oxidation state of one and the same element is both increased and decreased. These reactions are called disproportionation reactions.

Oxidation	- :	Classical concept - Addition of oxygen (or) Removal of hydrogen.
1		Electronic concept - Loss of electrons (or) Increase in oxidation number.
Reduction	:	Classical concept - Addition of Hydrogen (or) Removal of oxygen.
		Electronic concept - Gain of electrons (or) Decrease in oxidation number
Redox Reaction	:	The reaction that involve the oxidation and reduction as its two half reactions are called redox reactions.
Oxidising Agent	:	Classical Concept: In a redox reaction, the substance which oxidises the other (or) reduces itself is called oxidising agent.
		Electron Transfer concept: The substance that gains electrons.
Reducing Agent	:	Classical Concept: In a redox reaction, the substance which reduces the other (or) oxidises itself is called reducing agent.
		Electron Transfer concept : The substance that loss or donate electrons.

EVALUATION

T. Choose The Best Answer:

- 40 ml of methane is completely burnt using 80 ml of oxygen at room temperature The volume of gas left after cooling to room temperature is
 - (a) 40 ml CO₂ gas
 - (b) 40 ml CO₂ gas and 80 ml H₂O gas
 - (c) 60 ml CO₂ gas and 60 ml H₂O gas
 - (d) 120 ml CO₂ gas

[Ans. (a) 40 ml CO, gas]

Hint:
$$CH_{4(g)} + 2O_{2(g)} \longrightarrow CO_{2(g)} + 2H_2O_{(l)}$$

- An element X has the following isotopic composition $^{200}X = 90\%$, $^{199}X = 8\%$ and $^{202}X = 2\%$. The weighted average atomic mass of the element X is closest to
- (a) 201 u (b) 202 u (c) 199 u
- (d) 200 u

[Ans. (d) 200 u]

Hint: =
$$\frac{(200 \times 90) + (199 \times 8) + (202 \times 2)}{100}$$

= $199.96 = 200u$

- 3. **Assertion: Two mole of glucose contains** 12.044×10^{23} molecules of glucose
 - : Total number of entities present in Reason one mole of any substance is equal to

 6.02×10^{22} [First Mid-'18]

- (a) both assertion and reason are true and the reason is the correct explanation of assertion
- (b) both assertion and reason are true but reason is not the correct explanation of assertion
- (c) assertion is true but reason is false
- (d) both assertion and reason are false

[Ans. (c) assertion is true but reason is false]

Hint: Based on Avogadro's law. One mole of any substance is equal to 6.022×10^{23} .

- Carbon forms two oxides, namely carbon monoxide and carbon dioxide. The equivalent mass of which element remains constant?
 - (a) Carbon
- (b) oxygen
- [CRT-'22]
- (c) both carbon and oxygen
- (d) neither carbon nor oxygen [Ans. (b) oxygen]

Hint: React 1: $2C + O_2 \longrightarrow 2CO$

2 × 12g carbon combines with 32g of oxygen

 \therefore Equivalent mass of carbon $= \frac{2 \times 12}{32} \times 8 = 6$

React 2 : $C + O_2 \longrightarrow CO_2$

12g carbon combines with 32g of oxygen

PH: 8124201000 / 8124301000

 $\therefore \text{ Equivalent mass of carbon } \frac{12}{32} \times 8 = 3$

5. The equivalent mass of a trivalent metal element is 9 g eq⁻¹ the molar mass of its anhydrous oxide is

[QY-'24]

- (a) 102 g (b) 27 g
- (c) 270 g (d) 78 g

[Ans. (a) 102 g]

Hint: Atomic mass of the metal oxide is equal to 2 multiple atomic mass of metal + 3 multiple atomic mass of oxygen

6. The number of water molecules in a drop of water weighing 0.018 g is [First Mid-'18; July-'23; Mar-'24]

- (a) 6.022×10^{26}
- (b) 6.022×10^{23}
- (c) 6.022×10^{20}
- (d) 9.9×10^{22}

[Ans. (c) 6.022×10^{20}]

Hint: $0.001 \times 6.022 \times 10^{23}$

- 1 g of an impure sample of magnesium carbonate (containing thermally decomposable impurities) on complete thermal decomposition gave 0.44 g of carbon dioxide gas. The percentage of impurity in the sample is
 - (a) 0%
- (b) 4.4%
- (c) 16%
- (d) 8.4 %

[Ans. (c) 16%]

Hint: impurity is equal to $1 \times 100/1.84$.

- When 6.3 g of sodium bicarbonate is added to 30 g of acetic acid solution, the residual solution is found to weigh 33 g. The number of moles of carbon dioxide released in the reaction is
 - (a) 3
- (b) 0.75
- (c) 0.075(d) 0.3

[Ans. (c) 0.075]

Hint: Number of moles of CO₂ is equal to given weight/ molecular weight.

- When 22.4 litres of H_2 (g) is mixed with 11.2 litres of Cl₂(g), each at 273 K at 1 atm the moles of HCl (g), formed is equal to

 - (a) 2 moles of HCl (g) (b) 0.5 moles of HCl (g)
 - (c) 1.5 moles of HCl (g) (d) 1 moles of HCl (g)

[Ans. (d) 1 moles of HCl (g)]

Hint: $H_2(g) + Cl_2(g) \longrightarrow 2HCl$

1 mole of an ideal gas occupies at 22.4 *l*.

10. Hot concentrated sulphuric acid is a moderately strong oxidising agent. Which of the following reactions does not show oxidising behaviour?

(a)
$$Cu + 2H_2SO_4 \longrightarrow CuSO_4 + SO_2 + 2H_2O$$

- (b) C+ $2H_2SO_4 \longrightarrow CO_2 + 2SO_2 + 2H_2O$
- (c) $BaCl_2 + H_2SO_4 \longrightarrow BaSO_4 + 2HCl$
- (d) none of the above

[Ans. (c) BaCl₂ + H₂SO₄ \longrightarrow BaSO₄ + 2HCl]

- 11. Choose the disproportionation reaction among the following redox reactions.
 - (a) $3Mg_{(s)} + N_{2(g)} \longrightarrow Mg_3N_{2(s)}$
 - (b) $P_{4(s)} + 3 \text{ NaOH} + 3H_2O \longrightarrow PH_{3(g)} +$ 3NaH₂PO_{2(aq)}
 - (c) $Cl_{2(g)} + 2KI_{(aq)} \longrightarrow 2KCl_{(aq)} + I_2$
 - (d) $Cr_2O_{3(s)} + 2Al_{(s)} \longrightarrow Al_2O_{3(s)} + 2Cr_{(s)}$
 - [Ans. (b) $P_{4(s)}$ + 3 NaOH + $3H_2O$ \longrightarrow $PH_{3(g)} + 3NaH_2PO_{2(aq)}$]
- 12. The equivalent mass of potassium permanganate in alkaline medium is

$$MnO_4^- + 2H_2O + 3e^- \longrightarrow MnO_2 + 4OH^-$$

- (a) 31.6
- (b) 52.7

(c) 79

(d) None of these

[Ans. (b) 52.7]

Hint: The reduction reaction of the oxidising agent (MnO₄) involves gain of 3 electrons.

Hence the equivalent mass =

$$\frac{\text{Molar mass of KMnO}_4}{3} = \frac{158.1}{3} = 52.7.$$

- 13. Which one of the following represents 180g of [Qv-'19; Sep-'21]
 - (a) 5 Moles of water
- (b) 90 moles of water
- (c) $\frac{6.022 \times 10^{23}}{180}$ molecules of water
- (d) 6.022×10^{24} molecules of water

[Ans. (d) 6.022×10^{24} molecules of water]

Hint: $10 \times 6.022 \times 10^{23}$

14. 7.5 g of a gas occupies a volume of 5.6 litres at 0° C and 1 atm pressure. The gas is

[Hy-'18; May-'22; Qy-'23]

- (a) NO
- (b) N_2O
 - (c) CO
- (d) CO₂

[Ans. (a) NO]

Hint:
$$\frac{7.5g}{5.6l} \times 22.4l = 30g$$

Molar mass of NO (14 + 16) = 30g.

- **15.** Total number of electrons present in 1.7 g of ammonia is [First Mid-'18; Aug-'22; Hy-'23]
 - (a) 6.022×10^{23}
- (b) $\frac{6.022 \times 10^{22}}{1.7}$
- (c) $\frac{6.022 \times 10^{24}}{1.7}$
- (d) $\frac{6.022 \times 10^{23}}{1.7}$

[Ans. (a) 6.022×10^{23}]

Hint: Number of moles is equal to Atomic weight / valency

16. The correct increasing order of the oxidation state of sulphur in the anions

$$SO_4^{2-}$$
, SO_3^{2-} , $S_2O_4^{2-}$, $S_2O_6^{2-}$ is

- (a) $SO_3^{2-} < SO_4^{2-} < S_2O_4^{2-} < S_2O_6^{2-}$
- (b) $SO_4^{2-} < S_2O_4^{2-} < S_2O_6^{2-} < SO_3^{2-}$
- (c) $S_2O_4^{2-} < SO_3^{2-} < S_2O_6^{2-} < SO_4^{2-}$
- (d) $S_2O_6^{2-} < S_2O_4^{2-} < SO_4^{2-} < SO_3^{2-}$

[Ans. (c)
$$S_2O_4^{2-} < SO_3^{2-} < S_2O_6^{2-} < SO_4^{2-}$$
]

Hint:
$$S_2^{+3}O_4^{2-} < SO_3^{2-} < S_2^{+5}O_6^{2-} < SO_4^{2-}$$

- 17. The equivalent mass of ferrous oxalate is
 - (a) $\frac{\text{molar mass of ferrous oxalate}}{1}$
 - (b) $\frac{\text{molar mass of ferrous oxalate}}{2}$
 - (c) $\frac{\text{molar mass of ferrous oxalate}}{3}$
 - (d) none of these

[Ans. (c)] molar mass of ferrous oxalate
3

Hint:
$$FeC_2^{2+3+}O_4$$
 Oxidising $\rightarrow Fe^{3+} + CO_2^{4+}$
 $n = 1 + 2(1) = 3$

- 18. If Avogadro number were changed from 6.022×10^{23} to 6.022×10^{20} , this would change
 - (a) the ratio of chemical species to each other in a balanced equation
 - (b) the ratio of elements to each other in a compound
 - (c) the definition of mass in units of grams
 - (d) the mass of one mole of carbon

[Ans. (d) the mass of one mole of carbon]

- 19. Two 22.4 litre containers A and B contains 8 g of O₂ and 8 g of SO₂ respectively at 273 K and 1 atm pressure, then
 - (a) Number of molecules in A and B are same
 - (b) Number of molecules in B is more than that in A.
 - (c) The ratio between the number of molecules in A to number of molecules in B is 2:1
 - (d) Number of molecules in B is three times greater than the number of molecules in A.

[Ans. (c) The ratio between the number of molecules in A to number of molecules in B is 2:1]

- **20.** What is the mass of precipitate formed when 50 ml of 8.5 % solution of AgNO₃ is mixed with 100 ml of 1.865 % potassium chloride solution?
 - (a) 3.59 g (b) 7 g
- (c) 14 g
- (d) 28 g

[Ans. (a) 3.59 g]

Hint: Mass of AgNO₃ is equal to number of moles multiple molar mass.

- 21. The mass of a gas that occupies a volume of 612.5 ml at room temperature and pressure (25° C and 1 atm pressure) is 1.1 g. The molar mass of the gas is
 - (a) 66.25 g mol⁻¹
- (b) 44 g mol⁻¹
- (c) 24.5 g mol⁻¹
- (d) 662.5 g mol⁻¹

[Ans. (b) 44 g mol^{-1}]

Hint:
$$= \frac{612.5 \times 10^{-3} l}{24.5 \text{L mol}^{-1}} = -0.025 \text{ moles}$$

$$\text{Molar mass} = \frac{\text{mass}}{\text{No. of. moles}}$$

$$= \frac{1.1 \text{ g}}{0.025 \text{ mol}} = 44 \text{ g mol}^{-1}.$$

- **22.** Which of the following contain same number of carbon atoms as in 6 g of carbon-12.
 - (a) 7.5 g ethane
- (b) 8 g methane
- (c) both (a) and (b)
- (d) none of these

[Ans. (c) both (a) and (b)]

- 23. Which of the following compound(s) has/have percentage of carbon same as that in ethylene (C_2H_4) [Qy-'19; Sep-'21; Mar-'23]
 - (a) propene
- (b) ethyne
- (c) benzene
- (d) ethane

[Ans. (a) propene]

enquiry@surabooks.com

24. Which of the following is/are true with respect to carbon -12.

- (a) relative atomic mass is 12 u
- (b) oxidation number of carbon is +4 in all its compounds.
- (c) 1 mole of carbon-12 contain 6.022×10^{22} carbon atoms.
- (d) all of these

[Ans. (a) relative atomic mass is 12 u]

25. Which one of the following is used as a standard for atomic mass. [Govt. MQP-'18; July-'24]

(a) $_{6}C^{12}$

(b) $_{7}C^{12}$

(c) $_{6}C^{13}$

(d) $_{6}C^{14}$

[Ans. (a) $_{6}C^{12}$]

II. WRITE BRIEF ANSWER TO THE FOLLOWING QUESTIONS.

26. Define relative atomic mass. [First Mid-'18; July-'24]

Ans. The relative atomic mass is defined as the ratio of the average atomic mass to the unified atomic mass unit. Relative atomic mass (A_r)

 $= \frac{\text{Average mass of the atom}}{\text{Unified atomic mass}}$

27. What do you understand by the term mole?

[June-'19; CRT-'22; July -'23 & '24; Hy-'23]

- **Ans.** The mole is defined as the amount of a substance which contains 6.022×10^{23} particles such as atoms, molecules or ions. It is denoted by the symbol "n".
- 28. Define equivalent mass.

- **Ans.** Gram equivalent mass is defined as the mass of an element (compound or ion) that combines or displaces 1.008g hydrogen or 8g oxygen or 35.5g chlorine.
- 29. What do you understand by the term oxidation number?
- **Ans.** Oxidation number refers to the number of charges an atom would have in a molecule or an ionic compound, if electrons were transferred completely. The oxidation numbers reflect the number of electrons transferred.

30. Distinguish between oxidation and reduction.

[Hy-'19; Sep-'21; Mar-'23]

Ans.

	Oxidation	Reduction
(i)	Addition of oxygen and removal of hydrogen	Additional of hydrogen and removal of oxygen
(ii)	This process involves loss of electrons $Fe^{2+} \longrightarrow Fe^{3+} + e^{-}$	This process involves gain electrons. $Cu^{2+} + 2^{e-} \longrightarrow Cu$
(iii)	Oxidation number increases	Oxidation number decreases
(iv)	$Ca + S \longrightarrow Ca^{2+} + 2e^{-}$	$Zn^{2+} + 2e^{-} \longrightarrow Zn$
(v)	Removal of Metal $2KI + H_2O_2 \longrightarrow 2KOH + I_2$	Addition of metal $HgCl_2 + Hg \longrightarrow Hg_2Cl_2$

- 31. Calculate the molar mass of the following compounds.
 - i) Urea [CO(NH₂)₂]
 - ii) Acetone [CH₃COCH₃]
 - iii) Boric acid [H₃BO₃]
 - iv) Sulphuric acid [H₂SO₄]

Ans. (i) urea $[CO(NH_2)_2]$:

 $C : 1 \times 12.01 = 12.01$ $O : 1 \times 16 = 16.00$ $N : 2 \times 14.01 = 28.02$ $H : 4 \times 1.01 = 4.04$

 $\frac{4 \times 1.01}{60.07} = \frac{4.04}{60.07} \text{ g mol}^{-1}$

(ii) acetone [CH₃COCH₃]

 $C : 3 \times 12.01 = 36.03$

 $H : 6 \times 1.01 = 6.06$

O: $1 \times 16 = \frac{16.00}{58.09} \text{ g mol}^{-1}$

(iii) boric acid [H₃BO₃]:

 $H : 3 \times 1.01 = 3.03$

 $B : 1 \times 10 = 10.00$

O: $3 \times 16 = 48.00 \over 61.03$

 $\overline{61.03} \text{ g mol}^{-1}$

(iv) sulphuric acid [H₂SO₄]:

 $H : 2 \times 1.01 = 2.02$

 $S: 1 \times 32.06 = 32.06$

 $O: 4 \times 16 = 64.00$

98.08 g mol⁻¹

32. The density of carbon dioxide is equal to $1.965 \, \mathrm{kgm^{-3}}$ at 273 K and 1 atm pressure. Calculate the molar mass of CO_2 .

Ans. Given:

The density of CO_2 at 273 K and 1 atm pressure = 1.965 kgm⁻³

Molar mass of $CO_2 = ?$

At 273 K and 1 atm pressure, 1 mole of CO_2 occupies a volume of 22.4 L

Mass of 1 mole of CO,

$$= \frac{1.965 \text{Kg}}{\text{lm}^3} \times 22.4 \text{L}$$

$$= \frac{1.965 \times 10^3 \text{ g} \times 22.4 \times 10^{-3} \text{ m}^3}{\text{lm}^3}$$

$$= 44.01 \text{ g}$$

Molar mass of $CO_2 = 44 \text{ gmol}^{-1}$.

- **33.** Which contains the greatest number of moles of oxygen atoms
 - i) 1 mol of ethanol
 - ii) 1 mol of formic acid
 - iii) 1 mol of H,O
- Ans. (i) 1 mol of ethanol: C_2H_5OH (ethanol) Molar mass = 24 + 6 + 16 = 4646g of ethanol contains $1 \times 6.022 \times 10^{23}$ number of oxygen atoms.

- (ii) 1 mol of formic acid: HCOOH (Formic acid)
 Molar mass = 2 + 12 + 32 = 46
 46g of HCOOH contains 2 × 6.022 × 10²³
 number of oxygen atoms
- (iii) 1 mol of H_2O : H_2O (Water) Molar mass = 2 + 16 = 1818g of water contains $1 \times 6.022 \times 10^{23}$ number of oxygen atoms.
- .. mol of formic acid contains the greatest number of oxygen atoms.
- **34.** Calculate the average atomic mass of naturally occurring magnesium using the following data

Isotope	Isotopic atomic mass	Abundance (%)
Mg^{24}	23.99	78.99
Mg^{25}	24.99	10.00
Mg^{26}	25.98	11.01

Ans. Isotopes of Mg

Atomic mass =
$$Mg^{24} = 23.99 \times \frac{78.99}{100} = 18.95$$

Atomic mass = $Mg^{25} = 24.99 \times \frac{10}{100} = 2.499$
Atomic mass = $Mg^{26} = 25.98 \times \frac{11.01}{100} = 2.860$
Average atomic mass = 24.309

PH: 8124201000 / 8124301000

Average atomic mass of Mg = 24.31u.

- **35.** In a reaction $x + y + z_2 \longrightarrow xyz_2$ identify the Limiting reagent if any, in the following reaction mixtures.
 - (a) 200 atoms of x + 200 atoms of y + 50 molecules of z_2
 - (b) 1 mol of x + 1 mol of y + 3 mol of z_2
 - (c) 50 atoms of x + 25 atoms of y + 50 molecules of z_2
 - (d) 2.5 mol of x + 5 mol of y + 5 mol of z_2

Ans. Reaction: $x + y + z_2 \longrightarrow xyz_2$

Questio		r of moles of allowed to rea			r of moles of med during r		Limiting
	X	y	\mathbf{z}_2	X	y	\mathbf{z}_2	reagent
(a)	200 atoms	200 atoms	50 molecules	50 atoms	50 atoms	50 molecules	z_2
(b)	1 mol	1 mol	3 mol	1 mol	1 mol	1 mol	x and y
(c)	50 atom	25 atom	50 molecules	25 atom	25 atom	25 molecules	у
(d)	2.5 mol	5 mol	5 mol	2.5 mol	2.5 mol	2.5 mol	Х

36. Mass of one atom of an element is 6.645×10^{-23} g. How many moles of element are there in 0.320 kg.

Ans. Given:

mass of one atom = 6.645×10^{-23} g

∴ mass of 1 mole of atom

$$= 6.645 \times 10^{-23} \text{ g} \times 6.022 \times 10^{23}$$

$$= 40 g$$

:. number of moles of element in 0.320 kg

$$= \frac{1 \text{ mole}}{40 \text{g}} \times 0.320 \text{kg}$$

$$= \frac{1 \text{ mol} \times 320g}{40g}$$

= 8 mol

37. What is the difference between molecular mass and molar mass? Calculate the molecular mass and molar mass for carbon monoxide.

Ans.

	Molecular mass	Molar mass
(i)	Molecular mass is defined as the ratio of the mass of a molecule to the unified atomic mass unit.	Molar mass is defined as the mass of one mole of a substance.
(ii)	The relative molecular mass of any compound is calculated by adding the relative atomic masses of its constituent atoms	The molar mass of a compound is equal to the sum of the relative atomic masses of its constituents.
(iii)	Its unit is u or amu	Its unit is g mol ⁻¹
(iv)	Molecular mass of CO: (1 × at.mass of C) + (1 × at.mass of O) 1 × 12.01 amu + 1 × 16 amu = 28.01 amu	Molar mass of CO: 1 × 12.01 + 1 × 16 = 28.01 g mol ⁻¹

- 38. What is the empirical formula of the following?
 - i) Fructose (C₆H₁₂O₆) found in honey
 - ii) Caffeine (C₈H₁₀N₄O₂) a substance found in tea and coffee.

[First Mid-'18; Qy-'18; Sep-'21; July-'24]

Ans.

Compound	Molecular formula	Empirical formula
Fructose	$C_6 H_{12} O_6$	CH ₂ O
Caffeine	$C_8 H_{10} N_4 O_2$	$C_4 H_5 N_2 O$

- 39. The reaction between aluminium and ferric oxide can generate temperatures up to 3273 K and is used in welding metals. (Atomic mass of Al = 27 u Atomic mass of O = 16 u)
 - $2Al + Fe_2O_3 \longrightarrow Al_2O_3 + 2Fe$; If, in this process, 324 g of aluminium is allowed to react with 1.12 kg of ferric oxide.
 - i) Calculate the mass of Al₂O₃ formed.
 - ii) How much of the excess reagent is left at the end of the reaction? [Govt. MQP-'18]

Ans. (i)
$$2Al + Fe_2O_3 \longrightarrow Al_2O_3 + 2Fe$$

 $54g \quad 160g \quad 102g \quad 112g$

As per balanced equation 54g Al is required for 112g of Iron and 102g of Al₂O₃.

$$\therefore$$
 324g of Al will give $\frac{102}{54} \times 324 = 612g$ of Al₂O₃

(ii) 54g of Al required 160g of Fe₂O₃ for welding reaction

$$\therefore 324g \text{ of Al will require } \frac{160}{54} \times 324 = 960g \text{ of }$$
Fe₂O₃

$$\therefore \text{ Excess Fe}_2\text{O}_3 \text{ - unreacted Fe}_2\text{O}_3$$
$$= 1120 - 960 = 160\text{g}.$$

∴ 160 g of excess reagent is left at the end of the reaction.

40. How many moles of ethane is required to produce 44 g of $CO_{2(g)}$ after combustion.

[First Mid-'18; Qy.- '19]

Ans. Balanced equation for the combustion of ethane

$$C_2H_6 + \frac{7}{2}O_2 \longrightarrow 2CO_2 + 3H_2O$$

$$\Rightarrow$$
 2C₂H₆ + 7O₂ \longrightarrow 4CO₂ + 6H₂O

To produce 4 moles of CO₂, 2 moles of ethane is required

∴ To produce 1 mole (44 g) of CO₂ required number of moles of ethane

$$= \frac{2 \text{ mol ethane}}{4 \text{ mol-} CQ_2} \times 1 \text{ mol-} CQ_2$$

$$=\frac{1}{2}$$
 mole of ethane

= 0.5 mole of ethane.

41. Hydrogen peroxide is an oxidising agent. It oxidises ferrous ion to ferric ion and reduced itself to water. Write a balanced equation.

Ans.

42. Calculate the empirical and molecular formula of a compound containing 76.6% carbon, 6.38 % hydrogen and rest oxygen its vapour density is 47. [Mar-'19; Sep-2020; Aug-'22; Qy-'23]

Ans.

Element	Percentage	Atomic mass	Relative number of atoms	Simple ratio	Whole no
С	76.6	12	$\frac{76.6}{12} = 6.38$	$\frac{6.38}{1.06} = 6$	6
Н	6.38	1	$\frac{6.38}{1} = 6.38$	$\frac{6.38}{1.06} = 6$	6
О	17.02	16	$\frac{17.02}{16} = 1.06$	$\frac{1.06}{1.06} = 1$	1

Empirical formula =
$$C_6 H_6 O$$

n = $\frac{Molar \ mass}{Calculated \ empirical \ formula \ mass}$
= $\frac{2 \times vapour \ density}{94} = \frac{2 \times 47}{94} = 1$, [since Molar mass = 2 × Vapour density]

PH: 8124201000 / 8124301000

Molecular formula = $n \times empirical formula$

- :. Molecular formula = $(C_6H_6O) \times 1 = C_6H_6O$.
- **43.** A Compound on analysis gave Na = 14.31% S = 9.97% H = 6.22% and O = 69.5% calculate the molecular formula of the compound, if all the hydrogen in the compound is present in combination with oxygen as water of crystallization. (molecular mass of the compound is 322).

 [Mar-'23]

Ans.

Element	%	Relative number of atoms	Simple ratio		
Na	14.31	$\frac{14.31}{23} = 0.62$	$\frac{0.62}{0.31} = 2$		
S	9.97	$\frac{9.97}{32} = 0.31$	$\frac{0.31}{0.31} = 1$		
Н	6.22	$\frac{6.22}{1} = 6.22$	$\frac{6.22}{0.31} = 20$		
О	69.5	$\frac{69.5}{16} = 4.34$	$\frac{4.34}{0.31} = 14$		

[First Mid-'18]

$$\therefore \qquad \text{Empirical formula} = \operatorname{Na_2} \operatorname{SH_{20}O_{14}}$$

$$n = \frac{\operatorname{Molecular \ mass}}{\operatorname{Empirical \ formula \ mass}} = \frac{322}{322} = 1$$

$$\operatorname{Empirical \ formula \ mass} = (23 \times 2) + (32 \times 1) + (20 \times 1) + (14 \times 6)$$

$$= 46 + 32 + 20 + 224 = 322$$

$$\operatorname{Molecular \ formula} = \operatorname{Na_2} \operatorname{SH_{20}O_{14}}$$

Since all the hydrogen in the compound present as water.

∴ Molecular formula is Na₂SO₄.10H₂O

44. Balance the following equations by oxidation number method

i)
$$K_2Cr_2O_7 + KI + H_2SO_4 \longrightarrow K_2SO_4 + Cr_2(SO_4)_3 + I_2 + H_2O_4$$

ii)
$$\text{KMnO}_4 + \text{Na}_2\text{SO}_3 \longrightarrow \text{MnO}_2 + \text{Na}_2\text{SO}_4 + \text{KOH}$$

iii)
$$Cu + HNO_3 \longrightarrow Cu(NO_3)_2 + NO_2 + H_2O$$

iv)
$$KMnO_4 + H_2C_2O_4 + H_2SO_4 \longrightarrow K_2SO_4 + MnSO_4 + CO_2 + H_2O$$

Ans. (i) $K_2 Cr_2 O_7 + K I + H_2 SO_4 \longrightarrow K_2 SO_4 + Cr_2 (SO_4)_3 + I_2 + H_2 O_4$

$$\begin{split} & \textbf{K}_2\textbf{Cr}_2\textbf{O}_7 + \textbf{6}\textbf{K}\textbf{I} + \textbf{H}_2\textbf{SO}_4 & \longrightarrow \textbf{K}_2\textbf{SO}_4 + \textbf{Cr}_2(\textbf{SO}_4)_3 + \textbf{I}_2 + \textbf{H}_2\textbf{O} \\ & \textbf{K}_2\textbf{Cr}_2\textbf{O}_7 + \textbf{6}\textbf{K}\textbf{I} + \textbf{H}_2\textbf{SO}_4 & \longrightarrow \textbf{K}_2\textbf{SO}_4 + \textbf{Cr}_2(\textbf{SO}_4)_3 + \textbf{3I}_2 + \textbf{H}_2\textbf{O} \\ & \textbf{K}_2\textbf{Cr}_2\textbf{O}_7 + \textbf{6}\textbf{K}\textbf{I} + \textbf{7}\textbf{H}_2\textbf{SO}_4 & \longrightarrow \textbf{4}\textbf{K}_2\textbf{SO}_4 + \textbf{Cr}_2(\textbf{SO}_4)_3 + \textbf{3I}_2 + \textbf{7}\textbf{H}_2\textbf{O} \end{split}$$

(ii)
$$K \stackrel{+7}{M} nO_4 + Na_2 \stackrel{+4}{S} O_3 \longrightarrow \stackrel{+4}{M} nO_2 + Na_2 \stackrel{+6}{S} O_4 + KOH$$

$$\downarrow \qquad \qquad \downarrow \qquad$$

$$2KMnO_4 + 3Na_2SO_3 \longrightarrow MnO_2 + Na_2SO_4 + KOH$$

$$2KMnO_4 + 3Na_2SO_3 \longrightarrow 2MnO_2 + 3Na_2SO_4 + KOH$$

$$2KMnO_4 + 3Na_2SO_3 + H_2O \longrightarrow 2MnO_2 + 3Na_2SO_4 + 2KOH$$

(iii)
$$Cu + HNO_3 \longrightarrow Cu(NO_3)_2 + NO_2 + H_2O$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \qquad \qquad$$

$$\begin{array}{l} \textbf{Cu} + \textbf{2HNO}_3 & \longrightarrow \textbf{Cu}(\textbf{NO}_3)_2 + \textbf{NO}_2 + \textbf{H}_2\textbf{O} \\ \textbf{Cu} + 2\textbf{HNO}_3 + \textbf{2HNO}_3 & \longrightarrow \textbf{Cu}(\textbf{NO}_3)_2 + \textbf{2NO}_2 + 2\textbf{H}_2\textbf{O} \\ \textbf{Cu} + 4\textbf{HNO}_3 & \longrightarrow \textbf{Cu}(\textbf{NO}_3)_2 + 2\textbf{NO}_2 + 2\textbf{H}_2\textbf{O} \end{array}$$

(iv)
$$K \stackrel{+7}{M} \text{nO}_4 + \text{H}_2 \stackrel{+3}{\text{C}}_2 \text{O}_4 + \text{H}_2 \text{SO}_4 \longrightarrow K_2 \text{SO}_4 + \stackrel{+2}{M} \text{nSO}_4 + \stackrel{+4}{\text{C}} \text{O}_2 + \text{H}_2 \text{O}$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad$$

$$\begin{array}{l} \mathbf{2KMnO_4} + \mathbf{5H_2C_2O_4} + \mathrm{H_2SO_4} & \longrightarrow \mathrm{K_2SO_2} + \mathrm{MnSO_4} + \mathrm{CO_2} + \mathrm{H_2O} \\ 2\mathrm{KMnO_4} + 5\mathrm{H_2C_2O_4} + \mathrm{H_2SO_4} & \longrightarrow \mathrm{K_2SO_4} + \mathbf{2MnSO_4} + \mathbf{10CO_2} + \mathrm{H_2O} \\ 2\mathrm{KMnO_4} + 5\mathrm{H_2C_2O_4} + 3\mathrm{H_2SO_4} & \longrightarrow \mathrm{K_2SO_4} + 2\mathrm{MnSO_4} + 10\mathrm{CO_2} + 8\mathrm{H_2O} \end{array}$$

enquiry@surabooks.com PH: 8124201000 / 8124301000

45. Balance the following equations by ion electron method.

i)
$$KMnO_4 + SnCl_2 + HCl \longrightarrow MnCl_2 + SnCl_4 + H_2O + KCl$$

ii)
$$C_2O_4^{2-} + Cr_2O_7^{2-} \longrightarrow Cr^{3+} + CO_2$$
 (in acid medium)

iii)
$$Na_2S_2O_3 + I_2 \longrightarrow Na_2S_4O_6 + NaI$$

iv)
$$Zn + NO_3^- \longrightarrow Zn^{2+} + NO$$
 (in acid medium)

[First Mid-'18]

Ans. (i) Half reaction are:

$$\begin{array}{ccc}
& \stackrel{+7}{\text{M nO}_4^-} \longrightarrow \text{Mn}^{2+} & \dots \dots (1) \\
\text{and} & \text{Sn}^{2+} \longrightarrow \text{Sn}^{4+} & \dots \dots (2) \\
& \text{(1)} \Rightarrow \text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \longrightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}
\end{array}$$

$$(1) \Rightarrow MnO_4 + 8H^4 + 5e \longrightarrow Mn^{-4} + 4H_2C$$

$$(2) \Rightarrow Sn^{2+} \longrightarrow Sn^{4+} + 2e^{-}$$

$$(1) \times 2 \Rightarrow$$
 $2MnO_4^- + 16H^+ + 10e^- \longrightarrow 2Mn^{2+} + 8H_2O$

$$(2) \times 5 \Rightarrow \underbrace{5Sn^{2+} \longrightarrow 5Sn^{4+} + 10e^{-}}_{\Rightarrow 2MnO_{4}^{-} + 5Sn^{2+} + 16H^{+} \longrightarrow 2Mn^{2+} + 5Sn^{4+} + 8H_{2}O}$$

(ii)

$$\overset{+6}{\text{C}} \text{ r}_2 \text{O}_7^{2-} \longrightarrow \text{Cr}^{3+} \qquad \dots (2)$$

$$(1) \Rightarrow \qquad \qquad C_2O_4^{2-} \longrightarrow 2CO_2 + 2e^- \qquad \qquad(3)$$

$$Cr_2O_7^{2-} + 14H^+ \longrightarrow 2Cr^{3+} + 7H_2O$$
(4)

$$3C_2O_4^{2-} \longrightarrow 6CO_2 + 6e^2 \qquad \dots (5)$$

$$3C_{2}O_{4}^{2-} \longrightarrow 6CO_{2} + 6e^{2} \qquad(5)$$

$$Cr_{2}O_{7}^{2-} + 14H^{+} + 6e^{2} \longrightarrow 2Cr^{3+} + 7H_{2}O \qquad(4)$$

$$\Rightarrow Cr_{2}O_{7}^{2-} + 3C_{2}O_{4}^{2-} + 14H^{+} \longrightarrow 2Cr^{3+} + 6CO_{2} + 7H_{2}O$$

(iii)

$$S_2O_3^{2-} \longrightarrow S_4O_6^{2-} \qquad(1)$$
half reaction $\Rightarrow I_2 \longrightarrow I^- \qquad(2)$

eaction
$$\Rightarrow$$
 $I_2 \xrightarrow{} I$ (2)
(1) \Rightarrow $2S_2O_3^{2-} \longrightarrow S_4O_6^{2-} + 2e^{-}$ (3)
(2) \Rightarrow $I_2 + 2e^{-} \longrightarrow 2I^{-}$ (4)
(3) $+$ (4) \Rightarrow $2S_2O_3^{2-} + I_2 \longrightarrow S_4O_6^{2-} + 2I^{-}$

$$(2) \Rightarrow I_2 + 2e^{-} \longrightarrow 2I^{-} \qquad \dots \dots (4)$$

$$(3) + (4) \Rightarrow \underbrace{2S_2O_3^{2-} + I_2 \longrightarrow S_4O_6^{2-} + 2I^{-}}_{6}$$

$$\overset{0}{Z} n \longrightarrow Z n^{2+} \qquad \qquad \dots \dots (1)$$

$$\stackrel{+5}{NO_3} \longrightarrow \stackrel{+2}{NO}$$
(2)

$$(1) \Rightarrow Zn \longrightarrow Zn^{2+} + 2e^{-} \qquad \dots (3)$$

$$(2) \Rightarrow NO_3^- + 3e^- + 4H^+ \longrightarrow NO + 2H_2O \qquad \dots (4)$$

$$(3) \times 3 \Rightarrow 3Zn \longrightarrow 3Zn^{2+} + 6e^{-} \qquad \dots (5)$$

$$(4) \times 2 \Rightarrow 2NO_3^- + 6e^- + 8H^+ \longrightarrow 2NO + 4H_2O \qquad(6)$$

$$3Zn + 2NO_3^- + 8H^+ \longrightarrow 3Zn^{2+} + 2NO + 4H_2O$$

Evaluate Yourself

- 1. By applying the knowledge of chemical classification, classify each of the following into elements, compounds or mixtures.
 - (i) Sugar
- (ii) Sea water
- (iii) Distilled water
- Carbon dioxide
- (v) Copper wire
- (vi) Table salt
- (vii) Silver plate
- (viii) Naphthalene balls
- Ans. (i) Element Copper wire, Silver plate
 - (ii) Compound Sugar, distilled water, carbon dioxide, Table salt, Naphthalene balls

(iv)

- (iii) Mixture Sea water
- 2. Calculate the relative molecular mass of the following.
 - (i) Ethanol(C₂H₅OH)
 - (ii) Potassium permanganate (KMnO₄)
 - (iii) Potassium dichromate (K,Cr,O₇)
 - (iv) Sucrose (C_1, H_2, O_{11})

Ans. (i)
$$C_2H_5OH: (2 \times 12) + (5 \times 1) + (1 \times 16) + (1 \times 1) = 46 g$$

(ii) KMnO₄:
$$(1 \times 39) + (1 \times 55) + (4 \times 16) = 158 \text{ g}$$

(iii)
$$K_2Cr_2O_7$$
: $(2 \times 39) + (2 \times 52) + (7 \times 16) = 294$ g

(iv)
$$C_{12}H_{22}O_{11}: (12 \times 12) + (22 \times 1) + (11 \times 16) = 342 g$$

- 3. a) Calculate the number of moles present in 9 g of ethane.
 - b) Calculate the number of molecules of oxygen gas that occupies a volume of 224 ml at 273 K and 3 atm pressure.
- **Ans.** (a) Molar mass of ethane,

$$C_2H_6 = (2 \times 12) + (6 \times 1) = 30 \text{ g mol}^{-1}$$

No of moles = $\frac{\text{mass}}{\text{molar mass}} = \frac{9g}{30g \text{ mol}^{-1}} = 0.3 \text{ mole}$

(b) At 273 K and 1 atm pressure 1 mole of a gas occupies a volume of 22.4 L

Therefore.

number of moles of oxygen, that occupies a volume of 224 ml at 273 K and 3 atm pressure

$$= \frac{1 \text{ mole}}{273 \text{ K} \times 1 \text{ atm} \times 22.4 \text{ L}} \times 0.224 \text{ L} \times 273 \text{ K} \times 3 \text{ atm}$$

= 0.03 mole

1 mole of oxygen contains 6.022×10^{23} molecules 0.03 mole of oxygen contains = $6.022 \times 10^{23} \times 0.03$

- = 1.807×10^{22} molecules of oxygen
- 4. a) 0.456 g of a metal gives 0.606g of its chloride. Calculate the equivalent mass of the metal.

Ans. Mass of the metal = $W_1 = 0.456g$

Mass of the metal chloride = $W_2 = 0.606g$

- :. Mass of chlorine = $W_2 W_1 = 0.606 0.456 = 0.15g$
- 0.15g of chlorine combine with 0.456g of metal
- :. 35.5g of chlorine will combine with

$$\frac{0.456}{0.15} \times 35.5 = 107.92 \text{g eq}^{-1}$$

 b) Calculate the equivalent mass of potassium dichromate. The reduction half reaction in acid medium is,

$$Cr_2O_7^{-2} + 14H^+ + 6e^- \longrightarrow 2Cr^{3+} + 7H_2O$$

Ans. Equivalent mass of a oxidising agent

Molar mass


No. of moles of electrons gained by one mole of the reducing agent

$$= \frac{294.18 \,\text{mol}^{-1}}{6 \,\text{eq mol}^{-1}} = 49.03$$
$$= 49.0 \,\text{g eq}^{-1}$$

5. Balance the following equation using oxidation number method

$$As_2S_3 + HNO_3 + H_2O \rightarrow H_3AsO_4 + H_2SO_4 + NO$$

Ans.



Equate the total no. of electrons in the reactant side by cross multiplying,

 \Rightarrow 3 As₂ S₃ + 28 HNO₃ \rightarrow H₃AsO₄ + H₂SO₄ + NO Based on reactant side, balance the products

 \Rightarrow 3 As₂ S₃ + 28 HNO₃ \rightarrow 6 H₃AsO₄ + 9 H₂SO₄ + 28 NO **Product side** :36 hydrogen atoms & 88 oxygen atoms **Reactant side** :28 hydrogen atoms & 84 oxygen atoms Difference is 8 hydrogen atoms & 4 oxygen atoms

∴ Add 4 H₂O molecule on the reactant side.

Balanced equation is,

$$3 \text{ As}, S_3 + 28 \text{ HNO}_3 + 4 \text{ H}_2\text{O} \rightarrow 6 \text{ H}_3 \text{ As} \text{ O}_4 + 9 \text{ H}_2 \text{ SO}_4 + 28 \text{ NO}$$

6. A Compound on analysis gave the following percentage composition C=54.55%, H=9.09%, O=36.36%. Determine the empirical formula of the compound.

Ans.

Element	Percentage Composition	Atomic mass	Relative no. of atoms $= \frac{\text{Percentage}}{\text{Atomic mass}}$	Simple ratio	
С	54.55 %	12	54.55/12 = 4.55	4.55 / 2.27 = 2	
Н	9.09 %	1	9.09 / 1 = 9.09	9.09 / 2.27 = 4	
О	36.36 %	16	36.36/16 = 2.27	2.27/2.27 = 1	
Empirical formula (C ₂ H ₄ O)					

- 7. Experimental analysis of a compound containing the elements x,y,z on analysis gave the following data. x = 32 %, y = 24 %, z = 44 %. The relative number of atoms of x, y and z are 2, 1 and 0.5, respectively. (Molecular mass of the compound is 400 g) Find out.
 - i) The atomic masses of the element x,y,z.
 - ii) Empirical formula of the compound and
 - iii) Molecular formula of the compound.

Ans.

Element	Percentage Composition	Relative no. of atoms = Percentage Atomic mass	Atomic mass = Percentage Relative no. of atoms	Simple ratio		
X	32 %	2	16	4		
Y	24 %	1	24	2		
Z	44 %	0.5	88	1		
Empirical formula (X ₄ Y ₂ Z)						

PH: 8124201000 / 8124301000

Calculated empirical formula mass

$$= (16 \times 4) + (24 \times 2) + 88$$

$$= 64 + 48 + 88 = 200$$

$$= \frac{\text{molar mass}}{\text{calculated empirical formula mass}}$$

$$= \frac{400}{200}$$

$$= 2$$

:. Molecular formula $(X_4Y_2Z)_2 = X_8Y_4Z_2$

18

8. The balanced equation for a reaction is given below [CRT-'22]

$$2x+3y \rightarrow 4l + m$$

When 8 moles of x react with 15 moles of y, then

- Which is the limiting reagent? i)
- ii) Calculate the amount of products formed.
- iii) Calculate the amount of excess reactant left at the end of the reaction.

Ans.

Content	Read	etant	Products	
	X	у	l	m
Stoichiometric coefficient	2	3	4	1
No. of moles allowed to react	8	15	_	
No. of moles of reactant reacted and product formed	8	12	16	4
No. of moles of un-reacted reactants and the product formed	_	3	16	4

Limiting reagent: x

Product formed: 16 moles of l & 4 moles of m Amount of excess reactant: 3 moles of v

Government Exam Questions and Answers

CHOOSE THE CORRECT ANSWER 1 MARK

- The equivalent mass of a divalent metal element is 10 g eq⁻¹. The molar mass of its anhydrous oxide [Govt. MQP-'18]
 - (a) 46 g
- (b) 36 g
- (c) 52 g
- (d) none of these

[Ans. (c) 52 g]

Hint: Atomic mass of divalent metal is equal to 2 multiple of atomic mass of metal + 2 multiple of atomic mass of oxygen

Match the list I with List II correctly by using the code given below the list. [Qy-'18]

Lis	List I (no. of moles)		List II (Amount)
A	0.1 mole	1	4480 <i>ml</i> of CO ₂
В	0.2 mole	2	200 mg of hydrogen gas
C	0.25 mole	3	9 ml of water
D	0.5 mole	4	1.51×10^{23} molecules of
			oxygen

- A B C D
- 3 2 4 1 (a)
- **(b)** 2
- 3 2 1 4 (c)

[Ans. (d) 2 1 4 3]

Hint: Number of moles is equal to Mass/ Molar mass Number of moles is equal to Volume/ molar volume

- 3. The oxidation number of chromium in dichromate (ion) is [Qy-'18]
 - (a) +4
- (b) +6
- (c)+5
- (d) 0

[Ans. (b) + 6]

- The empirical formula of glucose is: [Hy-'19]
 - (a) CH₂O (b) CHO
- (c) CH₂O₂ (d) CH₃O₂ [Ans. (a) CH₂O]
- The oxidation number of carbon in CH₂F₂ is _____. [June-'19]
- (a) +4

5.

- (b) -4
- (c) 0
- [Ans. (c) 0]

(d) +2

- The relative molecular mass of ethanol is [Sep-2020]
 - (a) $0.46 \,\mathrm{g}$ (b) $4.6 \,\mathrm{g}$
 - - (c) 460 g
- (d) 46 g [Ans. (d) 46 g]
- Basicity of Sulphuric acid is
- [QY-'24]

- (a) 3
- (b) 1
- (c) 2
- (d) None
- [Ans. (c) 2]

Answer the Questions

2 MARKS

1. Write the electronic concept of oxidation and reduction reactions. [Qy & Hy-'18]

Ans. The process can be explained on the basis of electrons. The reaction involving loss of electron is termed oxidation

$$Fe^{2+} \rightarrow Fe^{3+} + e^{-}$$
 (loss of electron-oxidation).

The reaction involving gain of electron is termed reduction.

 $Cu^{2+} + 2e^{-} \rightarrow Cu$ (gain of electron-reduction)

2. How many moles of hydrogen is required to produce 10 moles of ammonia? [Hy-'18; Qy.-'24]

Ans. (i) The balanced stoichiometric equation for the formation of ammonia is

$$N_2(g) + 3 H_2(g) \longrightarrow 2 NH_3(g)$$

- (ii) To produce 2 moles of ammonia, 3 moles of hydrogen are required
- .. To produce 10 moles of ammonia

$$= \frac{3 \text{ moles of H}_2}{2 \text{ moles of NH}_3} \times 10 \text{ moles of NH}_3$$

= 15 moles of hydrogen are required.

3. Calculate oxidation number of oxygen in H_2O_2 .

[Mar-'19]

Ans. Hydrogen peroxide (H_2O_2)

$$2 (+1) + 2x = 0; \Rightarrow 2x = -2; \Rightarrow x = -1$$

4. What is combination reaction? Give example.

[Hv-'19]

Ans. Redox reactions in which two substances combine to form a single compound are called combination reaction. **Ex:**



5. Calculate the oxidation states of oxygen in H_2O_2 and KO_2 . [Qy-'19]

Ans. Hydrogen peroxide (H_2O_2) is -1.


$$2 (+1) + 2x = 0$$
; $\Rightarrow 2x = -2$; $\Rightarrow x = -1$

Super oxides such as KO_2 is -1/2

$$+1 + 2x = 0$$
; $\Rightarrow 2x = -1$; $\Rightarrow x = -1/2$.

6. Define basicity. Find the basicity of ortho-phosphoric acid. [Sep_2020]

- **Ans.** (i) Basicity: The number of replaceable hydrogen atoms present in a molecule of the acid is referred to as its basicity.
 - (ii) Basicity of ortho-phosphoric acid H₃PO₄



The number of Hydrogen atoms bonded to the oxygen atoms in this compound is 3. Therefore, the basicity of ortho-phosphoric acid is 3.

7. Define Gram equivalent mass.

[May-'22]

Ans. Gram equivalent mass is defined as the mass of an element (compound or ion) that combines or displaces 1.008 g hydrogen or 8 g oxygen or 35.5 g chlorine.

ANSWER THE QUESTIONS

3 MARKS

1. Statement 1: Two mole of glucose contains 12.044×10^{23} molecules of glucose

Statement 2: Total number of entities present in one mole of any substance is equal to 6.02×10^{22} . [Govt. MQP-'18]

Whether the above statements are true? Is there any relation between these two statements?

- **Ans.** The statements 1 & 2 are true. But there is no relation between statement 1 and statement 2.
- 2. Calculate the total number of electrons present in 17g of ammonia. [Govt. MQP-'18]

Ans. No. of electrons present in one ammonia (NH_3) molecule (7 + 3) = 10

No. of moles of NH₃ =
$$\frac{\text{Mass}}{\text{Molar mass}} = \frac{17g}{17g \text{ mol}^{-1}} = 1 \text{ mol}$$

No. of molecules present in 1 mol of NH₃

$$=6.022 \times 10^{23}$$

No. of electrons present in 1 mol of NH₃

$$= 10 \times 6.022 \times 10^{23}$$

$$=6.022 \times 10^{24}$$

enquiry@surabooks.com

3. Calculate the amount of water produced by the combustion of 32 g of methane. [Qy-'18]

Ans.
$$CH_4(g) + 2O_2(g) \rightarrow CO_2(g) + 2H_2O(g)$$

As per the stoichiometric equation,

Combustion of 1 mole (16 g) CH_4 produces 2 moles $(2 \times 18 = 36 \text{ g})$ of water.

$$CH_4$$

(12) + (4×1) = 16 g mol⁻¹

$$H_2O$$
 $(2\times1) + (1\times16) = 18 \text{ g mol}^{-1}$

Combustion of 32 g CH₄ produces

$$\frac{36 \text{ g H}_2\text{O}}{16 \text{ g CH}_4} \times \frac{2}{32 \text{ g CH}_4} = 72 \text{ g of water}$$

4. Calculate the equivalent mass of H₂SO₄.

[Mar.-'19]

Ans.
$$H_2SO_4$$
 basicity = 2eq mol⁻¹
Molar mass of H_2SO_4 = $(2 \times 1) + (1 \times 32) + (4 \times 16)$
= 98 g mol⁻¹
Gram equivalent of H_2SO_4 = $\frac{98}{2}$ = 49 g eq⁻¹

5. $X_2 + 3Y_2 \rightarrow 2XY_3$ In this reaction 2 moles of X_2 and 4.5 moles of Y_2 react to give products. Which is the limiting agent (reagent) and calculate the no. of moles of X_2 , Y_2 and XY_3 in the reaction mixture?

[Qy.-'19]

Ans.
$$X_2 + 3Y_2 \rightarrow 2XY_3$$

No. of moles 2 4.5 ?

SC 1 3 2

ratio 2/1 4.5/3 -
2(ER) 1.5(LR) -

mole-mole $= \frac{nX_2}{1} = \frac{nY_2}{3} = \frac{nXY_3}{2}$
 $= \frac{2}{1} = \frac{4.5}{3} = \frac{nXY_3}{2}$
 $= \frac{4.5}{3} = \frac{nXY_3}{2} = 3$ moles

No. of moles of $2XY_3 = 3$ moles

6. Calculate the oxidation number of underlined elements. (i) CO₂ (ii) H₂SO₄ [May-'22]

Ans. (i) \underline{CO}_2 - X + 2(-2) = 0

$$X = +4$$

(ii)
$$H_2\underline{S}O_4 - 2(+1) + X + 4(-2) = 0$$

 $2 + X - 8 = 0$

$$X = +6$$

- Calculate the oxidation state of the underlined elements in the following compounds. [CRT-'22]
 - (i) \underline{SO}_2 (ii) \underline{CH}_2F_2 (iii) \underline{OF}
- Ans. (i) $\underline{SO}_2 = X+2(-2)0 \Rightarrow x = +4 \Rightarrow S = +4$
 - (ii) $CH_2F_2 = X+2(+1)+2(-1)=0 \Rightarrow x=0 \Rightarrow C=0$
 - (iii) $\underline{O}F_2 = X+2(-1) = 0 \Rightarrow x = +2 \Rightarrow O = +2$

Answer All the Ouestions 5 MARKS

1. Define oxidation number. Balance the following equation using oxidation number method.

$$As_2S_3 + HNO_3 + H_2O \longrightarrow H_3AsO_4 + H_2SO_4 + NO [Govt. MQP-'18]$$

Ans. Oxidation number: It is defined as the imaginary charge left on the atom when all other atoms of the compound have been removed in their usual oxidation states that are assigned according to set of rules.

Solution:

Step 1 :
$$As_2S_3 + HNO_3 + H_2O \longrightarrow H_3AsO_4 + H_2SO_4 + NO$$

Step 2: Assign the oxidation numbers and identify the redox couples.

Decreased by 3

$$+3 -2 +1 +5 -2 +1 -2$$
 $As_2S_3 + H N O_3 + H_2O$
 $+1 +5 -2 +1 +6 -2 +2 -2$
 $H_3AsO_4 + H_2SO_4 + NO$
Increased by 2

Step 3: Multiply As₂S₃ by 3 and HNO₃ by 2.

$$3As_2S_3 + 2HNO_3 + H_2O \longrightarrow H_3AsO_4 + H_2SO_4 + NO$$

Step 4 : Balance all the elements in the equation (As, S and N) except H and O.


$$3As_2S_3 + 2HNO_3 + H_2O \longrightarrow 6H_3AsO_4 + 9H_2SO_4 + 2NO$$

Step 5 : Balance the complete equation including O & H.

2. What are disproportionation reaction. (or) What | 5. are auto redox reactions? Give an example.

[June-'19]

Ans. Disproportionation reaction (Auto redox reactions): In some redox reactions, the same compound can undergo both oxidation



and reduction. In such reactions, the oxidation state of one and the same element is both increased and decreased. These reactions are called disproportionation reactions.

3. Define limiting reagent.

Ans. Limiting reagent: When a reaction is carried out using non-stoichiometric quantities of the reactants, the product yield will be determined by the reactant that is completely consumed. It limits the further reaction from taking place and is called as the limiting reagent.

4. What is disproportionation reaction? Give example. [Qy. - '23]

Ans. The reaction in which an element undergoes simultaneously both oxidation and reduction are called as disproportional reactions.

5. Calculate the empirical and molecular formula of the compound containing 80% Carbon, 20% Hydrogen. If the molecular mass of the compound is 30 then determine the molecular formula. [Qy.-19]

Ans. For C \Rightarrow 80/12 = 6.6 for H \Rightarrow 20/1 = 20. Now divide 6.6 and 20 by 6.6 to get simple whole no. ratio of C and H which will come 1:3 So emperical formula is CH₃ and its mass is 15

Now to calculate n we have 30/15 = 2So Molecular formula is $CH_3 \times 2 = C_2H_6$

6. Define auto-oxidation (disproportination) reaction and its examples. [Qy. - '23 & '24]

Ans. Metal displacement reactions:

- Place a zinc metal strip in an aqueous copper sulphate solution taken in a beaker. Observe the solution, the intensity of blue colour of the solution slowly reduced and finally disappeared.
- The zinc metal strip became coated with brownish metallic copper. This is due to the following metal displacement reaction.

PH: 8124201000 / 8124301000

7. An organic compound present in vinegar has 40 % carbon, 6.6 % hydrogen and 53.4 % oxygen. Find the empirical formula of the compound.


[Hy. - '23; Mar-'24]

Ans.

Element	Percentage	Atomic mass	Relative No. of moles	Simplest ratio mole	Simplest ratio (in whole no)
C	40	12	$\frac{40}{12} = 3.3$	$\frac{3.3}{3.3} = 1$	1
Н	6.6	1	$\frac{6.6}{1} = 6.6$	$\frac{6.6}{3.3} = 2$	2
О	53.4	16	$\frac{53.4}{16} = 3.3$	$\frac{3.3}{3.3} = 1$	1


The Empirical formula is CH₂O.

- 8. Explain the following redox reactions with an example. [Qy.-'24]
 - i) Decomposition reaction.
 - ii) Metal displacement reaction.
- **Ans.** (i) Decomposition reaction: Redox reactions in which a compound breaks down into two or more components are called decomposition reactions. These reactions are opposite to combination reactions. In these reactions, the oxidation number of the different elements in the same substance is changed.



(ii) Metal displacement reactions: Place a zinc metal strip in an aqueous copper sulphate solution taken in a beaker. Observe the solution, the intensity of blue colour of the solution slowly reduced and finally disappeared.

The zinc metal strip became coated with brownish metallic copper. This is due to the following metal displacement reaction.



ADDITIONAL QUESTIONS

Additional Choose the Correct Answers 1 MARK

- 1. Consider the following statements
 - 1 Matter possesses mass.
 - 2 22 carat gold is a mixture.
 - 3 Dry ice is a compound.

Which of the following statement(s) given above is/ are correct?

- (a) 1 & 3
- (b) only 1
- (c) 1 & 2
- (d) 1, 2 & 3

[Ans. (d) 1, 2 & 3]

- 2. The solid state of matter is converted into gas by
 - (a) sublimation
- (b) deposition
- (c) freezing
- (d) condensation

[Ans. (a) sublimation]

- 3. The characteristic feature of orderly arrangement of molecules belongs to
 - (a) Solids
- (b) Liquid
- (c) Gases
- (d) None of these

[Ans. (a) Solids]

- 4. The oxidation number of oxygen in O₂ is
 - (a) 0
- (b) +1
- (c) +2
- (d) -2

[Ans. (a) 0]

- **5.** The volume occupied by any gas at S.T.P. is ____
 - (a) 22.4 litres
- (b) 2.24 litres
- (c) 224 litres
- (d) 0.224 litres

[Ans. (a) 22.4 litres]

- 6. What will be the basicity of H₃BO₃, which is not a protic acid?
 - (a) one
- (b) two
- (c) three
- (d) four

[Ans. (a) one]

- 7. Which form of based on physical characteristics possess neither definite volume nor definite shape?
 - (a) Solids
- (b) Liquids
- (c) Gases
- (d) Both (a) and (b)

[Ans. (c) Gases]

- 8. Identify the incorrect statement about a compound.
 - (a) A molecule cannot be separated into its constituent elements by physical methods of separation
 - (b) A molecule of a compound has atoms of different elements
 - (c) A compound retains the physical properties of its constituent element
 - (d) The ratio of atoms of different elements in a compound is fixed

[Ans. (c) A compound retains the physical properties of its constituent element]

- 9. Which among the following statement(s) describe an element?
 - i) It is pure substance which could be split into two or more simpler substances.
 - ii) It is a pure substance which cannot be split into simpler substance
 - iii) It's composition is not uniform
 - iv) All the above
 - (a) only (iv)
- (b) only (ii)
- (c) (ii) and (iii)
- (d) (i) and (iii)

[Ans. (b) only (ii)]

- **10.** Atomicity of nitrogen is
 - (a) 1
- (b) 2
- (c) 3
- (d) zero

[Ans. (b) 2]

- 11. Atoms are too small with diameter of 10^{-10} m and weigh approximately
 - (a) 10^{-27} kg
- (b) 10^{-27} g
- (c) 10^{-31} kg
- (d) 10^{-31} g

[Ans. (a) 10^{-27} kg]

- **12.** 1 amu (or) 1u ≈
 - (a) $1.6605 \times 10^{-25} \text{ kg}$
- (b) $1.6605 \times 10^{-26} \text{ kg}$
- (c) $1.6605 \times 10^{-27} \text{ kg}$
- (d) $1.6605 \times 10^{-28} \text{ kg}$

[Ans. (c) 1.6605×10^{-27} kg]

- 13. 12 g of carbon-12 contains _____ carbon atoms.
 - (a) 6.022×10^{23}
- (b) 6

(c) 12

(d) 12.022×10^{23}

[Ans. (a) 6.022×10^{23}]

- 14. Atomicity of nitrogen is
 - (a) 1
- (b) 2
- (c) 3
- (d) zero

[Ans. (b) 2]

Hint: Atomicity is defined as total number of atoms present in the molecule

15. Statement I: Equivalent mass of Mg is determined by Oxide Method.

Statement II: Molecular mass is calculated using vapour density.

- (a) Both the statements are individually true
- (b) Both the statements are individually true and statement II is the correct explanation of statement I.
- (c) Statement I is true but statement II is false.
- (d) Statement I is false but statement II is true.

[Ans. (a) Both the statements are individually true]

16. Match list I with list II and identify the correct code.

List I			List II
A	Bronze	1	Element
В	Table salt	2	Homogeneous mixture
C	Gold	3	Alloy
D	Petrol	4	Compound

- A B C D
- (a) 1 4 2 3
- **(b)** 3 4 1 2
- (c) 2 3 4 1
- (d) 4 2 3

[Ans. (b) 3 4 1 2]

- 17. $Fe^{2+} \longrightarrow Fe^{3+} + e^-$ is a(n) reaction.
 - (a) redox
- (b) reduction
- (c) oxidation
- (d) decomposition

[Ans. (c) oxidation]

18. Assertion: An element that has a fractional atomic mass.

Reason : An element exist as isotope.

- (a) Both assertion and reason are correct and reason is the correct explanation for assertion.
- (b) Both assertion and reason are correct but reason is not the correct explanation for assertion
- (c) Assertion is true but reason is false.
- (d) Both assertion and reason are false. [Ans. (a) Both assertion and reason are correct and reason is the correct explanation for assertion.]
- 19. The oxidation number of hydrogen in LiH is ____.
 - (a) + 1
- (b) -1
- (c) +2
- (d) -2

[Ans. (b) -1]

- **20.** The empirical formula and molecular mass of a compound are CH₂O and 180 g respectively. What will be the molecular formula of the compound?
 - (a) $C_9H_{19}O$
- (b) CH₂O
- (c) $C_6H_{12}O_6$
- (d) $C_2H_4O_2$

[Ans. (c) C₆H₁₂O₆]

Hint: Molecular formula is equal to empirical formula multiplied by molecular weight

- 21. In the reaction $NH_3 + H_2O \longrightarrow NH_4^+ + O\overline{H}$, NH_3 is acidic in nature. The reason for its acidity is
 - (a) Acceptance of one H⁺ from water
 - (b) Release of one OH⁻ ion
 - (c) Due to the nitrogen atom
 - (d) All the above.

[Ans. (a) Acceptance of one H⁺ from water]

- **22.** If a beaker holds 576 g of water, what will be the gram molecules of water in that beaker?
 - (a) 23 gram molecule
- (b) 23%
- (c) 32%
- (d) 32 gram molecule

[Ans. (d) 32 gram molecule]

Sol: Molecular mass of $H_2O = 2 \times 1 + 16$

 $= 18 \text{ g mol}^{-1}$

18 g of water = 1 gram molecule

∴ 576 g of water = $\frac{1 \times 576}{18}$

= 32 gram molecules.

- 23. Unit of Avogadro's number is
 - (a) mol
- (b) g
- (c) mol^{-1}
- (d) no unit

[Ans. (c) mol^{-1}]

enquiry@surabooks.com

24. Match the following prefixes with their multiples.

Equ	Equivalent Mass (E)		lecular Mass (M)
A	E _{KMnO₄} (Acidic)	1	M/2
В	E _{KMnO₄} (Neutral)	2	M
C	$E_{H_3PO_2}$	3	M/3
D	$E_{H_3PO_3}$	4	M/5

- (a)
- **(b)**
- (c)

[Ans. (a) 4 3 2 1]

25. One 'U' stands for the mass of

- (a) An atom of carbon-12
- (b) $1/12^{th}$ of the carbon-12
- (c) 1/12th of hydrogen atom
- (d) One atom of any of the element

[Ans. (b) 1/12th of the carbon-12]

26. Assertion: The atomic masses of most of the elements are in fractions.

Reason : The atomic mass represents the ratio of the average mass of the atom to one avogram.

- (a) Both assertion and reason are correct and reason is the correct explanation for assertion.
- (b) Both assertion and reason are correct but reason is not the correct explanation for assertion
- (c) Assertion is true but reason is false.
- (d) Both assertion and reason are false. [Ans. (b) Both assertion and reason are correct but reason is not the correct explanation for assertion]

27. Identify disproportionation reaction.

- (a) $CH_4 + 2O_2 \longrightarrow CO_2 + 2H_2O$ (b) $CH_4 + 4Cl_2 \longrightarrow CCl_4 + 4HCl$
- (c) $2F_2 + 2OH^- \longrightarrow 2F^- + OF_2 + H_2O$
- (d) $2NO_2 + 2OH^- \longrightarrow NO_2 + NO_3 + H_2O$

[Ans. (d) $2NO_2 + 2OH^- \longrightarrow NO_2 + NO_3 + H_2O$]

28. The oxidation number of Cr in $Cr_2O_7^{2-}$ is _____. (a) +6 (b) -6 (c) +7 (d) -7

[Ans. (a) + 6]

29. The oxidation state of a element in its uncombined state is

- (a) zero
- (b)+1
- (c) -1
- (d) none

[Ans. (a) zero]

30. Assertion: The number of oxygen atoms in 16 g of oxygen and 16 g of ozone is same.

: Each of the species represent 1 g atom Reason of oxygen.

- (a) Both assertion and reason are correct and reason is the correct explanation for assertion.
- (b) Both assertion and reason are correct but reason is not the correct explanation for assertion
- (c) Assertion is true but reason is false.
- (d) Both assertion and reason are false.

[Ans. (a) Both assertion and reason are correct and reason is the correct explanation for assertion.]

31. Assertion: The ash is produced by burning paper in air is lighter than the original mass of paper.

: The residue is left after the combustion Reason of a chemical reaction that entities is always lighter.

- (a) Both assertion and reason are correct and reason is the correct explanation for assertion.
- (b) Both assertion and reason are correct but reason is not the correct explanation for assertion
- (c) Assertion is true but reason is false.
- (d) Both assertion and reason are false.

[Ans. (c) Assertion is true but reason is false.]

- 32. Assertion: Oxalic acid is a dibasic acid : It contains two basic radicals
 - (a) Both assertion and reason are correct and reason is the correct explanation for assertion.
 - (b) Both assertion and reason are correct but reason is not the correct explanation for assertion
 - (c) Assertion is true but reason is false.
 - (d) Both assertion and reason are false.

[Ans. (c) Assertion is true but reason is false.]

- **33.** How many moles of magnesium phosphate, Mg₃(PO₄), will contain 0.25 moles of oxygen atoms?
 - (a) 0.02
- (b) 3.125×10^{-2}
- (c) 1.25×10^{-2}
- (d) 2.5×10^{-2}

[Ans. (b) 3.125×10^{-2}]

Sol: 8 mol of O = 1 mol of
$$Mg_3(PO_4)_2$$

0.25 mol O = $\frac{1 \times 0.25}{8}$ mol of $Mg_3(PO_4)_2$
= 3.125×10^{-2} mol of $Mg_3(PO_4)_2$.

34. Assertion: Equal volumes of all the gases do not ! 39. Rusting of iron is an example of reaction. contain equal number of atoms.

Reason : Atom is the smallest particle which takes part in chemical reactions.

- (a) Both assertion and reason are correct and reason is the correct explanation for assertion.
- (b) Both assertion and reason are correct but reason is not the correct explanation for assertion
- (c) Assertion is true but reason is false.
- (d) Both assertion and reason are false.

[Ans. (b) Both assertion and reason are correct but reason is not the correct explanation for assertion]

35. Assertion: Fluorine has an oxidation state of 1 in all its compounds.

Reason: Fluorine is the most electronegative element of the periodic table.

- (a) Both assertion and reason are correct and reason is the correct explanation for assertion.
- (b) Both assertion and reason are correct but reason is not the correct explanation for assertion
- (c) Assertion is true but reason is false.
- (d) Both assertion and reason are false.

[Ans. (a) Both assertion and reason are correct and reason is the correct explanation for assertion.]

- **36.** Among the three metals, zinc, copper and silver, the electron releasing tendency decreases in the following order.
 - (a) zinc >silver>copper (b) zinc >copper >silver
 - (c) silver>copper>zinc (d) copper>silver>zinc

[Ans. (b) zinc >copper >silver]

- **37.** Consider the following statements :
 - (i) Oxidation number of He = zero
 - (ii) Increase in oxidation number results in reduction.
 - (iii) The substance undergoing increase in oxidation number is reducing agent.

Which among the above statement(s) is/are correct?

- (a) only (i)
- (b) (ii) and (iii)
- (c) (i) and (iii)
- (d) only (ii)

[Ans. (c) (i) and (iii)]

- **38.** What is the ratio of empirical formula mass to molecular formula mass of benzene?
 - (a) 1:6
- (b) 6:1
- (c) 2:3
- (d) 3:2

[Ans. (a) 1:6]

- - (a) Combustion
 - (b) decomposition
 - (c) reduction reaction and redox reaction
 - (d) hydrolysis

[Ans. (c) reduction reaction and redox reaction]

- 40. Maximum oxidation state is present in the central metal atom of which compound
 - (a) CrO₂Cl₂
- (c) $[Fe(CN)_{6}]^{3-}$
- (d) MnO

[Ans. (a) CrO,Cl,]

- 41. The compound in which mass percentage of carbon is 75% and that of hydrogen is 25% is

 - (a) C_2H_6 (b) C_2H_2 (c) CH_4

[Ans. (c) CH₄]

42. Which of the following statement(s) is/are not true about the following decomposition reaction.

$$2KClO_3 \longrightarrow 2KCl + 3O_2$$

- (i) Potassium is undergoing oxidation
- (ii) Chlorine is undergoing oxidation
- (iii) Oxygen is reduced
- (iv) None of the species are under going oxidation and reduction.
- (a) only (iv)
- (b) (i) and (iv)
- (c) (iv) and (iii)
- (d) All of these

[Ans. (b) (i) and (iv)]

43. Match the list-I with list-II and select the correct answer using the code given below the list.

	List-I		List-II
A	$Cr_2O_7^{2-}$	1	+5
В	MnO_4^{2-}	2	+6
C	VO ₃ ²⁻	3	+3
D	FeF ₆ ³⁻	4	+7

 \mathbf{C} D B A

- 3 (a)
- 4 3 2 1 **(b)**
- 2 4 3 1 (c)
- 3 (d)

[Ans. (c) 2 4 1 3]

44. Identify the correct statements with reference to the given reaction

$$P_4 + 3OH^- + 3H_2O \longrightarrow PH_3 + 3H_2PO_2^-$$

- (i) Phosphorous is undergoing reduction only
- (ii) Phosphorous is undergoing oxidation only

PH: 8124201000 / 8124301000 enquiry@surabooks.com

- (iii) Phosphorous is undergoing both oxidation and reduction.
- (iv) Hydrogen is undergoing neither oxidation nor reduction.
- (a) only (iii)
- (b) both (iii) and (iv)
- (c) only (i)
- (d) None of these

[Ans. (b) both (iii) and (iv)]

45. Identify the correct statement(s) with respect to the following reaction:

 $Zn + 2HCl \longrightarrow ZnCl_2 + H_2$

- (i) Zinc is acting as an oxidant
- (ii) Chlorine is acting as a reductant
- (iii) Hydrogen is not acting as an oxidant
- (iv) Zn is acting as a reductant
- (a) only (ii)
- (b) only (iv)
- (c) both (ii) and (iii)
- (d) both (ii) and (i)

[Ans. (b) only (iv)]

46. Match the items in column list-I with relevant items in list-II.

	List-I		List-II
A	Ions having positive charge	1	anion
В	Ions having negative charge	2	-1
С	Oxidation number of fluorine in NaF	3	0
D	The sum of oxidation number of all atoms in a neutral molecule	4	cation

- A D
- 3 (a)
- 3 **(b)**
- 2 4 1 (c)
- (d) 2 3

[Ans. (d) 4 1 2 3]

- 47. Give an example of molecule in which the ratio of the molecular formula is six times the empirical formula.
 - (a) $C_6H_{12}O_6$
- (b) CH₂O
- (c) CH₄

(d) Na_2CO_3 [Ans. (a) $C_6H_{12}O_6$]

- 48. Which formula of a compound is a whole number multiple of the empirical formula?
 - (a) matter
- (b) mass
- (c) energy
- (d) weight

[Ans. (a) matter]

49. Assertion (A): In the reaction between potassium permanganate and potassium iodide, permanganate ions act as oxidising agent.

- Reason (R) : Oxidation state of manganese changes from +2 to +7 during the reaction.
- (a) Both A and R are true and R explains A
- (b) Both A and R are true but R does not explain A
- (c) A is true but R is false
- (d) Both A and R are false

[Ans. (c) A is true but R is false]

50. The change in the oxidation number of S in H₂S and SO, in the following industrial reaction:

$$2H_2S_{(g)} + SO_{2(g)} \longrightarrow 3S_{(s)} + H_2O_{(g)}$$

- (a) -2 to 0, +4 to 0 (b) -2 to 0, +4 to -1
- (c) -2 to -1, +4 to 0
- (d) -2 to -1, +4 to -2

[Ans. (a) -2 to 0, +4 to 0]

Sol: Reduction (+4 to 0)

$$2H_2S + SO_2 \longrightarrow 3S + H_2O$$

Oxidation (-2 to 0)

51. In which of the following reactions, hydrogen peroxide acts as an oxidising agent?

(a)
$$I_2 + H_2O_2 + 2OH^- \longrightarrow 2I^- + 2H_2O + O_2$$

- (b) $PbS + 4H_2O_2 \longrightarrow PbSO_4 + 4H_2O$
- (c) $2MnO_4^- + 3H_2O_2 \longrightarrow 2MnO_2 + 3O_2 + 2H_2O$
- (d) $HOCl + H_2O_2 \longrightarrow H_2O^+ + Cl^- + O_2$ [Ans. (b) PbS + $4H_2O_2 \longrightarrow PbSO_4 + 4H_2O$]
- **52.** Two elements X and Y (atomic mass of X = 75; Y = 16) combine to give a compound having 76% of X. The formula of the compound is?
 - (a) XY
- (b) X_2Y (c) X_2Y_2 (d) X_2Y_3

[Ans. (d) X_2Y_3]


53. Assertion (A): Among halogens fluorine is the best oxidant.

Reason (R) : Fluorine is the most electronegative atom.

- (a) Both A and R are true and R explains A
- (b) Both A and R are true but R does not explain A
- (c) A is true but R is false
- (d) Both A and R are false

[Ans. (a) Both A and R are true and R explains A]

54. Identify the redox reaction taking place in a ! 58. Limiting reagent in a chemical reaction is the beaker.



Initial stage

Final stage

(a)
$$Zn_{(s)} + Cu^{2+}_{(aq)} \longrightarrow Zn^{2+}_{(aq)} + Cu_{(s)}$$

(b)
$$Cu_{(s)}^{(s)} + 2Ag_{(aq)}^{+} \longrightarrow Cu_{(aq)}^{2+} + 2Ag_{(s)}^{(s)}$$

(c)
$$Cu_{(s)} + Zn^{2+}_{(aq)} \longrightarrow Zn_{(s)} + Cu^{2+}_{(aq)}$$

(c)
$$Cu_{(s)} + Zn^{2+}_{(aq)} \longrightarrow Zn_{(s)} + Cu^{2+}_{(aq)}$$

(d) $2Ag_{(s)} + Cu^{2+}_{(aq)} \longrightarrow 2Ag^{+}_{(aq)} + Cu_{(s)}$

[Ans. (d)
$$2Ag_{(s)} + Cu^{2+}_{(aq)} \longrightarrow 2Ag^{+}_{(aq)} + Cu_{(s)}$$
]

Reason: Since Cu is more reactive than Ag, it displaces Ag+ ions from its salt solution. Which get deposited on the copper rod

55. Match the list I with List II and select the correct answer using the code given below the list.

	List-I		List-II
A	n	1	6.02×10^{23} Ne atoms
В	Vapour density	2	0.01 moles of solute in one L of solution
C	22.4 L at S.T.P	3	Molecular mass/2
D	Centimolar solution	4	Molecular mass/ empirical formula mass

- (a)
- **(b)**
- (c)
- (d)

[Ans. (b) 4 3 1 2]

- **56.** A compound has an empirical formula C₂H₄O. If the value of n = 2 the molecular formula of the compound is
 - (a) C_2H_4O
- (b) CH₂O
- (c) CH₂
- (d) $C_4H_8O_2$

[Ans. (d) C₄H_oO₂]

- **57.** If ten volumes of dihydrogen gas reacts with five volumes of dioxygen gas then, how many volumes of water vapour would be produced?
 - (a) 1
- (b) 2
- (c) 5
- (d) 10 [Ans. (d) 10]

Hint: $2H_{2(g)} + O_{2(g)} \longrightarrow 2H_2O_{(g)}$

- reactant in which
 - (a) left some amount unreacted after the completion of reaction
 - (b) reacts completely in the reaction
 - (c) does not react in the reaction
 - (d) all of these

[Ans. (b) reacts completely in the reaction]

59. Assertion: When 4 moles of H₂ reacts with 2 moles of O2, then 4 moles of water

Reason: O, will act as limiting reagent.

- (a) Both assertion and reason are true and reason is the correct explanation of assertion.
- (b) Both assertion and reason are true but reason is not the correct explanation of assertion.
- (c) Only assertion is true but reason is false.
- (d) Both assertion and reason are false.

[Ans. (b) Both assertion and reason are true but reason is not the correct explanation of assertion.]

60. Match the list-I with list-II and select the correct answer using the code given below the list.

List-I			List-II			
A	Molecular formula	1	Completely consumed			
B Stoichiometric Equation		2	Left unreacted			
<u> </u>		2	F			
C	Limiting reagent	3	$n \times \text{Empirical formula}$			
D	Excess reagent	4	Balanced equation			

- (a)
- 4
- (d)

[Ans. (b) 3 4 1 2]

61. Assertion: K_{20} . Al_2O_3 . SiO_2 . $6H_2O$ is the empirical formula of potash alum.

: It is a double salt.

- (a) Both assertion and reason are correct and reason is the correct explanation for assertion.
- (b) Both assertion and reason are correct but reason is not the correct explanation for assertion

PH: 8124201000 / 8124301000

- (c) Assertion is true but reason is false.
- (d) Both assertion and reason are false.

[Ans. (b) Both assertion and reason are correct but reason is not the correct explanation for assertion

62. Anything that has mass and occupies space is called

- (a) matter
- (b) weight
- (c) energy
- (d) system

[Ans. (a) matter]

63. The mass of one mole of a substance is

- (a) molecular mass
- (b) Atomic mass
- (c) molar mass
- (d) Nuclear mass

[Ans. (c) molar mass]

64. Equal volume of nitrogen and Hydrogen gases will react to form ammonia in favourable condition then the limiting reagent is

- (a) H₂
- (b) N₂
- (c) NH₃
- (d) No reactant is a limiting regent

[Ans. (b) N₂]

65. Which of the following is correct?

- (a) Elemental analysis of a compound gives the mass percentage of atoms present in the compound
- (b) Using the mass percentage, we can determine the empirical formula of the compound
- (c) Molecular formula of the compound can be arrived at from the empirical formula using the molar mass of the compound.
- (d) All the above are correct

[Ans. (d) All the above are correct]

66. All oxidation reactions are accompanied by reactions.

- (a) accession
- (b) addition
- (c) reduction
- (d) decomposition

[Ans. (c) reduction]

67. During which reactions do the oxidation numbers of elements changes?

- (a) metabolic reactions
- (b) reduction reactions
- (c) exchange reactions
- (d) redox reactions

[Ans. (d) redox reactions]

68. When an ion in a compound is replaced by an ion of another element then these are called ______ reactions.

- (a) displacement
- (b) ionic
- (c) chemical
- (d) physical

[Ans. (a) displacement]

ADDITIONAL SHORT ANSWERS

1. Mixture of salt and water is a solution while that of oil and water is not. Explain.

Ans. Solution is a homogeneous mixture of two or more components. Salt in water is homogeneous and therefore it is a solution. Whereas oil in water is heterogeneous or immiscible mixture and so is not a solution.

2. Why is air sometimes considered as a heterogeneous mixture?

Ans. Air sometimes considered as a heterogeneous mixture due to the presence of dust particles which form a separate phase.

3. List the differences between elements and compounds.

Ans.

	Elements	Compounds	
(i)	An element consists of	Compounds are made	
	only one type of atom.	up of molecules which	
		contain two or more	
		atoms of different	
		elements.	
(ii)	Element can exist	Properties of	
	as monatomic or	compounds are different	
	polyatomic units.	from those of their	
		constituent elements.	
(iii)	Eg: Monatomic unit -	Eg:	
	Gold (Au),	Carbon dioxide (CO ₂),	
	Copper (Cu);	Glucose (C ₆ H ₁₂ O ₆)	
	Polyatomic unit -	0 12 0	
	Hydrogen (H ₂)		

4. Write a note on 'mixture' based on the chemical classification of matter.

Ans. Two or more substances mix together in any ratio without any chemical intraction is called mixture.

(i) Homogeneous mixture:

A mixture having uniform composition throughout is called homogeneous mixture. **Eg**: salt solution, air etc.,

(ii) Heterogeneous mixture:

A mixture in which the composition is not uniform throughout and different components can be observed is called heterogeneous mixture. **Eg**: Mixture of salt and sugar, cereals and pulse etc.,

5. Explain the classification of matter based on chemical composition.

Ans. Chemical Classification: Pure substances are composed of simple atoms or molecules. They are further classified as elements and compounds.

(a) Element:

- □ An element consists of only one type of atom.
- Element can exist as monatomic or polyatomic units.
- □ **Eg : Monatomic unit** Gold (Au), Copper (Cu); **Polyatomic unit** Hydrogen (H₂)

enquiry@surabooks.com

(b) Compound:

- Compounds are made up of molecules which contain two or more atoms of different elements.
- **Eg:** Carbon dioxide (CO_2) , Glucose $(C_6H_{12}O_6)$.
- 6. How will you classify matter based on physical state?
- **Ans.** Physical Classification of Matter: Matter can be classified as solids, liquids and gases based on their physical state. The physical state of matter can be converted into one another by modifying the temperature and pressure suitably.
- 7. Which law co-relates the mass and volume of a gas?
- **Ans.** Avogadro's law. It states that equal volume of all gases under the same conditions of temperature and pressure contain equal number of molecules.
- 8. Bring about the dissimilarities in mole concept and molar mass by clearly analysing them.

Ans.

Mole		Molar Mass	
1.	It is defined as the amount of the substance that contains as many specified elementary particles as the number of atoms in 12g of C ¹² .	It is defined as the mass of one mole of the substance.	
2.	1 mole = 6.022×10^{23} particles	$Molar mass = \frac{Mass}{mol} g mol^{-1}$	

- 9. (i) If an acid is mono basic, how will you relate their equivalent mass and molecular mass.
 - (ii) What is the basicity of $H_4 P_2 O_7$? [Qy. '23]
 - (iii) Give any two examples for dibasic acids.
- **Ans.** (i) If an acid is mono basic, then its equivalent mass = Molecular mass.
 - (ii) Basicity of H₄P₂O₇ is 4 (Tetrabasic acid)
 - (iii) Examples of dibasic acid are H₂SO₄, H₃PO₃.
- 10. Define Avogadro number.
- **Ans.** The total number of entities present in one mole of any substance is equal to 6.022×10^{23} . This number is called Avogadro number.
- 11. Define molar volume.
- Ans. The volume occupied by one mole of any substance in the gaseous state at a given temperature and pressure is called molar volume. One mole of an Ideal gas is equal to 22.4 L (Or) 22400ml at STP conditions.

- **12.** Does one gram mole of a gas occupy 22.4 L under all conditions of temperature and pressure.
- **Ans.** No, one gram mole of a gas occupies 22.4 L only under STP conditions, i.e. at 273 K temperature and 760mm of pressure. $(1.0315 \times 10^{15} \text{ Pa})$
- **13.** Why are the atomic mass of most of the elements fractional?
- **Ans.** It is because most of the elements occur in nature as a **mixture of isotopes** and their atomic masses are the average relative atomic masses of the isotopes depending in their abundance.
- 14. Write down the formulae for calculating the equivalent mass of an acid, base and oxidising agent.
- Ans. (i) Equivalent Mass of Acids:

 $E = \frac{\text{Molar mass of the acid}}{\text{Basicity of the acid}}$

(ii) Equivalent Mass of Bases :

 $E = \frac{\text{Molar mass of the base}}{\text{Acidity of the base}}$

(iii) Equivalent Mass of Oxidising agent :

 $E = \frac{\text{Molar mass of the oxidising agent}}{\text{no. of moles of electrons gained}}$ by one mole of the oxidising agent

- **15.** What do you understand by stoichiometric coefficients in a chemical equation?
- **Ans.** The co-efficients of reactants and products involved in a chemical equation represented by the balanced form are known as stoichiometric co-efficients.

Eg: $N_{2(g)} + 3H_{2(g)} \longrightarrow 2NH_{3(g)}$

The stoichiometric co-efficients are 1, 3 and 2 respectively.

- 16. Write the simplest formula for the following.
 - (i) N_2O_4 (ii) $C_6H_{12}O_6$ (iii) H_2O (iv) H_2O_7
- Ans. (i) NO_2
- (ii) CH₂O
- (iii) H₂O
- (iv) HO.
- 17. Categorise the redox reactions that occur in our daily life.

Ans. • Fading of the colour of the clothes

- □ Burning of cooking gas, fuel, wood, etc.
- Rusting of Iron
- Extraction of Metals.

enquiry@surabooks.com

- 18. $2Cu_2S + 3O_2 \longrightarrow 2Cu_2O + 2SO_2$
 - (i) In this reaction which substance is getting oxidised and which substance is getting reduced?
 - (ii) Name the oxidising and reducing agents.
- **Ans.** (i) Oxygen is being added to Cu, (ie.,) Cu_2S is oxidised to Cu_2O and the other reactant O_2 is getting reduced.
 - (ii) Cu₂S is the a reducing agent.O₂ is an oxidising agent.
- 19. How would you know whether a redox reaction is taking place in an acidic, alkaline or neutral medium.
- **Ans.** □ If H⁺ any acid appears on either side of the chemical equation, the reaction occurs in acidic solution.
 - If OH[−] or any base appears on either side of the chemical equation, the reaction occurs in basic solution.
 - □ If neither H⁺, OH[−] nor any acid or base is present in the chemical equation, the solution is neutral.
- **20.** Zn rod is immersed in CuSO₄ solution. What will you observe after an hour? Explain you observation in terms of redox reaction.
- **Ans.** The blue colour of CuSO₄ solution will get discharged and reddish brown copper metal will be deposited on Zn rod.
 - □ This is because blue colour Cu²⁺ (in CuSO₄) gets reduced to Cu by accepting two electrons from Zn, which gets oxidised to colourless ZnSO₄.

Reduction
$$Zn_{(s)} + CuSO_{4(aq)} \longrightarrow ZnSO_{4(aq)} + Cu_{(s)}^{0}$$
Oxidation

21. What is molar Volume?

- **Ans.** Molar volume is the volume occupied by one mole of any substance in the gaseous state at STP. It is equal to 2.24×10^{-2} m³ (22.4 L)
- **22.** Define stoichiometry.
- **Ans.** Stoichiometry is the quantitative relationship between reactants and products in a balanced chemical equation in moles.

- **23.** What will be oxidation number of sulphur in $S_2O_8^{2-}$ ion and $S_4O_6^{2-}$ ion? [HOTS]
- **Ans. (i)** In $S_2O_8^{2-}$, there is one peroxide bond (-O-O-) therefore, two oxygen atoms having oxidation number -1 (i.e., O_2^{2-}) and for the other six oxygen atoms, the oxidation number is -2. $S_2O_8^{2-} = 2x + (-2 \times 6) + (-1 \times 2) = -2$

$$S_2O_8^{2-} = 2x + (-2 \times 6) + (-1 \times 2) = -2$$

 $2x = +12 \Rightarrow x = +6$

- (ii) In $S_4O_6^{2-}$, two S-atoms have $O = S_5^{+5} = S_6 = S_6^{-5} = S_6$
- **24.** Nitric acid is an oxidising agent and reacts with PbO but it does not react with PbO₂. Explain why?
- Ans. (i) Nitric acid in an oxidising agent. It oxidises an element from lower oxidation state to higher oxidation state. In PbO, lead is in lower oxidation state of +2. HNO₃ oxidises lead from Pb²⁺ to Pb⁴⁺ PbO + 2 HNO₃ \rightarrow Pb (NO₃)₂ + H₂ O
 - (ii) In PbO₂, lead is in +4 oxidation state and cannot be oxidised further. Therefore no reaction takes place.
- **25.** Which one of the two, ClO_2^- or ClO_4^- shows disproportionation reaction and why? [HOTS]
- **Sol:** The oxidation state of Cl in ClO₂⁻ is +3. So, chlorine can get oxidised as well as reduced and can act as reductant and oxidant.

The disproportionation reaction of ClO_2^- is

$$3ClO_2^- \longrightarrow Cl^- + ClO_3^-$$

In ClO_4^- , Cl is in its highest oxidation state, So it can only be an oxidant.

- **26.** Identify the type of redox reaction taking place in the following.
 - (i) $3Mg_{(s)}^{0} + N_{2(g)}^{0} \longrightarrow Mg_{3} N_{2(s)}^{-3}$
 - (ii) $\overset{_{+5}}{V_2} \overset{_{-2}}{O_{5(s)}} + 5\overset{_{0}}{Ca_{(s)}} \longrightarrow \overset{_{0}}{2} \overset{_{0}}{V_{(s)}} + \overset{_{+2}}{5}\overset{_{-2}}{CaO_{(s)}}$
 - (iii) $2\overset{+1}{K}\overset{+5}{Cl}\overset{-2}{O_{3(s)}}$ \longrightarrow $2\overset{+1}{K}\overset{-1}{Cl}_{(s)} + 3\overset{0}{O_{2(g)}}$
 - (iv) $\overset{0}{\text{Ca}}_{(s)} + \overset{+1}{2}\overset{-2}{\text{H}}_{2}\overset{-2}{\text{O}}_{(l)} \longrightarrow \overset{+2}{\text{Ca}}\overset{-2+1}{\text{OH}}_{2(aq)} + \overset{0}{\text{H}}_{2(g)}$

$$\begin{array}{ll} (v) & Br_{2(1)} + 2I_{(aq)} \longrightarrow 2Br_{(aq)}^{-} + I_{2(s)} \\ \\ (vi) & Cl_{2(g)} + 2OH_{(aq)}^{-} \longrightarrow ClO_{(aq)}^{-} + Cl_{(aq)}^{-} + H_{2}O_{(l)} \end{array}$$

Ans. (i) Combination reaction

- (ii) Displacement reaction
- (iii) Decomposition reaction
- (iv) Metal displacement reaction
- (v) Non-metal displacement reaction
- (vi) Disproportionation reaction.

27. How can we say sugar has solid and water has liquid?

Ans. When a sugar dissolves into tea or coffee, the liquid transforms the sugar into a liquid. So it can fit in with the liquid and slide in with the molecules. If you try to evaporate the water for long enough, you will turn the sugar back into a solid.

28. The approximate production of Na_2CO_3 per month is 424×10^6 g while that of methyl alcohol is 320×10^6 g. Which is produced more in terms of moles?

Ans. Na₂CO₃ mass =
$$424 \times 10^6$$
 g
Molecular mass of Na₂CO₃ = $(23 \times 2) + 12 + (16 + 3)$
= $46 + 12 + 18$
= 106 g

No. of moles of Na₂CO₃ =
$$\frac{\text{Mass of Na}_2\text{CO}_3}{\text{molecular mass of Na}_2\text{CO}_3}$$
$$= \frac{424 \times 10^6 \text{g}}{106\text{g}}$$
$$= 4 \times 10^6 \text{ moles}$$

Methyl alcohol mass = 320×10^6 g

Ans.

29. How many moles of glucose are present in 720g of glucose?

Ans. Glucose =
$$C_6H_{12}O_6$$

Molecular mass = $(12 \times 6) + (1 \times 12) + (16 \times 6)$
of glucose = $72 + 12 + 96 = 180$

Number of mole =
$$\frac{\text{Mass of glucose}}{\text{Molecular mass of glucose}}$$

of glucose = $\frac{720}{180}$ = 4 moles.

30. What do you understand by the terms acidity and Basicity?

Ans. Acidity: The number of hydroxyl ions present in one mole of a base is known as the acidity of the base.

Basicity: The number of replaceable hydrogen atoms present in a molecule of the acid is referred to as its basicity.

31. What is meant by plasma state? Give an example.

Ans. Gaseous state of matter at very high temperature containing gaseous ions and free electron is referred to as the plasma state eg., Lightning.

32. What is displacement reactions?

Ans. Redox reactions in which an ion (or an atom) in a compound is replaced by an ion (or atom) of another element are called displacement reactions. They are further classified into (i) metal displacement reactions (ii) non-metal displacement reactions.

33. What is decomposition reaction?

Ans. Redox reactions in which a compound breaks down into two or more components are called decomposition reactions. These reactions are opposite to combination reactions. In these reactions, the oxidation number of the different elements in the same substance is changed.

34. Define Average atomic mass?

Ans. Average atomic mass is defined as the average of the atomic masses of all atoms in their naturally occurring isotopes.

Examples: Chlorine consist of chlorine isotope $_{17}\text{Cl}^{35}$ and $_{17}\text{Cl}^{37}$ are in the ratio 77:23, the average relative atomic mass of chlorine is $(35 \times 77 + 37 \times 23) / 100 = 35.46 \text{ u.}$

PH: 8124201000 / 8124301000

35. Draw a flow chart to illustrate classification of matter.

Physical Classification

Solid Liquid Gas Mixtures

Eg: Ice Eg: Water Eg: Water Vapour homogeneous heterogeneous

Eg: Green tea Eg: Oil and Water

Eg: Copper Eg: Copper Sulphate

36. Discuss the characteristic the properties of physical classification of matter.

[LOTS]

Ans.

S.No	Properties	Solid	Liquid	Gas
1.	Volume	definite	definite	indefinite
2.	Shape	definite	indefinite	indefinite
3.	Compressibility	cannot be compressed	can be compressed	can be highly compressed
4.	Arrangement of molecules	regular and close to each other	random or irregular but almost close to each other.	random and wide apart
5.	Bonding	strong intermolecular bonds	relatively strong intermolecular bonds; slightly weaker than solid	very weak intermolecular bonds.
6.	Fluidity	cannot flow	can flow from higher to lower level	can flow in all directions
7.	Example	Ice	Water	Water vapour

Additional Long Answers

5 MARK

1. What is the condition for molar Volume?

Ans. The molar volume of any ideal gas at 273 degree kelvin and 1 atm pressure is equal to 22.4 L (or) 22400ml.

Ideal gas equation is PV is equal to nRT where P is pressure at 1 atm. and temperature 273 degree kelvin is called Standard Temperature and Pressure.

R is the gas constant and is equal to 0.082 dtm^3 . atm. $k^{-1}.\text{mol}^{-1}$.

Hence V is equal to nRT/P.

V is equal to 22.4L.

2. Define auto-oxidation (disproportination) reaction and its examples.

Ans. Displacement reaction: Redox reactions in which an ion (or an atom) in a compound is replaced by an ion (or atom) of another element are called displacement reactions. They are further classified into







(i) metal displacement reactions (ii) non-metal displacement reactions.

(i) Metal displacement reactions:

Place a zinc metal strip in an aqueous copper sulphate solution taken in a beaker. Observe the solution, the intensity of blue colour of the solution slowly reduced and finally disappeared. The zinc metal strip became coated with brownish metallic copper. This is due to the following metal displacement reaction.



(ii) Non-metal displacement:



3. Write any three rules assigning for the oxidation number?

Ans. The oxidation state of a free elements (i.e. in its uncombined state) is zero.

Example: each atom in H₂, Cl₂, Na, S₈ have the oxidation number of zero.

■ For a monatomic ion, the oxidation state is equal to the net charge on the ion.

Example : The oxidation number of sodium in Na⁺ is +1.

The oxidation number of chlorine in Cl^- is -1.

■ The algebric sum of oxidation states of all atoms in a molecule is equal to zero, while in ions, it is equal to the net charge on the ion.

- 4. Distinguish between the following.
 - (i) Atomic and molecular mass
- (ii) Atomic mass and atomic weight
- (iii) Empirical and molecular formula
- (iv) Moles and molecules.

Ans.

(i)	Atomic Mass	Molecular Mass		
	Atomic mass is the mass of a single atom, which is its	Molecular weight is the mass of one molecule.		
	collective mass of neutron, proton and electrons. Molecular mass can be calculated from the sum			
		masses of all atoms present in a compound.		
(ii)	Atomic Mass	Atomic Weight		
	Atomic mass is the mass of a single atom, which is its	Atomic weight is the average weight of an elements with		
	collective mass of neutron, proton and electrons.	respect to all its isotopes and their relative abundance.		
(iii)	Empirical Formula	Molecular Formula		
	It represents the simplest whole number ratio of various	The molecular formula shows the exact number of different		
	atoms present in one molecule of the compound.	types of atoms present in a molecules of a compound.		
	Empirical formula of Benzene is CH	Molecular formula of Benzene is C ₆ H ₆		
(iv)	Moles	Molecules		
	The amount of the substance that contains specified	Two or more atoms joint together by chemical bonds.		
	particles as the number of atoms in 12g of carbon - 12			
1	isotope			

NUMERICAL PROBLEMS

- 1. Calculate the number of atoms in each of the following.
 - (i) 52 g of He and (ii) 52 moles of He.
- **Sol:** (i) 1 mol of He \equiv 4g \equiv 6.022 \times 10²³ He atoms (ie) 4g of He contains 6.022 \times 10²³ He atoms

∴ 52g of He contains =
$$\frac{6.022 \times 10^{23} \times 52}{4}$$

= 7.83×10^{24}

52g of He contains 7.83×10^{24} He atoms.

(ii) 1 mol of He contains 6.022×10^{23} He atoms

∴ 52 moles of He contains =
$$\frac{6.022 \times 10^{23} \times 52}{1}$$

= 3.131×10²⁵

- 52 moles of He contains 3.132×10^{25} He atoms.
- 2. Calculate the mass of the following:
 - (i) 1 atom of silver (ii) 1 molecule of benzene
 - (iii) 1 molecule of water.
- **Sol:** (i) Molecular mass of silver (Ag) = 107.87 uMolar mass of $Ag = 107.87 \text{ g mol}^{-1}$

$$\therefore \text{ Mass of 1 atom of Ag} = \frac{\text{Molar mass}}{\text{Avogadro's number}}$$

$$= \frac{107.87 \text{ g mol}^{-1}}{6.022 \times 10^{23} \text{ mol}^{-1}}$$
$$= 17.91 \times 10^{-23} \text{ g}.$$

Mass of 1 atom of Ag = 17.91×10^{-23} g.

(ii) Molecular mass of benzene $(C_6H_6) = (6 \times 12.01 \text{ u}) + (6 \times 1\text{u}) = 78.06 \text{ u}$

Molar mass of benzene

Molar mass of benzene = 78.06 g mol^{-1}

Then, mass of 1 molecule of benzene

= Avogadro's number
=
$$\frac{78.06 \text{ g mol}^{-1}}{6.022 \times 10^{23} \text{ mol}^{-1}} = 12.96 \times 10^{-23} \text{ g}$$

Mass of 1 molecule of benzene = 12.94×10^{-23} g.

(iii) Molecular mass of water = $(2 \times 1u) + (1 \times 16u)$ = 18 u

Molar mass of water = 18 g mol^{-1}

Mass of 1 molecule of water

$$= \frac{\text{Molar mass of water}}{\text{Avogadro's number}}$$

$$= \frac{18g \text{ mol}^{-1}}{6.022 \times 10^{23} \text{ mol}^{-1}} = 2.99 \times 10^{-23} \text{ g}$$

Mass of 1 molecule of water = 2.99×10^{-23} g.

PH: 8124201000 / 8124301000

- 3. One million silver atoms weigh 1.79×10^{-16} g. Calculate the atomic mass of silver.
- **Sol:** No. of silver atoms = 1 million = 1×10^6 Mass of one million Ag atoms = 1.79×10^{-16} g Mass of 6.022×10^{23} atoms of silver

$$= \frac{1.79 \times 10^{-16} \,\mathrm{g}}{1 \times 10^{6}} \times 6.022 \times 10^{23} = 107.8 \,\mathrm{g}.$$

Atomic mass of silver = 6.022×10^{23} atoms of Ag \therefore The atomic mass of Ag = 107.8 g.

- 4. How much mass (in gram units) is represented by the following?
 - (i) 0.2 mol of NH_3 (ii) 3.0 mol of CO_2
 - (iii) 5.14 mol of H₅IO₆
- **Sol:** (i) Molar mass of NH₃ = $(1 \times 14 + 3 \times 1) = 17 \text{g mol}^{-1}$ Mass of 0.2 mol of NH₃ = 0.2 mol × 17g mol⁻¹ = 3.4 g
 - (ii) Molar mass of $CO_2 = (1 \times 12 + 2 \times 16)$ = 44 g mol⁻¹ Mass of 3 moles of $CO_2 = 3 \text{ mol} \times 44 \text{ g mol}^{-1}$ = 132 g
 - (iii) Molar mass of $H_5IO_6 = (5 \times 1 + 1 \times 127 + 6 \times 16)$ = **228 g mol**⁻¹ Mass of 5.14 mol of $H_5IO_6 = 5.14$ mol $\times 228$ g mol⁻¹ = **1171.9 g.**
- 5. What mass of N_2 will be required to produce 34g of NH_3 by the reaction, $N_2 + 3H_2 \longrightarrow 2NH_3$.
- **Sol:** The reaction is

$$\begin{array}{cccc} N_2 & + & 3H_2 & \longrightarrow & 2NH_3 \\ 1 \text{ mol} & 3 \text{ mol} & 2 \text{ mol} \\ 2 \times 14 & & 2(1 \times 14 + 3 \times 1) \\ 28g & & = 34g \end{array}$$

Thus, to produce 34.0 g ammonia, 28g of N_2 is required.

- **6.** Calculate the Formula Weights of the following compounds.
 - (a) NO_2 (b) Glucose $(C_6H_{12}O_6)$ (c) NaOH (d) $Mg(OH)_2$
- Sol: (a) NO_2 $1 \times AW \text{ of } N = 1 \times 14 = 14 \text{ amu}$ $2 \times AW \text{ of } O = 2 \times 16 = 32 \text{ amu}$ Formula weight of $NO_2 = 46 \text{ amu}$
 - (b) $C_6H_{12}O_6$ Glucose $6 \times AW \text{ of } C = 6 \times 12.01 = 72.06 \text{ amu}$ $12 \times AW \text{ of } H = 12 \times 1.008 = 12.096 \text{ amu}$

 $6 \times AW$ of $O = 6 \times 16 = 96.0$ amu Formula weight of Glucose is = 180.156 amu Formula weight of Glucose = 180 amu

- (c) NaOH

 1 × AW of Na = 1× 22.99 = 22.99 amu

 1 × AW of O = 1×16 = 16.00 amu

 1 × AW of H = 1×1.008 = 1.008 amu

 Formula weight of NaOH is = 39.998 amu

 Formula weight of NaOH is = 40 amu.
- (d) $Mg(OH)_2$ $1 \times AW \text{ of } Mg = 1 \times 24.305 = 24.305 \text{ amu}$ $2 \times AW \text{ of } O = 2 \times 16 = 32.000 \text{ amu}$ $2 \times AW \text{ of } H = 2 \times 1.008 = 2.016 \text{ amu}$ Formula weight of $Mg(OH)_2$ is = 58.321 amu Formula weight of $Mg(OH)_2$ is = 58 amu.
- 7. Calculate the equivalent weight of H₃PO₄ and Ca(OH)₂ on the basis of given reaction.

 H₃PO₄ + NaOH \longrightarrow NaH₂PO₄ + H₂O
 Ca(OH)₂ + HCl \longrightarrow Ca(OH)Cl + H₂O
- Sol: Equivalent weight of H_3PO_4 $= \frac{\text{Molecular mass}}{\text{No. of replaceable H}^+} = \frac{98}{1} = 98$ Equivalent weight of $Ca(OH)_2$ $= \frac{\text{Molecular mass}}{\text{No. of replacement OH}^-} = \frac{74}{1} = 74$
- 3. (i) Calculate the gram molecular mass of sugar having molecular formula $C_{12}H_{22}O_{11}$.
 - (ii) Calculate (a) The mass of 0.5g molecule of sugar and (b) Gram molecule of sugar in 547.2 g.

Sol: (i) Molecular mass of Sugar $(C_{12}H_{22}O_{11})$ = $12 \times 12 + 22 \times 1 + 11 \times 16 = 342g$

- (ii) (a) 1 gram molecule of sugar = 342 g $\therefore 0.5 \text{ g}$ molecule of sugar = 342×0.5 = 171 g
 - **(b)** 342 g of sugar = 1 gram molecule

$$547.2 \text{ of sugar} = \frac{1}{342} \times 547.2 = 1.6 \text{ gram molecule.}$$

- 9. Calculate the number of moles in the following.(i) 7.85 g of copper (ii) 4.66 mg of silicon
 - (iii) 65.6 mg of oxygen.
- **Sol:** (i) Moles of copper = $\frac{\text{Mass of copper}}{\text{atomic mass}}$ = $\frac{7.85}{63.546}$ = 0.123 mol.

- (ii) Moles of silicon = $\frac{\text{Mass of silicon}}{\text{atomic mass}}$ = $\frac{4.66 \times 10^{-3}}{28.1}$ = $1.658 \times 10^{-4} \text{ mol}$ (iii) Moles of oxygen = $\frac{\text{Mass of oxygen}}{\text{atomic mass}}$ 65.6×10^{-6}
- $= \frac{65.6 \times 10^{-6}}{16}$ $= 4.1 \times 10^{-6} \text{ mol.}$
- **10.** What will be the molecular formula for the compound, whose empirical formula is CH₂Cl and molar mass is 98.96 g?
- Sol: Empirical formula = CH_2Cl ; Empirical formula mass = $12.01 + 2 \times 1.008 + 35.453$ = 49.48 g $n = \frac{\text{molecular mass}}{\text{empirical formula mass}}$ $= \frac{98.96g}{49.48g} = 2$ Molecular formula = $n \times \text{Empirical formula}$
- $= 2 \times CH_2Cl = C_2H_4Cl_2$ 11. Calculate the oxidation number of nitrogen in
- **Sol:** (i) Nitrous acid: HNO₂ + 1 + x 2 × 2 = 0
 - (ii) Nitric acid: $HNO_3 + 1 + x 2 \times 3 = 0$ x = +5

nitrous acid and nitric acid

- **12.** Balance the following reaction by oxidation number method.
- **Sol:** $MnO_4^{-1} + H_2S + H^+ \longrightarrow Mn^{2+} + S$ (Acidic Medium)
 - (i) Write oxidation number of elements $\begin{array}{cccc} MnO_4^{-1} & + & H_2S & \longrightarrow & Mn^{2+} & + & S \\ (+7)(-2) & (+1)(-2) & & +2 & 0 \end{array}$
 - (ii) Balance the number of atoms of the elements in which oxidation number changes

$$MnO_4^{-1} + H_2S \longrightarrow Mn^{2+} + S$$

(+7) (-2) (+2) 0

(iii) Decide the oxidation and reduction reaction on the basis of difference of oxidation number. Increase in oxidation number by 2(Oxidation)

$$MnO_4^{-1} + H_2S + H^+ \longrightarrow Mn^{2+} + S$$

Decrease in oxidation number by 5(Reduction)

(iv) On multiplying oxidation reaction by 5 and reduction reaction by 2 to balance the change in oxidation number.

$$2MnO_4^{-1} + 5H_2S \longrightarrow 2Mn^{2+} + 5S$$

(v) Balance the electric charge and atoms which do not change in oxidation number (spectators).

$$2MnO_4^{-1} + 5H_2S + 6H^+ \longrightarrow 2Mn^{2+} + 5S + 8H_2O$$

$$2(-1) 5(0) + 6(+1) = 2(+2) + 5(0) + 8(0)$$

$$-2 + 6 = +4$$

$$+ 4 = +4$$

In the above reaction the reactants and products are balanced in terms of electric charge and mass equivalence.

13. A compound on analysis was found to contain C = 34.6%; H = 3.85% and O = 61.55%. Calculate its empirical formula.

Sol:

Element	%	Percentage mass At. mass	Molar Ratio	Simplest Whole Number Ratio
С	34.6	$\frac{34.6}{12} = 2.88$	$\frac{2.88}{2.88} = 1$	3
Н	3.85	$\frac{3.85}{1} = 3.85$	$\frac{3.85}{2.88} = 1.335$	4
О	61.55	$\frac{61.55}{16} = 3.85$	$\frac{3.85}{2.88} = 1.335$	4

The empirical formula of the compound = $C_3H_4O_4$.

- **14.** Calculate the mass of the oxygen atom in amu.
- Sol: Oxygen

Mass of Oxygen atom = 2.656×10^{-23} 1 a.m.u (or) 1 u is equal to 1.66075 multiple The mass of oxygen atom in amu = $\frac{2.656 \times 10^{-23}}{1.66075 \times 10^{-24}} \approx 15.992$ a.m.u

- 15. How many moles of barium suphate is precipitated when 1 mole of aluminium sulphate reacts completely with barium chloride?
- **Sol:** $Al_2 (SO_4)_3 + 3 BaCl_2 \rightarrow 3 Ba SO_4 + 2 AlCl_3$ When 1 mole of aluminium sulphate reacts with barium chloride, 3 moles of $BaSO_4$ is precipitated.

- 16. Calculate the molecular mass of the following:
 - a) KMnO₄
- b) Crystalline Oxalic acid
- c) Methane
- Sol: (a) KMnO₄

1 × atomic mass of K = 1 × 39 = 39
Mn = 1 × 55 = 55
O = 4 × 16 =
$$\frac{64}{158}$$

 \therefore Molecular mass of KMnO₄ = $1\overline{58}$

(b) Crystalline Oxalic acid

$$C \rightarrow 2 \times 12 = 24$$

$$O \rightarrow 4 \times 16 = 64$$

$$H \qquad 2 \times 1 = 2$$

$$4 \times 1 = 4$$

$$2 \times 16 = 32$$

$$126$$

:. Molecular mass of oxalic acid = 126

(c) Methane CH₄

$$C \rightarrow 1 \times 12 = 12$$

$$H \rightarrow 4 \times 1 = 4$$

$$16$$

- \therefore Molecular mass of $CH_4 = 16$
- 17. Calculate the number of atoms/molecules present in the following:
 - a) 10g of Hg
- b) 1.8g of water
- c) 100g of sulpurdioxide
- d) 1kg of acetic acid
- **Sol:** (a) 10g of Hg

Atomic mass of Hg = 200 g mol^{-1} 200 g of mercury contains 6.022×10^{23} atoms of mercury.

10 g of mercury contains =
$$\frac{10 \times 6.022 \times 10^{23}}{200}$$

= 0.301×10^{23}
= 3.01×10^{22}
atoms of mercury.

(b) 1.8g of water

1 mole of water = 18 g mol^{-1} 18 g of water contains 6.022×10^{23} molecules of water

1.8g of water contains
$$= \frac{1.8 \times 6.022 \times 10^{23}}{18}$$
$$= 0.6022 \times 10^{23}$$
$$= 6.022 \times 10^{22}$$
Molecules of water

(c) 100g of sulphur dioxide

Molecular mass of $SO_2 = 64$ 64g of sulphur dioxide contains = 6.022×10^{23} Molecules of SO_2

$$\therefore 100g \text{ of SO}_2 \text{ contains} = \frac{100 \times 6.022 \times 10^{23}}{64}$$

$$= 9.41 \times 10^{23}$$
molecules of SO₂

(d) 1Kg of acetic acid

Molecular mass of acetic acid = 60 60g of acetic acid contains = 6.022×10²³ Molecules of acetic acid

... 1000g of acetic acid contains

$$= \frac{1000 \times 6.022 \times 10^{23}}{60}$$
$$= 100.37 \times 10^{23}$$
molecules of acetic acid

- **18.** Calculate the number of moles present in the following:
 - a) 50 g of calcium chloride
 - b) 120 g of sodium hydroxide
 - c) 46 g of ethanol
 - d) 90 g of magnesium oxide
 - e) 19.5 g of potassium
- **Sol:** (a) 50 g of calcium chloride

Molar mass of calcium chloride = 111

No. of moles =
$$\frac{\text{Mass}}{\text{Molar mass}}$$

No. of moles = $\frac{50}{111}$ = 0.450 moles

(b) 120 g of sodium hydroxide

Molar mass of sodium hydroxide = 40

No. of moles =
$$\frac{\text{Mass}}{\text{Molar mass}}$$

No. of moles (n) = $\frac{120}{40}$ = 3 moles

(c) 46 g of ethanol

Molecular mass of ethanol = 46

No. of moles =
$$\frac{\text{Mass}}{\text{Molar mass}}$$

No. of moles (n) =
$$\frac{46}{46}$$
 = 1 mole

(d) 90 g of magnesium oxide

Molecular mass of MgO = 40

No. of moles =
$$\frac{\text{Mass}}{\text{Molar mass}}$$

No. of moles = $\frac{90}{40}$ = 2.25 moles

(e) 19.5 g of potassium

Atomic mass of pottassium = 39

No. of moles =
$$\frac{\text{Mass}}{\text{Molar mass}}$$

No. of moles = $\frac{19.5}{39}$ = 0.5 moles

19. Calculate the molar volume of the following:

- $88 g of CO_2$
- b) 5 moles of methane
- c) 460 g of formic acid
- d) 3.0115×10^{23} molecules of SO₂gas

Sol: (a) 88 g of CO₂

Molar mass of
$$CO_2$$
 = 44 g
Molar volume of 44 g (1mole) of CO_2
= 2.24× 10⁻² m³

The volume of 88g (2 moles) =
$$\frac{2.24 \times 10^{-2} \times 88}{44}$$

= 4.48×10^{-2} m³

(b) 5 moles of methane

Molar mass of methane = 16 gMolar volume of 16 g (1mole) of methane $= 2.24 \times 10^{-2} \text{ m}^3$

Volume of 5 moles (80g) of methane

$$= \frac{2.24 \times 10^{-2} \times 80}{16}$$
$$= 11.2 \times 10^{-2} \,\mathrm{m}^3$$

460 g of formic acid

Molar mass of formic acid = 46 gMolar volume of 46 g (1mole) of formic acid $= 2.24 \times 10^{-2} \text{ m}^3$

Molar volume of 460g of (10 moles) of formic

$$= \frac{2.24 \times 10^{-2} \times 460}{46}$$
$$= 22.4 \times 10^{-2} \text{ m}^3$$

(d) 3.0115×10^{23} molecules of SO_2 gas

$$6.023 \times 10^{23} \text{ molecules} = 1 \text{ mole}$$

$$3.0115 \times 10^{23} \text{ molecules} = \frac{1}{6.022 \times 10^{23}} \times 3.0115 \times 10^{23} \times 3.0115 \times 3.0115 \times 10^{23} \times 3.0115 \times 3.0115$$

Molar volume of 1 mole of $SO_2 = 2.24 \times 10^{-2} \text{m}^3$

Molar volume of 0.5 moles of SO₂

$$= 2.24 \times 10^{-2} \times 0.5$$
$$= 1.12 \times 10^{-2} \text{ m}^3$$

20. Calculate the equivalent mass of the following

- a) Zn b) Nitrate ion c) sodium
- Sol: (a) Zn

Equivalent mass =
$$\frac{\text{Atomic mass}}{\text{Valency}}$$

= $\frac{65}{2} = 32.5 \text{ g}$

(b) Nitrate ion (NO₃⁻)

Equivalent mass of an ion = $\frac{\text{Formula mass}}{\text{Change of ion}}$ Equivalent mass of $NO_3^- = \frac{62}{1} = 62 g$

(c) Sodium Equivalent mass = $\frac{\text{Atomic mass}}{\text{Valency}}$

Equivalent mass of sodium = $\frac{23}{1}$ = 23 g

21. 1.05 g of a metal gives on oxidation 1.5g of its oxide. Calculate its equivalent mass.

Sol: Mass of oxygen =
$$1.5 - 1.05$$

= 0.45 g

0.45g of oxygen combines with 1.05 g of metal.

 \therefore 8 g of oxygen combines with $\frac{8 \times 1.05}{0.45}$ g of metal = 18.66 g of metal

 \therefore equivalent mass of metal = 18.66g equ⁻¹

22. Calculate equivalent mass of the following

- Sodium hydroxide
- Aluminium hydroxide **b**)
- ammonium hydroxide
- Calcium hydroxide d)
- Magnesium hydroxide

Sol: (a) NaOH equivalent mass of NaOH = $\frac{40}{1}$ = 40

- (b) Aluminium hydroxide equivalent mass of Al(OH)₃ = $\frac{78}{2}$ = 26
- (c) Ammonium hydroxide equivalent mass of $NH_4OH = \frac{35}{1} = 35$
- (d) Calcium hydroxide equivalent mass of $Ca(OH)_2 = \frac{74}{2} = 37$
- (e) Magnesium hydroxide Mg(OH), equivalent mass of Magnesium hydroxide $=\frac{58}{2}=29$

23. Calculate Equivalent mass of the following

- Hydrochloric acid
- Nitric acid **b**)
- Acetic acid c)
- d) Crystalline oxalic acid
- Phosphorous acid

Sol: (a) Hydrochloric acid

Equivalent mass of an acid

$$= \frac{\text{Molar mass of the acid}}{\text{Basicity of the acid}}$$

Equivalent mass of HCl=
$$\frac{36.5}{1}$$

= 36.5

(b) Nitric acid Equivalent mass of $HNO_3 = \frac{Molar mass}{basicity}$

$$=\frac{63}{1}=63$$

 $= \frac{63}{1} = 63$ (c) Acetic acid (CH₃ COOH)
Equivalent max Equivalent mass of acetic acid = $\frac{\text{Molar mass}}{\text{hasicity}}$

$$=\frac{60}{1}=60$$

(d) Crystalline oxalic acid

Equivalent mass of oxalic acid = $\frac{\text{Molar mass}}{\text{basicity}}$

equivalent mass
$$=$$
 $\frac{126}{2} = 63$

(e) Phosphorous acid (H₃ PO₃)

Equivalent mass of phosphorous acid

$$= \frac{\text{Molar mass}}{\text{basicity}} = \frac{82}{2} = 41$$

PH: 8124201000 / 8124301000

 \therefore equivalent mass of H₂PO₂ = 41

24. 3.24 g of titanium reacts with oxygen to form 5.40 g of the metal oxide. Find the empirical formula of the metal oxide?

Sol: Weight of Titanium = 3.24 g; Weight of metal oxide = 5.40 g Weight of Oxygen = (5.40 - 3.24) = 2.16 g

Element	Percentage	Atomic mass	Relative No. of moles	Simple ratio mole	Simplest whole Number Ratio
Ti	3.24	48	$\frac{3.24}{48} = 0.0675$	$\frac{0.0675}{0.0675} = 1$	1
0	2.16	16	$\frac{2.16}{16} = 0.135$	$\frac{0.135}{0.0675} = 2$	2

.. The empirical formula is Ti O₂

25. A compound contains 11.99% N, 13.70% O, 9.25% B and 65.06% F. Find its empirical formula Sol:

Element	Percentage	Atomic mass	Relative No. of moles	Simple ratio mole	Simplest whole Number Ratio
N	11.99	14	$\frac{11.99}{14} = 0.856$	$\frac{0.856}{0.856} = 1$	1
О	13.70	16	$\frac{13.70}{16} = 0.856$	$\frac{0.856}{0.856} = 1$	1
В	9.25	10	$\frac{9.25}{10} = 0.925$	$\frac{0.925}{0.856} = 1$	1
F	65.06	19	$\frac{65.06}{19} = 3.424$	$\frac{3.424}{0.856} = 4$	4

[∴] Empirical formula of the compound in NOBF₄

26. A organic compound used for welding operation contains the following composition by mass: C = 92.3%, H = 7.7%. Find out the molecular formula of the compound. At STP, 10.0 L of this gas is found to weight 11.6g.

Sol: Determination of Molecular formula

Element	Percentage	Atomic mass	Relative No. of moles	Simple ratio mole	Simplest whole Number Ratio
С	92.3	12	$\frac{92.3}{12} = 7.7$	$\frac{7.7}{7.7} = 1$	1
Н	7.7	1	$\frac{7.7}{1} = 7.7$	$\frac{7.7}{7.7} = 1$	1

PH: 8124201000 / 8124301000

Empirical formula is CH

Molecular formula = $n \times emprical$ formula

Emperical formula mass $(1 \times 12) + (1 \times 1) = 12 + 1 = 13$

$$n = \frac{Molecular\ mass}{Empirical\ formula\ mass}$$

Molar mass =
$$\frac{\text{wt. of the substance} \times \text{Molar volume}}{\text{vol.of the substance}}$$
 at STF

Molar volume at STP = 2.24×10^{-2} m³ = 22.4 l = 22400 ml

Molar mass of the gas at STP =
$$\frac{11.6 \times 22.4}{10} = 25.9 = 26$$

$$n = \frac{26}{13} = 2$$

Molecular formula = n × (emp. formula) = $2 \times (CH) = C_2 H_2$

40

27. The organic compound Vitamin-C, has the following composition by mass: 40.92% C, 4.58% H, and the rest is oxygen. Determine its molecular formula. Molar mass of the substance is 176 g mol⁻¹.

Sol:

Element	Percentage	Atomic mass	Relative No. of moles	Simplest whole Number Ratio
С	40.92	12	$\frac{40.92}{12} = 3.41$	3
Н	4.58	1	$\frac{4.58}{1} = 4.58$	4
О	100 – [40.92 +4.58] =54.5	16	$\frac{54.5}{16} = 3.406$	3

Empirical formula is $C_3H_4O_3$

Empirical formula mass = $(12 \times 3) + (1 \times 4) + (3 \times 16) = 36 + 4 + 48 = 88$

Molecular formula = $n \times empirical$ formula

$$n = \frac{Molecular\ mass}{Empirical\ formula\ mass} = \frac{176}{88} = 2$$

n = 2

:. Molecular formula = $n \times (emp. formula) = 2 \times (C_3 H_4 O_3) = C_6 H_8 O_6$

REDOX REACTION ACTIVITY

A piece of cut apple becomes brown. Why? Can | Ans. you prevent it by a simple method?

Ans. Apple turns brown when cut since the surface is exposed to air and undergoes oxidation. It can be prevented by dipping sliced apples in lemon juice. Lemon juice is an antioxidant which takes in all the available oxygen and prevents it from reaching the apple's tissues.

Place an iron piece in a moist atmosphere and observe it after two days. Is there any deposition of new substance? Why does it happen? What is this phenomenon called?

Ans. When iron is exposed to moist air, the iron reacts with oxygen in the presence of moisture to from a reddish - brown chemical compound, iron - oxide. This phenomenon is called rusting. A new substance Iron (III) oxide is formed.

 $4 \text{ Fe} + 3\text{O}_2 + 6\text{H}_2\text{O} \rightarrow 4 \text{ Fe} (\text{OH})_3$

Calculate the oxidation number of underlined atoms of the following:

- 1. K_2MnO_4
- 2. $K_{2}CrO_{4}$
- 3. $\underline{NO_3}^-$
- 4. $H_4\underline{P}, O_7$
- $\underline{C}lO_3^-$
- 6. $\underline{\mathbf{A}}\mathbf{sO}_{3}^{3-}$

1. $K_{2}MnO_{4}$

Oxidation number of Mn be x

$$2(1) + x + 4(-2) = 0$$

$$2 + x - 8 = 0$$

$$x - 6 = 0$$

$$x = 6$$

Oxidation number of Mn in K_2 MnO₄ is +6.

2. $K_{2}CrO_{4}$

$$2(1) + x + 4(-2) = 0$$

$$2 + x - 8 = 0$$

$$x - 6 = 0$$

$$x = +6$$

Oxidation number of Cr in K₂ CrO₄ is +6.

3. \underline{NO}_3^-

$$x + 3(-2) = -1$$

 $x - 6 = -1$
 $x = -1 + 6 = +5$

Oxidation number of N in NO_3^- is +5.

4.
$$H_4 P_2 O_7$$

 $4(1) + 2x + 7(-2) = 0$
 $4 + 2x - 14 = 0$
 $2x - 10 = 0$
 $2x = 10$
 $x = 5$

Oxidation number of P in $H_4P_2O_7$ is +5.

5.
$$\underline{C}10_3^-$$

$$x+3(-2) = -1$$
$$x-6 = -1$$
$$x = +5$$

Oxidation number of Cl in ClO_3^- is +5.

6.
$$\underline{\mathbf{A}}\mathbf{sO}_3^{3-}$$

$$x+3(-2) = -3$$

$$x-6 = -3$$

$$x = -3+6$$

$$x = +3$$

Oxidation number of As in AsO_3^{3-} is +3.

- 4. An iron nail is placed in copper sulphate solution taken in the beaker. Observe it for some time? Find the changes that takes place and why?
- **Ans.** When iron nail is dipped in copper sulphate solution, the colour of copper sulphate tuns from blue to light green and reddish brown deposits is formed on iron nail. This is because iron is more reactive than copper, so it displaces Cu from CuSO₄ solution.

The displacement reaction can be written as

$$CuSO_4 + Fe \rightarrow FeSO_4 + Cu$$

5. The approximate production of Na_2CO_3 per month is 424×10^6 g while that of methyl alcohol is 320×10^6 g. Which is produced more in terms of moles?

Mass of Na₂CO₃ =
$$424 \times 10^6$$
g
No of moles (n) = Mass of the substance
Molar mass of the substance

$$= \frac{424 \times 10^6}{106}$$
$$= 4 \times 10^6 \,\text{moles}$$

Mass of
$$CH_3OH = 320 \times 10^6 g$$

No of moles =
$$\frac{\text{Mass of the substance}}{\text{Molar mass of the substance}}$$

= $\frac{320 \times 10^6}{32}$ = 10×10^6 moles

Methyl alcohol is produced more.

- 6. Find the molecular mass of FeSO₄7H₂O.
- **Ans.** Molecular mass of FeSO₄7H₂O

Atomic mass of Fe = 55.845

Atomic mass of S = 32.065

Atomic mass of $O_4 = 16 \times 4 = 64$

Atomic mass of $H_2 = 1.00794 \times 2 = 2.01588$

Atomic mass of O = 16

Molecular mass of

$$FeSO4.7H2O = 55.845 + 32.065 + 64 + 7$$

$$(2.01588 + 16)$$

$$= 278.02g/mol$$

7. The density of CO₂ is 1.977 kgm⁻³ at STP. Calculate the molecular mass of CO₂.

Ans. Density of $CO_2 = 1.977 \text{ Kgm}^{-3}$

$$\overrightarrow{PV} = nRT$$

No of moles = $\frac{Mass}{Molar Mass}$

$$PV = \frac{Mass}{Molar\ Mass} \times R \times T$$

$$Molar\ Mass = \frac{Mass}{V} \times \frac{R \times T}{P}$$

Density =
$$\frac{\text{Mass}}{\text{V}}$$

Molar Mass of
$$CO_2 = \frac{D \times R \times T}{P}$$

Standard Temperature = 273 K

Standard Pressure = 760mm of Hg = 1amu

$$= \frac{1.977 \times 0.0821 \times 273}{1}$$

= 44

8. How many moles of glucose are present in 720 g of glucose?

Ans. Mass of glucose = 720gMolecular weight of
glucose $(C_6H_{12}O_6)$ = 180No. of moles = $\frac{Mass}{Molar\ Mass}$ = $\frac{720}{180}$ = 4 moles

9. Calculate the weight of 0.2 mole of sodium carbonate.

Ans. No. of moles of $Na_2CO_3 = 0.2$ mole

Molar mass of $Na_2CO_3 = 106g/mol$ Mass = No of moles × molar mass of Na_2CO_3 = $0.2 \times 106 = 21.2g$

10. Calculate the equivalent mass of bicarbonate ion.

Ans. Bicarbonate ion = HCO_{3}^{-} Molar mass of HCO_{3}^{-} = 61 Equivalent mass of ion = $\frac{\text{Molar mass}}{\text{Charge of ion}}$ Equivalent mass of HCO_{3}^{-} = $\frac{61}{1}$ = 61

11. Calculate the equivalent mass of barium hydroxide

Ans. Equivalent mass of Ba(OH)₂

Molar mass of $Ba(OH)_2 = 171.34 \text{ g/mol}$

Acidity of the $Ba(OH)_2 = 2$

Equivalent mass of the Ba(OH)₂

$$= \frac{\text{Molar mass of the base}}{\text{Acidity of the base}}$$
$$= \frac{171}{2} = 85.5$$

12. Boric acid, H₃BO₃ is a mild antiseptic and is often used as an eye wash. A sample contains 0.543 mol H₃BO₃. What is the mass of boric acid in the sample.

Formula mass of boric acid H_3BO_3 = 61.834 amu
1 mole of H_3BO_3 = Molar mass of H_3BO_3 = 61.834 g

0.543 mole of H_3BO_3 = 61.834 × 0.543
= 33.57 g of H_3BO_3 The mass of

0.543 moles of H_3BO_3 = 33.57g

13. (i)
$$K_2Cr_2O_7 + KI + H_2SO_4 \longrightarrow K_2SO_4 + Cr_2(SO_4)_3 + I_2 + H_2O$$

Ans.

Decrease in O.N (Reduction)

Decreases by +3 $K_2Cr_2O_7 + KI + H_2SO_4 \rightarrow K_2SO_4 + Cr_2(SO_4)_3 + I_2 + H_2O$ +6

Increase in O N (oxidation)

Equalise the increase / decrease in O N by multiplying I species by 1

$$K_2Cr_2O_7 + 3KI + H_2SO_4 \longrightarrow K_2SO_4 + Cr_2(SO_4)_3 + 3I_2 + H_2O$$

Balance all other atoms except H and O

$$K_2Cr_2O_7 + 6KI + 7H_2SO_4 \longrightarrow 4K_2SO_4 + Cr_2(SO_4)_3 + 3I_2 + H_2O$$

Balance O atom by adding H₂O on the the side falling short of oxygen

$$K_2Cr_2O_7 + 6KI + 7H_2SO_4 \longrightarrow 4K_2SO_4 + Cr_2(SO_4)_3 + 3I_2 + H_2O + 6H_2O$$

So the balanced equation is

$$K_2Cr_2O_7 + 6KI + 7H_2SO_4 \longrightarrow 4K_2SO_4 + Cr_2(SO_4)_3 + 3I_2 + 7H_2O$$

(ii)
$$KMnO_4 + Na_2SO_3 \longrightarrow MnO_2 + Na_2SO_4 + KOH$$
 (Alkaline medium)

Decrease in O.N (Reduction)

Decreases by +3
$$KMnO_4 + Na_2SO_3 \rightarrow MnO_2 + Na_2SO_4 + KOH$$
+7
+4
Increases by +2
Increase in O N
(oxidation)

Equalise the increase / decrease in O N by multiplying Mn species by 2 and S species by 3

$$2KMnO_4 + 3Na_2SO_3 \longrightarrow 2MnO_2 + 3Na_2SO_4 + KOH$$

Balance all other atoms except H and O

Balance O atoms by adding H₂O molecules on the side falling short of oxygen atom.

$$2KMnO4 + 3Na2SO3 + H2O \longrightarrow 2MnO2 + 3Na2SO4 + 2KOH$$

(iii)
$$K_2Cr_2O_7 + KCI + H_2SO_4 \xrightarrow{} KHSO_4 + CrO_2CI_2 + H_2O$$

It is not a redox reaction.

(iv)
$$Cu + HNO_3 \longrightarrow Cu(NO_3)_2 + NO_2 + H_2O$$
Increase in O N
(Oxidation)

Increases by +2

 $Cu + HNO_3 \rightarrow Cu(NO_3)_2 + NO_2 + H_2O$
0

 $+5$
Decrease by +1
Decrease in O N
Reduction

Equalise the increase / decrease in O N by multiplying Cu species by +1 and N species by +2

$$Cu + 2HNO_3 \longrightarrow Cu(NO_3)_2 + 2NO_2 + H_2O$$

Balance all other atoms except H and O atoms

$$Cu + 2HNO_3 \longrightarrow Cu(NO_3)_2 + 2NO_2 + H_2O$$

Balance O atom by adding H₂O molecules on the side falling short of oxygen atoms.

$$Cu + 4HNO_3 \longrightarrow Cu(NO_3)_2 + 2NO_2 + H_2O$$

The balanced equation in

$$Cu + 4HNO_3 \longrightarrow Cu(NO_3)_2 + 2NO_2 + 2H_2O$$

(v)
$$P + HNO_3 \longrightarrow H_3PO_4 + NO_2 + H_2O$$

Increase in O N (Oxidation)

Decrease in O N
Reduction

Equalise the increase / decrease in O N by multiplying P species by +1 and N species by +5

(vi)
$$P + 5HNO_3 \longrightarrow H_3PO_4 + 5NO_2 + H_2O$$

All atoms are balanced
Balanced Equation is
 $P + 5HNO_3 \longrightarrow H_3PO_4 + 5NO_2 + H_2O$

(vii)
$$H_2C_2O_4 + KMnO_4 + H_2SO_4 \longrightarrow H_2SO_4 + MnSO_4 + CO_2 + H_2O$$

Increase in O N (Oxidation)

Increases by 1
$$H_2C_2O_4 + KMnO_4 + H_2SO_4 \rightarrow K_2SO_4 + MnSO_4 + CO_2 + H_2O_4 + K_2SO_4 + MnSO_4 + CO_2 + K_2SO_4 + MnSO_4 + K_2SO_4 + MnSO_5 + K_2SO_5 + MnSO_5 + K_2SO_5 + K_2SO_$$

Equalise the increase / decrease in O N by multiplying Cu species by 5 and Mn species by 1

$$5H_2C_2O_4 + KMnO_4 + H_2SO_4 \longrightarrow K_2SO_4 + MnSO_4 + 5CO_2 + H_2O_4$$

Balance all other atoms except H and O atoms

$$\begin{aligned} 5\mathrm{H}_2\mathrm{C}_2\mathrm{O}_4 + 2\mathrm{KMnO}_4 + 3\mathrm{H}_2\mathrm{SO}_4 & \longrightarrow \mathrm{K}_2\mathrm{SO}_4 \\ &+ 2\mathrm{MnSO}_4 + 10\mathrm{CO}_2 + \mathrm{H}_2\mathrm{O} \end{aligned}$$

Balance O atom by adding H₂O on the side falling short of oxygen atoms.

$$5\mathrm{H}_2\mathrm{C}_2\mathrm{O}_4 + 2\mathrm{KMnO}_4 + 3\mathrm{H}_2\mathrm{SO}_4 \longrightarrow \mathrm{K}_2\mathrm{SO}_4 \\ + 2\mathrm{MnSO}_4 + 10\mathrm{CO}_2 + \mathrm{H}_2\mathrm{O} + 7\mathrm{H}_2\mathrm{O}$$

The balanced equation in

$$\begin{aligned} 5\mathrm{H}_2\mathrm{C}_2\mathrm{O}_4 + 2\mathrm{KMnO}_4 + 3\mathrm{H}_2\mathrm{SO}_4 & \longrightarrow \mathrm{K}_2\mathrm{SO}_4 \\ &+ 2\mathrm{MnSO}_4 + 10\mathrm{CO}_2 + 8\mathrm{H}_2\mathrm{O} \end{aligned}$$

(viii)CuO + NH₃
$$\longrightarrow$$
 Cu + N₂ + H₂O

$$\mathrm{CuO} + \mathrm{NH_3} \longrightarrow \mathrm{Cu} + \mathrm{N_2} + \mathrm{H_2O}$$

$$Cu + 2e^- \longrightarrow Cu \text{ (Reduction)} \qquad \dots (1)$$

$$N^{-3} \longrightarrow N_2^0 + 3e^-$$
 (oxidation)(2)

Multiply Equation (2) by 2 to balance nitrogen atom

$$2N^{-3} \longrightarrow N_2^0 + 6e^- \qquad(3)$$

Multiply equation (1) by 3 to balance the number of electrons.

$$3Cu^{2+} + 6e^{-} \longrightarrow 3Cu$$
(4)

Add equation (3) and (4)

$$3Cu^{2+} + 2N^{-3} \longrightarrow 3Cu + N_2$$

Over all balanced equation

$$3CuO + 2NH_3 \longrightarrow 3Cu + N_2 + H_2O$$

Balance O atom by adding H_2O on the side falling short of it.

$$3\text{CuO} + 2\text{NH}_3 \longrightarrow 3\text{Cu} + \text{N}_2 + 3\text{H}_2\text{O}$$

PH: 8124201000 / 8124301000

(ix)
$$\operatorname{Zn} + \operatorname{HNO}_3 \longrightarrow \operatorname{Zn} (\operatorname{NO}_3)_2 + \operatorname{NH}_4 \operatorname{NO}_3 + \operatorname{H}_2 \operatorname{O}_3$$

$$Zn + HNO_3 \longrightarrow Zn (NO_3)_2 + NH_4NO_3 + H_2O$$

$$Zn^0 \longrightarrow Zn^{2+} + 2e^- \qquad \dots (1)$$

$$N^{+5} + 8e^{-} \longrightarrow N^{-3} \qquad \dots (2)$$

Multiply Equation (1) by 4 to balance the electrons

$$4Zn^0 \longrightarrow 4Zn^{2+} + 2e^- \qquad(3)$$

Add equation (3) and (2)

$$4Zn^0 \longrightarrow 4Zn^{2+} + 8e^{1}$$

$$N5+ + 8e^{1} \longrightarrow N^{3-}$$

$$4Zn^{0} + N^{5+} \longrightarrow 4Zn^{2+} + N^{3-}$$

Overall equation

$$4Zn + 10HNO_3 \longrightarrow 4Zn (NO_3)_2 + NH_4NO_3 + H_2O$$
Release all the stame expect O and H

Balance all the atoms except O and H

$$4Zn + 10HNO_{3} \longrightarrow 4Zn (NO_{3})_{2} + NH_{4}NO_{3} + H_{2}O$$

Balance oxygen atom by adding H₂O on the side falling short of oxygen atom.

$$4Zn + 10HNO_3 \xrightarrow{} 4Zn (NO_3)_2 + NH_4NO_3 + 3H_2O$$

14. (i) $MnO_{4}^{-} + Sn^{2+} \longrightarrow Mn^{2+} + Sn^{4+}$ **Ans.**

Decrease in O.N (Reduction)

Decreases by 5
$$MnO_{-4}^{-} + Sn^{2+} \rightarrow Mn^{2+} + Sn^{4+}$$

$$+7 + 2+ 2+ 2+ 1$$
Increase in O.N
(oxidation)

Equalise the increase / decrease in O.N by multiplying the oxidant and reductant by suitable numbers.

$$2MnO_{4}^{-} + 5Sn^{2+} \longrightarrow 2Mn^{2+} + 5Sn^{4+}$$

Balance all other atoms except O and H

$$2MnO^-_{4} + 5Sn^{2+} \longrightarrow 2Mn^{2+} + 5Sn^{4+}$$

Balance O atom by adding water on the side falling short of oxygen otoms.

$$2MnO_{4}^{-} + 5Sn^{2+} \longrightarrow 2Mn^{2+} + 5Sn^{4+} + 8H_{2}O$$

Balance H atom by adding H+ on the side falling short of hydrogen atoms.

$$2MnO_{4}^{-} + 5Sn^{2+} 16H^{+} \longrightarrow 2Mn^{2+} + 5Sn^{4+} + 8H_{2}O$$

(ii)
$$C_2O_4^{2-} + Cr_2O_7^{2-} \longrightarrow Cr^{3+} + CO_2$$

Increase in O.N (Oxidation)

Increases by +1
$$C_{2}O_{4}^{2-} + Cr_{2}O_{7}^{2-} \rightarrow Cr^{3+} + CO_{2}$$
+3
$$+6 \left[\begin{array}{c} Decreases by \\ +3 \end{array}\right]^{3+} +4$$
Decrease in O.N
(Reduction)

$$3C_2O_4^{2-} + Cr_2O_7^{2-} \longrightarrow 2Cr^{3+} + 6CO_2 + 7H_2O$$

$$3C_2O_4^{2-} + Cr_2O_7^{2-} + 14H^+ \longrightarrow 2Cr^{3+} + 6CO_2 + 7H_2O$$

(iii)
$$S_2O_3^{2-} + I_2 \longrightarrow S_2O_4^{2-} + I^-$$

Increase in O.N (Oxidation)

Increases by 1
$$S_2O_3^{2-} + I_2 \rightarrow S_2O_4^{2-} + I^-$$
+2 0 Decreases by 1 Decrease in O.N (Reduction)

$$\begin{split} &S_{2}O_{3}^{2-} + I_{2} & \longrightarrow S_{2}O_{4}^{2-} + I^{-} \\ &S_{2}O_{3}^{2-} + I_{2} & \longrightarrow S_{2}O_{4}^{2-} + 2I^{-} \\ &S_{2}O_{3}^{2-} + I_{2} + H_{2}O & \longrightarrow S_{2}O_{4}^{2-} + 2I^{-} \\ &S_{2}O_{3}^{2-} + I_{2} + H_{2}O & \longrightarrow S_{2}O_{4}^{2-} + 2I^{-} + 2I^{-} \\ &S_{2}O_{3}^{2-} + I_{2} + H_{2}O & \longrightarrow S_{2}O_{4}^{2-} + 2I^{-} + 2I^{-} \\ \end{split}$$

(iv)
$$Sb^{3+} + MnO_4^- \longrightarrow Sb^{5+} + Mn^{2+}$$

Decrease in O.N
Reduction

Decrease by 5

$$Sb^{3+} + MnO_4^- \rightarrow Sb^{5+} + Mn^{2+}$$
+3 Increase by 2
Increase in O.N
Oxidation

Equalise the increase / decrease in Oxidation number by multiplying with suitable numbers.

$$5Sb^{3+} + 2MnO_4^{-} \longrightarrow Sb^{5+} + Mn^{2+}$$

Balance all other atoms except O and H

$$5Sb^{3+} + 2MnO_4^- \longrightarrow 5Sb^{5+} + 2Mn^{2+}$$

Balance Oxygen atom by adding H₂O on the side falling short of oxygen.

$$5Sb^{3+} + 2MnO_4^- \longrightarrow 5Sb^{5+} + 2Mn^{2+} + 8H_2O$$

Balance hydrogen atom by adding H⁺ on the side falling short of hydrogen atoms.

$$5Sb^{3+} + 2MnO_4^{-} \longrightarrow 5Sb^{5+} + 2Mn^{2+} + 8H_2O + 16H^{+}$$

(v)
$$\operatorname{MnO_4}^{2-} \longrightarrow \operatorname{MnO_4}^{2-} + \operatorname{MnO_2}$$

 $\operatorname{MnO_4}^{2-} \longrightarrow \operatorname{MnO_4}^{-} + \operatorname{e^-} (\operatorname{Oxidation})$
 $+6 +7 + \cdots (1)$
 $\operatorname{MnO_4}^{2-} + 2\operatorname{e^-} \longrightarrow \operatorname{MnO_2} (\operatorname{Reduction})$
 $+6 +4 + \cdots (2)$

Multiply equation (1) by (2)

$$2MnO_4^{2-} \longrightarrow 2MnO_4^{-} + 2e^{-}$$
(3)

Add equation (2) and (3)

$$3MnO_4^{2-} \longrightarrow MnO_2 + 2MnO_4^{-}$$

Balance O atoms by adding H₂O on the side falling short of oxygen atoms

$$3MnO_4^{2-} \longrightarrow MnO_2 + 2MnO_4^{-} + 2H_2O$$

Balance H atoms by adding H⁺ on the side falling short of hydrogen atoms

$$3MnO_4^{2-} + 4H^+ \longrightarrow MnO_2 + 2MnO_4^- + 2H_2O$$

(vi)
$$MnO_4^- + Fe^{2+} \longrightarrow Mn^{2+} + Fe^{3+}$$

Decrease in O.N
Reduction

Decrease in +5

$$MnO_4^- + Fe^{2+} \rightarrow Mn^{2+} + Fe^{3+}$$
 $+7$
 $+2$
 $Increase in O.N$
Oxidation

Equalise the increase / decrease in Oxidation number by multiplying Mn species by 1 and Fe species by 5.

$$MnO_4^- + 5Fe^{2+} \longrightarrow Mn^{2+} + 5Fe^{3+}$$

Balance all other atoms except O and H

$$MnO_4^- + 5Fe^{2+} \longrightarrow Mn^{2+} + 5Fe^{3+}$$

Balance O atom by adding H₂O on the side falling short of hydrogen and equal number OH⁻ on the opposite side.

$$MnO_4^- + 5Fe^{2+} + 8H_2O \longrightarrow Mn^{2+} + 5Fe^{3+} + 4H_2O + 8OH^ MnO_4^- + 5Fe^{2+} + 8H_2O \longrightarrow Mn^{2+} + 5Fe^{3+} + 4H_2O + 8OH^-$$

(vii)
$$S_2O_3^{2-} + I_2 \longrightarrow S_2O_6^{2-} + I^-$$

Increase in O.N Oxidation

Increases by 3
$$+5$$

$$S_2O_3^{2-} + I_2 \omega S_2O_6^{2-} + I_{-1}$$

$$+2 \qquad 0 \qquad Decrease by 1$$
Decrease in O.N
Reduction

Equalise the increase / decrease in O.N by multiplying the S species by 1 and I species by 3.

$$S_2O_3^{2-} + 3I_2 \longrightarrow S_2O_6^{2-} + 3I^{-}$$

Balance all other atoms except O and H

$$S_2O_3^{2-} + 3I_2 \longrightarrow S_2O_6^{2-} + 6I^{-}$$

Balance O atom by adding H₂O on the side falling short of Oxygen.

$$S_2O_3^{2-} + 3I_2 + 3H_2O \longrightarrow S_2O_6^{2-} + 6I^{-}$$

PH: 8124201000 / 8124301000

Balance H atom by adding H+ ion on the side falling short of hydrogen.

$$S_2O_3^{2-} + 3I_2 + 3H_2O \longrightarrow S_2O_6^{2-} + 6I^- + 6H^+$$

Add equal number of OH- ion on the both side since the medium is alkaline

$$S_{2}O_{3}^{2-} + 3I_{2} + 3H_{2}O + 6OH^{-} \longrightarrow S_{2}O_{6}^{2-} + 6I^{-} + 6H^{+} + 6OH^{-}$$

$$S_{2}O_{3}^{2-} + 3I_{2} + 3H_{2}O + 6OH^{-} \longrightarrow S_{2}O_{6}^{2-} + 6I^{-} + 6H_{2}O$$

15. A compound contains 50% of X (atomic mass 10) and 50% Y (atomic mass 20). Give its molecular formula. Ans.

Element	Percentage	Atomic mass	Relative No. of moles	Simple Ratio Moles	Simplest whole number Ratio
X	50	10	$\frac{50}{10} = 5$	$\frac{5}{2.5} = 2$	2
Y	50	20	$\frac{50}{20} = 2.5$	$\frac{2.5}{2.5} = 1$	1

Its simplest formula = X_2Y

16. Determine the empirical formula of a compound containing K = 24.75%, Mn = 34.77% and rest is oxygen. *Ans.*

Element	Percentage	Atomic mass	Relative No. of moles	Simple Ratio Moles	Simplest whole number Ratio
K	24.75	39	$\frac{24.75}{39} = 0.63$	$\frac{0.63}{0.63} = 1$	1
Mn	34.77	55	$\frac{34.77}{55} = 0.63$	$\frac{0.63}{0.63} = 1$	1
О	$ \begin{array}{c} 100 - \\ (24.75 + 34.77) = 40.48 \end{array} $	16	$\frac{40.48}{16} = 2.53$	$\frac{2.53}{0.63} = 4$	4

The empirical formula is KMnO₄

