



Mathematics

12th Standard

VOLUME - I & II

Based on the Updated New Textbook

FREE
Practice Workbook
with
Practice Model
Question Papers

Salient Features

- Answers for **All Textual Questions** are given.
- Exhaustive Additional Questions & Answers in all chapters.
- Model Question Papers 1 to 6 (PTA) : Questions are incorporated in the appropriate sections.
- Govt. Model Question Paper-2019 [**Govt. MQP-2019**], Quarterly Exam - 2019 & 2023 [**Qy. - 2019 & 2023**], Half yearly Exam - 2019 & 2023 [**Hy. - 2019 & 2023**], Public Exam March - 2020; 2023 & 2024, May - 2022 [**Mar. -2020; 2023 & 2024, May - 2022**], Govt. Supplementary Exam September - 2020, August - 2021, July - 2022 & June - 2023 [**Sep -2020, Aug. - 2021, July - 2022 & June-2023**] and First & Second Revision Test - 2022 [**FRT & SRT - 2022**] questions are incorporated in the appropriate sections.
- Govt. Public Exam - March 2024 Question Paper is given with answers.



SURA PUBLICATIONS

Chennai

For Orders Contact



**80562 94222 / 81242 01000 / 81243 01000
96001 75757 / 78718 02000 / 98409 26027**

enquiry@surabooks.com Ph:8124201000 / 8124301000

All rights reserved © SURA Publications.

No part of this book may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, digitally, electronically, mechanically, photocopying, recorded or otherwise, without the written permission of the publishers. Strict action will be taken.

Code No : SG 322

Authors :

Mr. G.Selvaraj, M.Sc., M.Ed., M.Phil.,
Chennai

Edited by :

Mr. S. Sathish M.Sc., M.Phil.,

Reviewed by :

Mr.S. Niranjana, B.Tech., (NITT)PGDM (IIM)
Chennai

Our Guides for XI, XII Standard

- ❖ சுராவின் தமிழ் உரைநூல்
- ❖ Sura's Smart English
- ❖ Sura's Mathematics (EM/TM)
- ❖ Sura's Physics (EM/TM)
- ❖ Sura's Botany (EM/TM)
- ❖ Sura's Chemistry (EM/TM)
- ❖ Sura's Bio-Botany & Botany (EM/TM)
(Short Version & Long Version)
- ❖ Sura's Bio-Zoology & Zoology (EM/TM)
(Short Version & Long Version)
- ❖ Sura's Computer Science (EM/TM)
- ❖ Sura's Computer Applications (EM/TM)
- ❖ Sura's Commerce (EM/TM)
- ❖ Sura's Economics (EM/TM)
- ❖ Sura's Accountancy (EM/TM)
- ❖ Sura's Business Maths (EM)

PREFACE

Sir,

*An equation has no meaning, for me
unless it expresses a thought of GOD - Ramanujam*
[Statement to a friend]

Respected Principals, Correspondents, Head Masters / Head
Mistresses, Teachers,

From the bottom of our heart, we at SURA Publications
sincerely thank you for the support and patronage that you have
extended to us for more than a decade.

It is in our sincerest effort we take the pride of releasing **Sura's
Mathematics Guide Volume I and Volume II** for +2 Standard.
This guide has been authored and edited by qualified teachers
having teaching experience for over a decade in their respective
subject fields. This Guide has been reviewed by reputed Professors
who are currently serving as Head of the Department in esteemed
Universities and Colleges.

With due respect to Teachers, I would like to mention that
this guide will serve as a teaching companion to qualified teachers.
Also, this guide will be an excellent learning companion to students
with exhaustive exercises and in-text questions in addition to precise
answers for textual questions.

In complete cognizance of the dedicated role of Teachers, I
completely believe that our students will learn the subject effectively
with this guide and prove their excellence in Board Examinations.

I once again sincerely thank the Teachers, Parents and Students
for supporting and valuing our efforts.

God Bless all.

Subash Raj, B.E., M.S.
- Publisher
Sura Publications

All the Best

Head Office:

Sura Publications

1620, 'J' Block, 16th Main Road,
Anna Nagar, Chennai - 600 040.

Phones : 044 - 4862 9977, 044 - 4862 7755.

e-mail : orders@surabooks.com

website : www.surabooks.com

For Orders Contact



80562 94222
81242 01000
81243 01000
96001 75757
78718 02000
98409 26027

18/04/2024

CONTENT

Chapter No	Title	Page No
Volume - I		
1.	Applications of Matrices and Determinants	1-46
2.	Complex Numbers	47-80
3.	Theory of Equations	81-102
4.	Inverse Trigonometric Functions	103-130
5.	Two Dimensional Analytical Geometry-II	131-170
6.	Applications of Vector Algebra	171-220
Volume - II		
7.	Applications of Differential Calculus	221-274
8.	Differentials and Partial Derivatives	275-304
9.	Applications of Integration	305-336
10.	Ordinary Differential Equations	337-374
11.	Probability Distributions	375-404
12.	Discrete Mathematics	405-420
Public Exam March 2024 Question Paper with Answers		421-428

SYLLABUS

Chapter No	Title	Month	Chapter No	Title	Month
1.	Applications of Matrices and Determinants (1.1 - 1.5)	June	7.	Applications of Differential Calculus (7.1 - 7.10)	October
2.	Complex Numbers (2.1 - 2.8)	July	8.	Differentials and Partial Derivatives (8.1 - 8.3)	
3.	Theory of Equations (3.1 - 3.9)		8.	Differentials and Partial Derivatives (8.4 - 8.6)	Nov.
4.	Inverse Trigonometric Functions (4.1 - 4.5)		9.	Applications of Integration (9.1-9.5)	
I MID TERM TEST (June, July)			9.	Applications of Integration (9.6-9.9)	Dec.
4.	Inverse Trigonometric Functions (4.6 - 4.10)	August	10.	Ordinary Differential Equations (10.1 - 10.8)	
5.	Two Dimensional Analytical Geometry-II (5.1 - 5.7)		HALFYEARELY EXAMINATION (June to December)		
6.	Applications of Vector Algebra (6.1 - 6.5)		11.	Probability Distributions (11.1-11.6)	January
6.	Applications of Vector Algebra (6.6 - 6.10)	Sep.	12.	Discrete Mathematics (12.1 - 12.3)	
QUARTERLY EXAMINATION (June to September)			ANNUAL EXAMINATION (Full Portions)		

TO ORDER WITH US

SCHOOLS and TEACHERS:

We are grateful for your support and patronage to 'SURA PUBLICATIONS'
Kindly prepare your order in your School letterhead and send it to us.
For Orders contact: 81242 01000 / 81243 01000

DIRECT DEPOSIT

A/c Name : **Sura Publications**
Our A/c No. : **36550290536**
Bank Name : **STATE BANK OF INDIA**
Bank Branch : Padi
IFSC : SBIN0005083

A/c Name : **Sura Publications**
Our A/c No. : **21000210001240**
Bank Name : **UCO BANK**
Bank Branch : Anna Nagar West
IFSC : UCBA0002100

A/c Name : **Sura Publications**
Our A/c No. : **6502699356**
Bank Name : **INDIAN BANK**
Bank Branch : Asiad Colony
IFSC : IDIB000A098

A/c Name : **Sura Publications**
Our A/c No. : **1154135000017684**
Bank Name : **KVB BANK**
Bank Branch : Anna Nagar
IFSC : KVBL0001154

A/c Name : **Sura Publications**
Our A/c No. : **13240200032412**
Bank Name : **FEDERAL BANK**
Bank Branch : Anna Nagar
IFSC : FDRL0001324

A/c Name : **Sura Publications**
Our A/c No. : **50200031530945**
Bank Name : **HDFC BANK**
Bank Branch : Cenotaph Road, Teynampet
IFSC : HDFC0001216

After Deposit, please send challan and order to our address.
email to : orders@surabooks.com / Whatsapp : 81242 01000.



For Google Pay :
98409 26027



For PhonePe :
98409 26027



DEMAND DRAFT / CHEQUE

Please send Demand Draft / cheque in favour of 'SURA PUBLICATIONS'
payable at **Chennai**. The Demand Draft / cheque should be sent with your order
in School letterhead.

STUDENTS :

Order via Money Order (M/O) to



SURA PUBLICATIONS

1620, 'J' Block, 16th Main Road, Anna Nagar, Chennai - 600 040.

Phones : 044-4862 9977, 044-4862 7755.

Mobile : 96001 75757 / 81242 01000 / 81243 01000.

email : orders@surabooks.com Website : www.surabooks.com

Volume - I

MATHEMATICS

CONTENTS

1. Applications of Matrices and Determinants	3 - 46
2. Complex Numbers.....	47 - 80
3. Theory of Equations	81 - 102
4. Inverse Trigonometric Functions.....	103 - 130
5. Two Dimensional Analytical Geometry-II	131 - 170
6. Applications of Vector Algebra	171 - 220

CHAPTER

1

APPLICATIONS OF MATRICES AND DETERMINANTS

MUST KNOW DEFINITIONS

- ✦ If $|A| \neq 0$, then A is a non-singular matrix and if $|A| = 0$, then A is a singular matrix.
- ✦ The adjoint matrix of A is defined as the transpose of the matrix of co-factors of A.
- ✦ If $AB = BA = I_n$, then the matrix B is called the inverse of A.
- ✦ If a square matrix has an inverse, then it is unique.
- ✦ A^{-1} exists if and only if A is non-singular.
- ✦ Singular matrix has no inverse.
- ✦ If A is non – singular and $AB = AC$, then $B = C$ (left cancellation law).
- ✦ If A is non – singular and $BA = CA$ then $B = C$ (Right cancellation law).
- ✦ If A and B are any two non-singular square matrices of order n , then $\text{adj} (AB) = (\text{adj} B) (\text{adj} A)$
- ✦ A square matrix A is called orthogonal if $AA^T = A^T A = I$
- ✦ Two matrices A and B of same order are said to be **equivalent** if one can be obtained from the other by the applications of elementary transformations ($A \sim B$).
- ✦ A non – zero matrix is in a **row – echelon form** if all zero rows occur as bottom rows of the matrix and if the first non – zero element in any lower row occurs to the right of the first non – zero entry in the higher row.
- ✦ The **rank** of a matrix A is defined as the order of a highest order non – vanishing minor of the matrix A [$\rho(A)$].
- ✦ The **rank** of a non – zero matrix is equal to the number of non – zero rows in a row – echelon form of the matrix.
- ✦ An **elementary matrix** is a matrix which is obtained from an identity matrix by applying only one elementary transformation. Every non-singular matrix can be transformed to an identity matrix by a sequence of elementary row operations.
- ✦ A system of linear equations having atleast one solution is said to be **consistent**.
- ✦ A system of linear equations having no solutions is said to be **inconsistent**.

IMPORTANT FORMULAE TO REMEMBER

- ✦ Co – factor of a_{ij} is $A_{ij} = (-1)^{i+j} M_{ij}$, where M_{ij} is the minor of a_{ij}
- ✦ For every square matrix A of order n , $A (\text{adj } A) = (\text{adj } A)A = |A| I_n$
 $AA^{-1} = A^{-1}A = I_n$
- ✦ If A is non – Singular then
 - (i) $|A^{-1}| = \frac{1}{|A|}$ (ii) $(A^T)^{-1} = (A^{-1})^T$
 - (iii) $(\lambda A^{-1}) = \frac{1}{\lambda} A^{-1}$ where λ is a non – zero scalar.

Reversal law for inverses :

- ✦ $(AB)^{-1} = B^{-1}A^{-1}$ where A, B are non – singular matrices of same order.

Law of double inverse :

- ✦ If A is non - singular, A^{-1} is also non – singular and $(A^{-1})^{-1} = A$.
- ✦ If A is a non - singular square matrix of order n , then
 - (i) $(\text{adj } A)^{-1} = \text{adj } (A^{-1}) = \frac{1}{|A|} \cdot A$
 - (ii) $|\text{adj } A| = |A|^{n-1}$
 - (iii) $\text{adj } (\text{adj } A) = |A|^{n-2}A$
 - (iv) $\text{adj } (\lambda A) = \lambda^{n-1} \text{adj } (A)$ where λ is a non – zero scalar
 - (v) $|\text{adj } (\text{adj } A)| = |A|^{(n-1)^2}$
 - (vi) $(\text{adj } A)^T = \text{adj } (A^T)$
- ✦ If a matrix contains at least one non – zero element, then $\rho(A) \geq 1$.
- ✦ The rank of identity matrix I_n is n . If A is an $m \times n$ matrix then $\rho(A) \leq \min \{m, n\}$.
- ✦ A square matrix A of order n is invertible if and only if $\rho(A) = n$.
- ✦ Transforming a non-singular matrix A to the form I_n , by applying row operations is called Gauss – Jordan method.

Matrix – Inversion method :

- ✦ The solution for $AX = B$ is $X = A^{-1}B$ where A and B are square matrices of same order and non – singular

Cramer's Rule :

- ✦ If $\Delta = 0$, Cramer's rule cannot be applied $x_1 = \frac{\Delta_1}{\Delta}, x_2 = \frac{\Delta_2}{\Delta}, x_3 = \frac{\Delta_3}{\Delta}$

Gaussian Elimination method :

Transform the augmented matrix of the system of linear equations into row – echelon form and then solve by back substitution method.

Rouches capelli Theorem :

A system of equations $AX = B$ is consistent if and if $\rho(A) = \rho([A|B])$

- (i) If $\rho(A) = \rho([A|B]) = n$, the number of unknowns, then the system is consistent and has a unique solution.
- (ii) If $\rho(A) = \rho([A|B]) = n - k, k \neq 0$ then the system is consistent and has infinitely many solutions.
- (iii) If $\rho(A) \neq \rho([A|B])$, then the system is inconsistent and has no solution.

Homogeneous system of linear equations :

(i) If $\rho(A) = \rho([A|B]) = n$, then the system has a unique solution which is the trivial solution for trivial solution, $|A| \neq 0$

(ii) If $\rho(A) = \rho([A|B]) < n$, the system has a non – trivial solution.

For non – trivial solution, $|A| = 0$.

$$\star \quad A^{-1} = \pm \frac{1}{\sqrt{|adj A|}} \cdot adj A \quad \star \quad A = \pm \frac{1}{\sqrt{|adj A|}} \cdot adj (adj A)$$

EXERCISE 1.1**1. Find the adjoint of the following :**

(i) $\begin{bmatrix} -3 & 4 \\ 6 & 2 \end{bmatrix}$ (ii) $\begin{bmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{bmatrix}$ (iii) $\frac{1}{3} \begin{bmatrix} 2 & 2 & 1 \\ -2 & 1 & 2 \\ 1 & -2 & 2 \end{bmatrix}$

Sol. (i) Let $A = \begin{pmatrix} -3 & 4 \\ 6 & 2 \end{pmatrix}$
 $adj A = \begin{pmatrix} 2 & -4 \\ -6 & -3 \end{pmatrix}$

[Interchange the elements in the leading diagonal and change the sign of the elements in off diagonal]

(ii) Let $A = \begin{pmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{pmatrix}$

$$adj A = \begin{pmatrix} + \begin{vmatrix} 4 & 1 \\ 7 & 2 \end{vmatrix} - \begin{vmatrix} 3 & 1 \\ 3 & 2 \end{vmatrix} + \begin{vmatrix} 3 & 4 \\ 3 & 7 \end{vmatrix} \\ - \begin{vmatrix} 3 & 1 \\ 7 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 3 & 2 \end{vmatrix} - \begin{vmatrix} 2 & 3 \\ 3 & 7 \end{vmatrix} \\ + \begin{vmatrix} 3 & 1 \\ 4 & 1 \end{vmatrix} - \begin{vmatrix} 2 & 1 \\ 3 & 1 \end{vmatrix} + \begin{vmatrix} 2 & 3 \\ 3 & 4 \end{vmatrix} \end{pmatrix}^T$$

$$= \begin{bmatrix} + (8-7) - (6-3) + (21-12) \\ - (6-7) + (4-3) - (14-9) \\ + (3-4) - (2-3) + (8-9) \end{bmatrix}^T = \begin{bmatrix} 1 & -3 & 9 \\ 1 & 1 & -5 \\ -1 & 1 & -1 \end{bmatrix}^T$$

$$adj A = \begin{bmatrix} 1 & 1 & -1 \\ -3 & 1 & 1 \\ 9 & -5 & -1 \end{bmatrix}$$

(iii) Let $A = \begin{bmatrix} 2 & 2 & 1 \\ -2 & 1 & 2 \\ 1 & -2 & 2 \end{bmatrix}$ and $\lambda = \frac{1}{3}$

Since $adj (\lambda A) = \lambda^{n-1} (adj A)$

we get $adj \left(\frac{1}{3} \begin{bmatrix} 2 & 2 & 1 \\ -2 & 1 & 2 \\ 1 & -2 & 2 \end{bmatrix} \right) = \left(\frac{1}{3} \right)^2$

$$adj \begin{pmatrix} 2 & 2 & 1 \\ -2 & 1 & 2 \\ 1 & -2 & 2 \end{pmatrix}$$

∴ Required adjoint matrix

$$\begin{aligned} &= \frac{1}{9} \begin{bmatrix} + \begin{vmatrix} 1 & 2 \\ -2 & 2 \end{vmatrix} - \begin{vmatrix} -2 & 2 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} -2 & 1 \\ 1 & -2 \end{vmatrix} \\ - \begin{vmatrix} 2 & 1 \\ -2 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} - \begin{vmatrix} 2 & 2 \\ 1 & -2 \end{vmatrix} \\ + \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} - \begin{vmatrix} 2 & 1 \\ -2 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 2 \\ -2 & 1 \end{vmatrix} \end{bmatrix}^T \\ &= \frac{1}{9} \begin{bmatrix} (2+4) - (-4-2) + (4-1) \\ - (4+2) + (4-1) - (-4-2) \\ + (4-1) - (4+2) + (2+4) \end{bmatrix}^T \\ &= \frac{1}{9} \begin{bmatrix} 6 & 6 & 3 \\ -6 & 3 & 6 \\ 3 & -6 & 6 \end{bmatrix}^T = \frac{1}{9} \begin{bmatrix} 6 & -6 & 3 \\ 6 & 3 & -6 \\ 3 & 6 & 6 \end{bmatrix} \\ &= \frac{3}{9} \begin{bmatrix} 2 & -2 & 1 \\ 2 & 1 & -2 \\ 1 & 2 & 2 \end{bmatrix} \end{aligned}$$

[Taking 3 common from each entry]

$$= \frac{1}{3} \begin{bmatrix} 2 & -2 & 1 \\ 2 & 1 & -2 \\ 1 & 2 & 2 \end{bmatrix}$$

2. Find the inverse (if it exists) of the following :

(i) $\begin{bmatrix} -2 & 4 \\ 1 & -3 \end{bmatrix}$ (ii) $\begin{bmatrix} 5 & 1 & 1 \\ 1 & 5 & 1 \\ 1 & 1 & 5 \end{bmatrix}$ (iii) $\begin{bmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{bmatrix}$

Sol. (i) Let $A = \begin{bmatrix} -2 & 4 \\ 1 & -3 \end{bmatrix}$

$$|A| = \begin{vmatrix} -2 & 4 \\ 1 & -3 \end{vmatrix} = 6 - 4 = 2 \neq 0$$

Since A is non-singular, A^{-1} exists

$$A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$\text{Now, adj } A = \begin{bmatrix} -3 & -4 \\ -1 & -2 \end{bmatrix}$$

[Inter change the entries in leading diagonal and change the sign of elements in the off diagonal]

$$\therefore A^{-1} = \frac{1}{2} \begin{bmatrix} -3 & -4 \\ -1 & -2 \end{bmatrix}$$

$$(ii) \text{ Let } A = \begin{bmatrix} 5 & 1 & 1 \\ 1 & 5 & 1 \\ 1 & 1 & 5 \end{bmatrix}$$

Expanding along R_1 ,

$$\begin{aligned} |A| &= 5 \begin{vmatrix} 5 & 1 \\ 1 & 5 \end{vmatrix} - 1 \begin{vmatrix} 1 & 1 \\ 1 & 1 \end{vmatrix} + 1 \begin{vmatrix} 1 & 5 \\ 1 & 1 \end{vmatrix} \\ &= 5(25 - 1) - 1(5 - 1) + 1(1 - 5) \\ &= 5(24) - 1(4) + 1(-4) \\ &= 120 - 4 - 4 = 120 - 8 = 112 \neq 0 \end{aligned}$$

Since A is non singular, A^{-1} exists.

$$\begin{aligned} \text{adj } A &= \begin{bmatrix} + \begin{vmatrix} 5 & 1 \\ 1 & 5 \end{vmatrix} - \begin{vmatrix} 1 & 1 \\ 1 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 5 \\ 1 & 1 \end{vmatrix} \\ - \begin{vmatrix} 1 & 1 \\ 1 & 5 \end{vmatrix} + \begin{vmatrix} 5 & 1 \\ 1 & 1 \end{vmatrix} - \begin{vmatrix} 5 & 1 \\ 1 & 1 \end{vmatrix} \\ + \begin{vmatrix} 1 & 1 \\ 5 & 1 \end{vmatrix} - \begin{vmatrix} 5 & 1 \\ 1 & 1 \end{vmatrix} + \begin{vmatrix} 5 & 1 \\ 1 & 5 \end{vmatrix} \end{bmatrix}^T \\ &= \begin{bmatrix} +(25-1)-(5-1)+(1-5) \\ -(5-1)+(25-1)-(5-1) \\ +(1-5)-(5-1)+(25-1) \end{bmatrix}^T \\ &= \begin{bmatrix} 24 & -4 & -4 \\ -4 & 24 & -4 \\ -4 & -4 & 24 \end{bmatrix}^T = \begin{bmatrix} 24 & -4 & -4 \\ -4 & 24 & -4 \\ -4 & -4 & 24 \end{bmatrix} \end{aligned}$$

Taking 4 common from every entry we get,

$$\begin{aligned} \text{adj } A &= 4 \begin{bmatrix} 6 & -1 & -1 \\ -1 & 6 & -1 \\ -1 & -1 & 6 \end{bmatrix} \\ \therefore A^{-1} &= \frac{1}{|A|} \text{adj } A = \frac{1}{112} \cdot 4 \begin{bmatrix} 6 & -1 & -1 \\ -1 & 6 & -1 \\ -1 & -1 & 6 \end{bmatrix} \end{aligned}$$

$$= \frac{1}{28} \begin{bmatrix} 6 & -1 & -1 \\ -1 & 6 & -1 \\ -1 & -1 & 6 \end{bmatrix}$$

$$(iii) \text{ Let } A = \begin{bmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{bmatrix}$$

Expanding along R_1 we get,

$$\begin{aligned} |A| &= 2 \begin{vmatrix} 4 & 1 \\ 7 & 2 \end{vmatrix} - 3 \begin{vmatrix} 3 & 1 \\ 3 & 2 \end{vmatrix} + 1 \begin{vmatrix} 3 & 4 \\ 3 & 7 \end{vmatrix} \\ &= 2(8 - 7) - 3(6 - 3) + 1(21 - 12) \\ &= 2(1) - 3(3) + 1(9) \\ &= 2 - 9 + 9 = 2 \neq 0 \end{aligned}$$

Since A is a non-singular matrix, A^{-1} exists

$$\text{adj } A = \begin{bmatrix} + \begin{vmatrix} 4 & 1 \\ 7 & 2 \end{vmatrix} - \begin{vmatrix} 3 & 1 \\ 3 & 2 \end{vmatrix} + \begin{vmatrix} 3 & 4 \\ 3 & 7 \end{vmatrix} \\ - \begin{vmatrix} 3 & 1 \\ 7 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 3 & 2 \end{vmatrix} - \begin{vmatrix} 2 & 3 \\ 3 & 7 \end{vmatrix} \\ + \begin{vmatrix} 3 & 1 \\ 4 & 1 \end{vmatrix} - \begin{vmatrix} 2 & 1 \\ 3 & 1 \end{vmatrix} + \begin{vmatrix} 2 & 3 \\ 3 & 4 \end{vmatrix} \end{bmatrix}^T$$

$$= \begin{bmatrix} +(8-7)-(6-3)+(21-12) \\ -(6-7)+(4-3)-(14-9) \\ +(3-4)-(2-3)+(8-9) \end{bmatrix}^T = \begin{bmatrix} 1 & -3 & 9 \\ 1 & 1 & -5 \\ -1 & 1 & -1 \end{bmatrix}^T$$

$$\text{adj } A = \begin{bmatrix} 1 & 1 & -1 \\ -3 & 1 & 1 \\ 9 & -5 & -1 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$\Rightarrow A^{-1} = \frac{1}{2} \begin{bmatrix} 1 & 1 & -1 \\ -3 & 1 & 1 \\ 9 & -5 & -1 \end{bmatrix}$$

$$3. \text{ If } F(\alpha) = \begin{bmatrix} \cos \alpha & 0 & \sin \alpha \\ 0 & 1 & 0 \\ -\sin \alpha & 0 & \cos \alpha \end{bmatrix}, \text{ show that}$$

$$[F(\alpha)]^{-1} = F(-\alpha) \quad [\text{Hy - 2019; FRT - 2022; Mar - 2023}]$$

$$\text{Sol. Given that } F(\alpha) = \begin{bmatrix} \cos \alpha & 0 & \sin \alpha \\ 0 & 1 & 0 \\ -\sin \alpha & 0 & \cos \alpha \end{bmatrix}$$

Expanding along R_1 we get,

$$|F(\alpha)| = \cos \alpha \begin{vmatrix} 1 & 0 \\ 0 & \cos \alpha \end{vmatrix} - 0 + \sin \alpha \begin{vmatrix} 0 & 1 \\ -\sin \alpha & 0 \end{vmatrix}$$

$$= \cos \alpha (\cos - 0) + \sin \alpha (0 + \sin \alpha)$$

$$= \cos^2 + \sin^2 \alpha = 1 \neq 0$$

Since $F(\alpha)$ is a non-singular matrix, $[F(\alpha)]^{-1}$ exists.

Now, $\text{adj } (F(\alpha)) =$

$$\begin{bmatrix} + \begin{vmatrix} 1 & 0 \\ 0 & \cos \alpha \end{vmatrix} - \begin{vmatrix} 0 & 0 \\ \sin \alpha & \cos \alpha \end{vmatrix} + \begin{vmatrix} 0 & 1 \\ -\sin \alpha & 0 \end{vmatrix} \\ - \begin{vmatrix} 0 & \sin \alpha \\ 0 & \cos \alpha \end{vmatrix} + \begin{vmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{vmatrix} - \begin{vmatrix} \cos \alpha & 0 \\ -\sin \alpha & 0 \end{vmatrix} \\ + \begin{vmatrix} 0 & \sin \alpha \\ 1 & 0 \end{vmatrix} - \begin{vmatrix} \cos \alpha & \sin \alpha \\ 0 & 0 \end{vmatrix} + \begin{vmatrix} \cos \alpha & 0 \\ 0 & 1 \end{vmatrix} \end{bmatrix}^T$$

$$= \begin{bmatrix} +(\cos \alpha - 0) & -(0) & +(0 + \sin \alpha) \\ -(0) & +(\cos^2 \alpha + \sin^2 \alpha) & -(0) \\ +(0 - \sin \alpha) & -(0) & +(\cos - 0) \end{bmatrix}^T$$

$$= \begin{bmatrix} \cos \alpha & 0 & +\sin \alpha \\ 0 & 1 & 0 \\ -\sin \alpha & 0 & \cos \alpha \end{bmatrix}^T = \begin{bmatrix} \cos \alpha & 0 & -\sin \alpha \\ 0 & 1 & 0 \\ \sin \alpha & 0 & \cos \alpha \end{bmatrix}$$

$$\therefore F(\alpha)^{-1} = \frac{1}{|F(\alpha)|} \text{adj } (F(\alpha))$$

$$[F(\alpha)]^{-1} = \frac{1}{1} \begin{bmatrix} \cos \alpha & 0 & -\sin \alpha \\ 0 & 1 & 0 \\ \sin \alpha & 0 & \cos \alpha \end{bmatrix}$$

$$= \begin{bmatrix} \cos \alpha & 0 & -\sin \alpha \\ 0 & 1 & 0 \\ \sin \alpha & 0 & \cos \alpha \end{bmatrix} \quad \dots (1)$$

$$\text{Now, } F(-\alpha) = \begin{bmatrix} \cos(-\alpha) & 0 & \sin(-\alpha) \\ 0 & 1 & 0 \\ -\sin(-\alpha) & 0 & \cos(-\alpha) \end{bmatrix}$$

$$= \begin{bmatrix} \cos \alpha & 0 & -\sin \alpha \\ 0 & 1 & 0 \\ \sin \alpha & 0 & \cos \alpha \end{bmatrix} \quad \dots (2)$$

$[\because \cos \alpha$ is an even function, $\cos(-\alpha) = \cos \alpha$ and $\sin \alpha$ is an odd function, $\sin(-\alpha) = -\sin \alpha]$

From (1) and (2)

$$[F(\alpha)]^{-1} = F(-\alpha)$$

Hence proved.

4. If $A = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$, show that $A^2 - 3A - 7I_2 = 0_2$.

Hence find A^{-1} .

Sol. Given $A = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$

$$A^2 = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} = \begin{bmatrix} 25-3 & 15-6 \\ -5+2 & -3+4 \end{bmatrix}$$

$$= \begin{bmatrix} 22 & 9 \\ -3 & 1 \end{bmatrix} \quad \therefore A^2 - 3A - 7I_2$$

$$= \begin{bmatrix} 22 & 9 \\ -3 & 1 \end{bmatrix} - 3 \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} - 7 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 22-15-7 & 9-9+0 \\ -3+3+0 & 1+6-7 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0_2$$

Hence proved.

$$\therefore A^2 - 3A - 7I_2 = 0$$

Post-multiplying by A^{-1} we get,

$$A^2 \cdot A^{-1} - 3AA^{-1} - 7I_2 A^{-1} = 0 \cdot A^{-1}$$

$$\Rightarrow A(AA^{-1}) - 3(AA^{-1}) - 7(A^{-1}) = 0$$

$$[\because I_2 A^{-1} = A^{-1} \text{ and } (0)A^{-1} = 0]$$

$$\Rightarrow AI - 3I - 7A^{-1} = 0$$

$$[\because AA^{-1} = I]$$

$$\Rightarrow AI - 3I = 7A^{-1}$$

$$\Rightarrow A^{-1} = \frac{1}{7} [A - 3I] \quad [\because AI = A]$$

$$\Rightarrow A^{-1} = \frac{1}{7} \left[\begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} - 3 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right]$$

$$\Rightarrow A^{-1} = \frac{1}{7} \begin{bmatrix} 5-3 & 3-0 \\ -1-0 & -2-3 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 2 & 3 \\ -1 & -5 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{7} \begin{bmatrix} 2 & 3 \\ -1 & -5 \end{bmatrix}$$

5. If $A = \frac{1}{9} \begin{bmatrix} -8 & 1 & 4 \\ 4 & 4 & 7 \\ 1 & -8 & 4 \end{bmatrix}$, prove that $A^{-1} = A^T$.

Sol.

To prove $A^{-1} = A^T$

$$AA^{-1} = AA^T$$

It is enough to prove $AA^T = I$

$$AA^T = \frac{1}{9} \begin{bmatrix} -8 & 1 & 4 \\ 4 & 4 & 7 \\ 1 & -8 & 4 \end{bmatrix} \cdot \frac{1}{9} \begin{bmatrix} -8 & 1 & 4 \\ 1 & 4 & -8 \\ 4 & 7 & 4 \end{bmatrix}$$

$$= \frac{1}{81} \begin{bmatrix} 64+1+16 & -32+4+28 & -8-8+16 \\ -32+4+28 & 16+16+49 & 4-32-28 \\ -8-8+16 & 4-32+28 & 1+64+16 \end{bmatrix}$$

$$= \frac{1}{81} \begin{bmatrix} 81 & 0 & 0 \\ 0 & 81 & 0 \\ 0 & 0 & 81 \end{bmatrix} = \frac{81}{81} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

Therefore $A^{-1} = A^T$

6. If $A = \begin{bmatrix} 8 & -4 \\ -5 & 3 \end{bmatrix}$, verify that $A(\text{adj } A) = (\text{adj } A)A = |A| I_2$. [Sep. - 2020; Aug. - 2021]

Sol. Given $A = \begin{bmatrix} 8 & -4 \\ -5 & 3 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 3 & 4 \\ 5 & 8 \end{bmatrix}$$

[Interchange the elements in the leading diagonal and change the sign of the elements in the off diagonal]

$$|A| = 24 - 20 = 4$$

$$\therefore A(\text{adj } A) = \begin{bmatrix} 8 & -4 \\ -5 & 3 \end{bmatrix} \begin{bmatrix} 3 & 4 \\ 5 & 8 \end{bmatrix}$$

$$= \begin{bmatrix} 24-20 & 32-32 \\ -15+15 & -20+24 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \quad \dots(1)$$

$$(\text{adj } A)(A) = \begin{bmatrix} 3 & 4 \\ 5 & 8 \end{bmatrix} \begin{bmatrix} 8 & -4 \\ -5 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 24-20 & -12+12 \\ 40-40 & -20+24 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \quad \dots(2)$$

$$|A| I_2 = 4 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \quad \dots(3)$$

From (1), (2) and (3), it is proved that

$A(\text{adj } A) = (\text{adj } A)A = |A| I_2$ is verified.

7. If $A = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & -3 \\ 5 & 2 \end{bmatrix}$, verify that $(AB)^{-1} = B^{-1}A^{-1}$. [Qy. - 2023]

Sol. Given $A = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & -3 \\ 5 & 2 \end{bmatrix}$

$$AB = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix} \begin{bmatrix} -1 & -3 \\ 5 & 2 \end{bmatrix} = \begin{bmatrix} -3+10 & -9+4 \\ -7+25 & -21+10 \end{bmatrix}$$

$$= \begin{bmatrix} 7 & -5 \\ 18 & -11 \end{bmatrix}$$

$$|AB| = -77 + 90 = 13 \neq 0 \Rightarrow (AB)^{-1} \text{ exists}$$

$$|A| = 15 - 14 = 1 \neq 0 \Rightarrow A^{-1} \text{ exists}$$

$$|B| = -2 + 15 = 13 \neq 0 \Rightarrow B^{-1} \text{ exists}$$

$$(AB)^{-1} = \frac{1}{|AB|} \text{adj } (AB) = \frac{1}{13} \begin{pmatrix} -11 & 5 \\ -18 & 7 \end{pmatrix} \dots(1)$$

$$B^{-1} = \frac{1}{|B|} (\text{adj } B) = \frac{1}{13} \begin{pmatrix} 2 & 3 \\ -5 & -1 \end{pmatrix}$$

$$A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{1} \begin{pmatrix} 5 & -2 \\ -7 & 3 \end{pmatrix}$$

$$= \begin{pmatrix} 5 & -2 \\ -7 & 3 \end{pmatrix}$$

$$\therefore B^{-1}A^{-1} = \frac{1}{13} \begin{pmatrix} 2 & 3 \\ -5 & -1 \end{pmatrix} \begin{pmatrix} 5 & -2 \\ -7 & 3 \end{pmatrix}$$

$$= \frac{1}{13} \begin{pmatrix} 10-21 & -4+9 \\ -25+7 & 10-3 \end{pmatrix}$$

$$= \frac{1}{13} \begin{pmatrix} -11 & 5 \\ -18 & 7 \end{pmatrix} \quad \dots(2)$$

From (1) and (2) it is prove that

$$(AB)^{-1} = B^{-1}A^{-1}.$$

8. If $\text{adj } (A) = \begin{bmatrix} 2 & -4 & 2 \\ -3 & 12 & -7 \\ -2 & 0 & 2 \end{bmatrix}$, find A.

Sol. Given $\text{adj } A = \begin{bmatrix} 2 & -4 & 2 \\ -3 & 12 & -7 \\ -2 & 0 & 2 \end{bmatrix}$

$$\text{We know that } A = \pm \frac{1}{\sqrt{|\text{adj } A|}} \text{adj } (\text{adj } A) \quad \dots(1)$$

$$|\text{adj } A| = 2 \begin{vmatrix} 12 & -7 \\ 0 & 2 \end{vmatrix} + 4 \begin{vmatrix} -3 & -7 \\ -2 & 2 \end{vmatrix} + 2 \begin{vmatrix} -3 & 12 \\ -2 & 0 \end{vmatrix}$$

[Expanded along R_1]

$$= 2(24-0) + 4(-6-14) + 2(0+24)$$

$$= 2(24) + 4(-20) + 2(24) = 48 - 80 + 48$$

$$= 96 - 80 = 16 \quad \dots(2)$$

Now, $\text{adj } (\text{adj } A)$

$$\begin{bmatrix} + \begin{vmatrix} 12 & -7 \\ 0 & 2 \end{vmatrix} - \begin{vmatrix} -3 & -7 \\ -2 & 2 \end{vmatrix} + \begin{vmatrix} -3 & 12 \\ -2 & 0 \end{vmatrix} \\ - \begin{vmatrix} -4 & 2 \\ 0 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 2 \\ -2 & 2 \end{vmatrix} - \begin{vmatrix} 2 & -4 \\ -2 & 0 \end{vmatrix} \\ + \begin{vmatrix} -4 & 2 \\ 12 & -7 \end{vmatrix} - \begin{vmatrix} 2 & 2 \\ -3 & -7 \end{vmatrix} + \begin{vmatrix} 2 & -4 \\ -3 & 12 \end{vmatrix} \end{bmatrix}^T$$

$$\begin{aligned}
 &= \begin{bmatrix} +(24-0)-(-6-14)+(0+24) \\ -(-8-0)+(4+4)-(0-8) \\ +(28-24)-(-14+6)+(24-12) \end{bmatrix}^T \\
 &= \begin{bmatrix} 24 & 20 & 24 \\ 8 & 8 & 8 \\ 4 & 8 & 12 \end{bmatrix}^T = \begin{bmatrix} 24 & 8 & 4 \\ 20 & 8 & 8 \\ 24 & 8 & 12 \end{bmatrix} \\
 &= 4 \begin{bmatrix} 6 & 2 & 1 \\ 5 & 2 & 2 \\ 6 & 2 & 3 \end{bmatrix} \quad \dots(3)
 \end{aligned}$$

Substituting (2) and (3) in (1) we get,

$$A = \pm \frac{1}{\sqrt{16}} \cdot 4 \begin{bmatrix} 6 & 2 & 1 \\ 5 & 2 & 2 \\ 6 & 2 & 3 \end{bmatrix}$$

$$A = \pm \frac{4}{4} \begin{bmatrix} 6 & 2 & 1 \\ 5 & 2 & 2 \\ 6 & 2 & 3 \end{bmatrix} = \pm \begin{bmatrix} 6 & 2 & 1 \\ 5 & 2 & 2 \\ 6 & 2 & 3 \end{bmatrix}$$

9. If $\text{adj}(A) = \begin{bmatrix} 0 & -2 & 0 \\ 6 & 2 & -6 \\ -3 & 0 & 6 \end{bmatrix}$, find A^{-1} .

[PTA - 6; FRT -2022; June & Qy. - 2023]

Sol. Given $\text{adj}(A) = \begin{bmatrix} 0 & -2 & 0 \\ 6 & 2 & -6 \\ -3 & 0 & 6 \end{bmatrix}$

We know that $A^{-1} = \pm \frac{1}{\sqrt{|\text{adj} A|}} (\text{adj} A) \dots(1)$

$$\begin{aligned}
 |\text{adj} A| &= 0 + 2 \begin{vmatrix} 6 & -6 \\ -3 & 6 \end{vmatrix} + 0 \\
 &= 2(36 - 18) = 2(18) = 36
 \end{aligned}$$

[Expanded along R_1]

$$\begin{aligned}
 \therefore A^{-1} &= \pm \frac{1}{\sqrt{36}} \begin{bmatrix} 0 & -2 & 0 \\ 6 & 2 & -6 \\ -3 & 0 & 6 \end{bmatrix} \\
 &= \pm \frac{1}{6} \begin{bmatrix} 0 & -2 & 0 \\ 6 & 2 & -6 \\ -3 & 0 & 6 \end{bmatrix}
 \end{aligned}$$

10. Find $\text{adj}(\text{adj}(A))$ if $\text{adj} A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix}$.

Sol. Given $\text{adj} A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix}$

Now $\text{adj}(\text{adj} A) = \begin{bmatrix} + \begin{vmatrix} 2 & 0 \\ 0 & 1 \end{vmatrix} - \begin{vmatrix} 0 & 0 \\ -1 & 1 \end{vmatrix} + \begin{vmatrix} 0 & 2 \\ -1 & 0 \end{vmatrix} \\ - \begin{vmatrix} 0 & 1 \\ 0 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 1 \\ -1 & 1 \end{vmatrix} - \begin{vmatrix} 1 & 0 \\ -1 & 0 \end{vmatrix} \\ + \begin{vmatrix} 0 & 1 \\ 2 & 0 \end{vmatrix} - \begin{vmatrix} 1 & 1 \\ 0 & 0 \end{vmatrix} + \begin{vmatrix} 1 & 0 \\ 0 & 2 \end{vmatrix} \end{bmatrix}^T$

$$\begin{aligned}
 &= \begin{bmatrix} +(2-0) & -(0) & +(0+2) \\ - (0) & +(1+1) & - (0) \\ +(0-2) & - (0) & +(2-0) \end{bmatrix}^T = \begin{bmatrix} 2 & 0 & 2 \\ 0 & 2 & 0 \\ -2 & 0 & 2 \end{bmatrix}^T \\
 \text{adj}(\text{adj} A) &= \begin{bmatrix} 2 & 0 & -2 \\ 0 & 2 & 0 \\ 2 & 0 & 2 \end{bmatrix}
 \end{aligned}$$

11. $A = \begin{bmatrix} 1 & \tan x \\ -\tan x & 1 \end{bmatrix}$, show that

$$A^T A^{-1} = \begin{bmatrix} \cos 2x & -\sin 2x \\ \sin 2x & \cos 2x \end{bmatrix}.$$

Sol. $A = \begin{bmatrix} 1 & \tan x \\ -\tan x & 1 \end{bmatrix}$

$$|A| = \begin{vmatrix} 1 & \tan x \\ -\tan x & 1 \end{vmatrix} = 1 + \tan^2 x = \sec^2 x$$

$$\text{adj} A = \begin{bmatrix} 1 & -\tan x \\ \tan x & 1 \end{bmatrix}$$

We know, $A^{-1} = \frac{1}{|A|} \text{Adj} A$

$$\begin{aligned}
 &= \frac{1}{\sec^2 x} \begin{bmatrix} 1 & -\tan x \\ \tan x & 1 \end{bmatrix} = \cos^2 x \begin{bmatrix} 1 & -\tan x \\ \tan x & 1 \end{bmatrix} \\
 &= \cos^2 x \begin{bmatrix} 1 & -\frac{\sin x}{\cos x} \\ \frac{\sin x}{\cos x} & 1 \end{bmatrix} \\
 A^{-1} &= \begin{bmatrix} \cos^2 x & -\cos x \sin x \\ \cos x \sin x & \cos^2 x \end{bmatrix}
 \end{aligned}$$

$$A = \begin{bmatrix} 1 & \tan x \\ -\tan x & 1 \end{bmatrix}$$

$$A^T = \begin{bmatrix} 1 & -\frac{\sin x}{\cos x} \\ \frac{\sin x}{\cos x} & 1 \end{bmatrix}$$

$$A^T A^{-1} = \begin{bmatrix} 1 & -\frac{\sin x}{\cos x} \\ \frac{\sin x}{\cos x} & 1 \end{bmatrix} \begin{bmatrix} \cos^2 x & -\cos x \sin x \\ \cos x \sin x & \cos^2 x \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2 x - \sin^2 x & -2 \sin x \cos x \\ 2 \sin x \cos x & \cos^2 x - \sin^2 x \end{bmatrix} = \begin{bmatrix} \cos 2x & -\sin 2x \\ \sin 2x & \cos 2x \end{bmatrix}$$

12. Find the matrix A for which

$$A \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} = \begin{bmatrix} 14 & 7 \\ 7 & 7 \end{bmatrix}$$

Sol. Given $A \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} = \begin{bmatrix} 14 & 7 \\ 7 & 7 \end{bmatrix}$

Let $B = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$ and $C = \begin{bmatrix} 14 & 7 \\ 7 & 7 \end{bmatrix}$

$$\therefore AB = C$$

Post multiply by B^{-1} we get

$$A(BB^{-1}) = CB^{-1}$$

$$\Rightarrow A = CB^{-1} \quad [\because BB^{-1} = I]$$

$$|B| = \begin{vmatrix} 5 & 3 \\ -1 & -2 \end{vmatrix} = -10 + 3 = -7 \neq 0$$

$\therefore B^{-1}$ exists

$$B^{-1} = \frac{1}{|B|} \text{adj } B = \frac{-1}{7} \begin{bmatrix} -2 & -3 \\ 1 & 5 \end{bmatrix}$$

$$A = CB^{-1}$$

$$= \begin{bmatrix} 14 & 7 \\ 7 & 7 \end{bmatrix} \cdot \left(\frac{-1}{7} \right) \begin{bmatrix} -2 & -3 \\ 1 & 5 \end{bmatrix}$$

$$= 7 \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \left(\frac{-1}{7} \right) \begin{bmatrix} -2 & -3 \\ 1 & 5 \end{bmatrix}$$

$$= - \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} -2 & -3 \\ 1 & 5 \end{bmatrix} = - \begin{bmatrix} -4+1 & -6+5 \\ -2+1 & -3+5 \end{bmatrix}$$

$$= - \begin{bmatrix} -3 & -1 \\ -1 & 2 \end{bmatrix} \Rightarrow A = \begin{bmatrix} 3 & 1 \\ 1 & -2 \end{bmatrix}$$

13. Given $A = \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 3 & -2 \\ 1 & 1 \end{bmatrix}$ and

$C = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$, find a matrix X such that $AXB = C$.

Sol. Given $A = \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 3 & -2 \\ 1 & 1 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$

$$\text{Also, } A X B = C$$

Pre-multiply by A^{-1} we get,

$$(A^{-1} A) X B = A^{-1} C$$

$$\Rightarrow X B = A^{-1} \cdot C \quad [\because A^{-1} A = I]$$

Post Multiply by B^{-1} we get

$$(X B) B^{-1} = (A^{-1} C) B^{-1} \Rightarrow X = (A^{-1} C) B^{-1}$$

$$|A| = \begin{vmatrix} 1 & -1 \\ 2 & 0 \end{vmatrix} = 0 + 2 = 2 \neq 0$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{2} \begin{bmatrix} 0 & 1 \\ -2 & 1 \end{bmatrix}$$

$$|B| = \begin{vmatrix} 3 & -2 \\ 1 & 1 \end{vmatrix} = 3 + 2 = 5 \neq 0$$

$$\therefore B^{-1} = \frac{1}{|B|} \text{adj } B = \frac{1}{5} \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix}$$

$$A^{-1} C = \frac{1}{2} \begin{bmatrix} 0 & 1 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} 0+2 & 0+2 \\ -2+2 & -2+2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 2 & 2 \\ 0 & 0 \end{bmatrix}$$

$$= \frac{1}{2} (2) \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\therefore X = (A^{-1} C) \cdot B^{-1}$$

$$= \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \frac{1}{5} \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix}$$

$$= \frac{1}{5} \begin{bmatrix} 1-1 & 2+3 \\ 0+0 & 0+0 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 0 & 5 \\ 0 & 0 \end{bmatrix}$$

$$= \frac{1}{5} (5) \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\therefore X = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

14. If $A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$, show that $A^{-1} = \frac{1}{2} (A^2 - 3I)$.

Sol.

Given $A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$

$$|A| = 0 - 1 \begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} + 1 \begin{vmatrix} 1 & 0 \\ 1 & 1 \end{vmatrix}$$

$$= -1(0-1) + 1(1-0) = 1 + 1 = 2$$

$$|A| = 2 \neq 0,$$

Hence A^{-1} exists

$$\begin{aligned} \text{adj } A &= \begin{bmatrix} + \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} & - \begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} & + \begin{vmatrix} 1 & 0 \\ 1 & 1 \end{vmatrix} \\ - \begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} & + \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} & - \begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix} \\ + \begin{vmatrix} 1 & 1 \\ 0 & 1 \end{vmatrix} & - \begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix} & + \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} \end{bmatrix}^T \\ &= \begin{bmatrix} (0-1) & -(0-1) & +(1-0) \\ -(0-1) & +(0-1) & -(0-1) \\ +(1-0) & -(0-1) & +(0-1) \end{bmatrix}^T \\ &= \begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix}^T = \begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix} \\ \therefore A^{-1} &= \frac{1}{2} \begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix} \quad \dots(1) \end{aligned}$$

$$\begin{aligned} \text{Now } A^2 &= \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0+1+1 & 0+0+1 & 0+1+0 \\ 0+0+1 & 1+0+1 & 1+0+0 \\ 0+1+0 & 1+0+0 & 1+1+0 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix} \\ A^2 - 3I &= \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix} - 3 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 2-3 & 1-0 & 1-0 \\ 1-0 & 2-3 & 1-0 \\ 1-0 & 1-0 & 2-3 \end{bmatrix} \\ &= \frac{1}{2} (A^2 - 3I) = \frac{1}{2} \begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix} \quad \dots (2) \end{aligned}$$

From (1) and (2), it is proved that $A^{-1} = \frac{1}{2} [A^2 - 3I]$

- 15. Decrypt the received encoded message $\begin{bmatrix} 2 & -3 \end{bmatrix} \begin{bmatrix} 20 & 4 \end{bmatrix}$ with the encryption matrix $\begin{bmatrix} -1 & -1 \\ 2 & 1 \end{bmatrix}$ and the decryption matrix as its inverse, where the system of codes are described by the numbers 1 – 26 to the letters A – Z respectively, and the number 0 to a blank space.**

Sol. Let the encryption matrix be $A = \begin{bmatrix} -1 & -1 \\ 2 & 1 \end{bmatrix}$

$$|A| = -1 + 2 = 1 \neq 0$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{1} \begin{bmatrix} 1 & 1 \\ -2 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ -2 & -1 \end{bmatrix}$$

Hence the decryption matrix is $\begin{bmatrix} 1 & 1 \\ -2 & -1 \end{bmatrix}$

Coded row matrix	Decoding matrix	Decoded row matrix
$\begin{bmatrix} 2 & -3 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -2 & -1 \end{bmatrix}$	$= \begin{bmatrix} 2+6 & 2+3 \end{bmatrix} = \begin{bmatrix} 8 & 5 \end{bmatrix}$
$\begin{bmatrix} 20 & 4 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -2 & -1 \end{bmatrix}$	$= \begin{bmatrix} 20-8 & 20-4 \end{bmatrix} = \begin{bmatrix} 12 & 16 \end{bmatrix}$

So, the sequence of decoded row matrices is

$\begin{bmatrix} 8 & 5 \end{bmatrix}, \begin{bmatrix} 12 & 16 \end{bmatrix}$

Now the 8th English alphabet is H.

5th English alphabet is E.

12th English alphabet is L.

and the 16th English alphabet is P.

Thus the receiver reads the message as “HELP”.

EXERCISE 1.2

- 1. Find the rank of the following matrices by minor method:**

(i) $\begin{bmatrix} 2 & -4 \\ -1 & 2 \end{bmatrix}$ (ii) $\begin{bmatrix} -1 & 3 \\ 4 & -7 \\ 3 & -4 \end{bmatrix}$

[Mar. - 2024]

(iii) $\begin{bmatrix} 1 & -2 & -1 & 0 \\ 3 & -6 & -3 & 1 \end{bmatrix}$ [PTA - 5]

(iv) $\begin{bmatrix} 1 & -2 & 3 \\ 2 & 4 & -6 \\ 5 & 1 & -1 \end{bmatrix}$ [July - 2022]

(v) $\begin{bmatrix} 0 & 1 & 2 & 1 \\ 0 & 2 & 4 & 3 \\ 8 & 1 & 0 & 2 \end{bmatrix}$ [Aug. - 2021]

Sol. (i) Let $A = \begin{bmatrix} 2 & -4 \\ -1 & 2 \end{bmatrix}$

A is a matrix of order 2×2

$$\therefore \rho(A) \leq \min(2, 2) = 2$$

The highest order of minor of A is 2

$$\text{It is } \begin{vmatrix} 2 & -4 \\ -1 & 2 \end{vmatrix} = 4 - 4 = 0$$

$$\text{So, } \rho(A) < 2$$

$$\text{Next consider the minor of order 1 } |2| = 2 \neq 0$$

$$\therefore \rho(A) = 1$$

$$(ii) \text{ Let } A = \begin{bmatrix} -1 & 3 \\ 4 & -7 \\ 3 & -4 \end{bmatrix}$$

A is a matrix of order 3×2

$$\therefore \rho(A) \leq 2$$

We find that there is a second order minor,

$$\begin{vmatrix} -1 & 3 \\ 4 & -7 \end{vmatrix} = 7 - 12 = -5 \neq 0$$

$$\therefore \rho(A) = 2.$$

$$(iii) \text{ Let } A = \begin{bmatrix} 1 & -2 & -1 & 0 \\ 3 & -6 & -3 & 1 \end{bmatrix}$$

A is a matrix of order (2×4)

$$\therefore \rho(A) \leq \min(2, 4) = 2$$

The highest order of minor of A is 2

$$\text{It is } \begin{vmatrix} 1 & -2 \\ 3 & -6 \end{vmatrix} = -6 + 6 = 0$$

$$\text{Also, } \begin{vmatrix} -1 & 0 \\ -3 & 1 \end{vmatrix} = -1 + 0 = -1 \neq 0.$$

$$\therefore \rho(A) = 2.$$

$$(iv) \text{ Let } A = \begin{bmatrix} 1 & -2 & 3 \\ 2 & 4 & -6 \\ 5 & 1 & -1 \end{bmatrix}$$

A is a matrix of order 3×3

$$\therefore \rho(A) \leq \min(3, 3) = 3$$

The highest order of minor of A is 3.

$$(I.E) \begin{vmatrix} 1 & -2 & 3 \\ 2 & 4 & -6 \\ 5 & 1 & -1 \end{vmatrix} = 1 \begin{vmatrix} 4 & -6 \\ 1 & -1 \end{vmatrix} + 2 \begin{vmatrix} 2 & -6 \\ 5 & -1 \end{vmatrix} + 3 \begin{vmatrix} 2 & 4 \\ 5 & 1 \end{vmatrix}$$

[Expanded along R_1]

$$= 1(-4 + 6) + 2(-2 + 30) + 3(2 - 20)$$

$$= 1(2) + 2(28) + 3(-18)$$

$$= 2 + 56 - 54 = 58 - 54 = 4 \neq 0$$

$$\therefore \rho(A) = 3.$$

$$(v) \text{ Let } A = \begin{bmatrix} 0 & 1 & 2 & 1 \\ 0 & 2 & 4 & 3 \\ 8 & 1 & 0 & 2 \end{bmatrix}$$

A is a matrix of order 3×4

$$\therefore \rho(A) \leq \min(3, 4) = 3$$

The highest order of minor of A is 3

$$(I.E) \begin{vmatrix} 0 & 1 & 2 \\ 0 & 2 & 4 \\ 8 & 1 & 0 \end{vmatrix} = 0 + 0 + 8(4 - 4) = 0$$

[Expanded along C_1]

$$\text{Also, } \begin{vmatrix} 0 & 2 & 1 \\ 0 & 4 & 3 \\ 8 & 0 & 2 \end{vmatrix} = 0 + 0 - 8 \begin{vmatrix} 2 & 1 \\ 4 & 3 \end{vmatrix}$$

[Expanded along C_1]

$$= -8(6 - 4) = -8(2) = -16 \neq 0$$

$$\therefore \rho(A) = 3$$

2. Find the rank of the following matrices by row reduction method :

$$(i) \begin{bmatrix} 1 & 1 & 1 & 3 \\ 2 & -1 & 3 & 4 \\ 5 & -1 & 7 & 11 \end{bmatrix} \quad [\text{PTA-1; FRT -2022}]$$

$$(ii) \begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 1 & -2 & 3 \\ 1 & -1 & 1 \end{bmatrix} \quad [\text{May - 2022}]$$

$$(iii) \begin{bmatrix} 3 & -8 & 5 & 2 \\ 2 & -5 & 1 & 4 \\ -1 & 2 & 3 & -2 \end{bmatrix}$$

Sol. (i) Let $A = \begin{bmatrix} 1 & 1 & 1 & 3 \\ 2 & -1 & 3 & 4 \\ 5 & -1 & 7 & 11 \end{bmatrix}$

$$A = \begin{bmatrix} 1 & 1 & 1 & 3 \\ 2 & -1 & 3 & 4 \\ 5 & -1 & 7 & 11 \end{bmatrix} \xrightarrow{R_2 \rightarrow R_2 - 2R_1} \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & -3 & 1 & -2 \\ 5 & -1 & 7 & 11 \end{bmatrix}$$

$$\xrightarrow{R_3 \rightarrow R_3 - 5R_1} \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & -3 & 1 & -2 \\ 0 & -6 & 2 & -4 \end{bmatrix}$$

$$\xrightarrow{R_3 \rightarrow R_3 - 2R_2} \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & -3 & 1 & -2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

The last equivalent matrix is in row echelon form
it has two non-zero rows

$$\therefore \rho(A) = 2.$$

$$(ii) \text{ Let } A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 1 & -2 & 3 \\ 1 & -1 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 1 & -2 & 3 \\ 1 & -1 & 1 \end{bmatrix} \xrightarrow{\begin{matrix} R_2 \rightarrow R_2 - 3R_1 \\ R_3 \rightarrow R_3 - R_1 \\ R_4 \rightarrow R_4 - R_1 \end{matrix}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & -4 & 4 \\ 0 & -3 & 2 \end{bmatrix}$$

$$\begin{array}{l}
 R_3 \rightarrow R_3 \div 4 \rightarrow \begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & -1 & 1 \\ 0 & -3 & 2 \end{bmatrix} \xrightarrow{R_4 \rightarrow R_4 - 3R_3} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & -1 & 1 \\ 0 & 0 & -1 \end{bmatrix} \\
 R_3 \rightarrow 7R_3 - R_2 \rightarrow \begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & 0 & 2 \\ 0 & 0 & -1 \end{bmatrix} \xrightarrow{R_4 \rightarrow 2R_4 + R_3} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{bmatrix}
 \end{array}$$

The last equivalent matrix is in row echelon form
it has two non-zero rows

$$\therefore \rho(A) = 3$$

(iii) Let $A = \begin{bmatrix} 3 & -8 & 5 & 2 \\ 2 & -5 & 1 & 4 \\ -1 & 2 & 3 & -2 \end{bmatrix}$

$$\begin{array}{l}
 A \xrightarrow{R_3 \leftrightarrow R_1} \begin{bmatrix} -1 & 2 & 3 & -2 \\ 2 & -5 & 1 & 4 \\ 3 & -8 & 5 & 2 \end{bmatrix} \\
 \xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 + 2R_1 \\ R_3 \rightarrow R_3 + 3R_1 \end{array}} \begin{bmatrix} -1 & 2 & 3 & -2 \\ 0 & -1 & 7 & 0 \\ 0 & -2 & 14 & -4 \end{bmatrix} \\
 \xrightarrow{R_3 \rightarrow R_3 \div 2} \begin{bmatrix} -1 & 2 & 3 & -2 \\ 0 & -1 & 7 & 0 \\ 0 & -1 & 7 & -2 \end{bmatrix} \\
 \xrightarrow{R_3 \rightarrow R_3 - R_2} \begin{bmatrix} -1 & 2 & 3 & -2 \\ 0 & -1 & 7 & 0 \\ 0 & 0 & 0 & -2 \end{bmatrix}
 \end{array}$$

The last equivalent matrix is in row-echelon form. It has three non-zero rows.

$$\therefore \rho(A) = 3$$

3. Find the inverse of each of the following by Gauss – Jordan method :

(i) $\begin{bmatrix} 2 & -1 \\ 5 & -2 \end{bmatrix}$ [Govt. MQP-2019]

(ii) $\begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \\ 6 & -2 & -3 \end{bmatrix}$ (iii) $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 3 \\ 1 & 0 & 8 \end{bmatrix}$

Sol. (i) Let $A = \begin{bmatrix} 2 & -1 \\ 5 & -2 \end{bmatrix}$

Applying Gauss – Jordan method, we get

$$\begin{array}{l}
 [A|I_2] = \left[\begin{array}{cc|cc} 2 & -1 & 1 & 0 \\ 5 & -2 & 0 & 1 \end{array} \right] \\
 \xrightarrow{R_1 \rightarrow R_1 \div 2} \left[\begin{array}{cc|cc} 1 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 5 & -2 & 0 & 1 \end{array} \right]
 \end{array}$$

$$\xrightarrow{R_2 \rightarrow R_2 - 5R_1} \left[\begin{array}{cc|cc} 1 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & \frac{1}{2} & -\frac{5}{2} & 1 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 \times 2} \left[\begin{array}{cc|cc} 1 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 1 & -5 & 2 \end{array} \right]$$

$$\xrightarrow{R_1 \rightarrow R_1 + \frac{1}{2}R_2} \left[\begin{array}{cc|cc} 1 & 0 & -2 & 1 \\ 0 & 1 & -5 & 2 \end{array} \right]$$

$$\therefore \text{We get } A^{-1} = \begin{bmatrix} -2 & 1 \\ -5 & 2 \end{bmatrix}$$

(ii) Let $A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \\ 6 & -2 & -3 \end{bmatrix}$

Applying Gauss – Jordan method, we get

$$[A|I_3] = \left[\begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 1 & 0 & -1 & 0 & 1 & 0 \\ 6 & -2 & -3 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 - R_1} \left[\begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 6 & -2 & -3 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - 6R_1} \left[\begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 4 & -3 & -6 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - 4R_2} \left[\begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -2 & -4 & 1 \end{array} \right]$$

$$\xrightarrow{R_1 \rightarrow R_1 + R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & -1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -2 & -4 & 1 \end{array} \right]$$

$$\xrightarrow{R_1 \rightarrow R_1 + R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & -3 & 1 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -2 & -4 & 1 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 + R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & -3 & 1 \\ 0 & 1 & 0 & -3 & -3 & 1 \\ 0 & 0 & 1 & -2 & -4 & 1 \end{array} \right]$$

$$\text{So, we get } A^{-1} = \begin{bmatrix} -2 & -3 & 1 \\ -3 & -3 & 1 \\ -2 & -4 & 1 \end{bmatrix}$$

(iii) Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 3 \\ 1 & 0 & 8 \end{bmatrix}$

Applying Gauss Jordan method, we get

$$[A|I_3] = \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 5 & 3 & 0 & 1 & 0 \\ 1 & 0 & 8 & 0 & 0 & 1 \end{array} \right]$$

$$\begin{array}{l} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - R_1 \end{array} \rightarrow \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & -3 & -2 & 1 & 0 \\ 0 & -2 & 5 & -1 & 0 & 1 \end{array} \right]$$

$$R_1 \rightarrow R_1 + R_3 \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 8 & 0 & 0 & 1 \\ 0 & 1 & -3 & -2 & 1 & 0 \\ 0 & -2 & 5 & -1 & 0 & 1 \end{array} \right]$$

$$R_3 \rightarrow R_3 + 2R_2 \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 8 & 0 & 0 & 1 \\ 0 & 1 & -3 & -2 & 1 & 0 \\ 0 & 0 & -1 & -5 & 2 & 1 \end{array} \right]$$

$$R_2 \rightarrow R_2 - 3R_3 \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 8 & 0 & 0 & 1 \\ 0 & 1 & 0 & 13 & -5 & -3 \\ 0 & 0 & -1 & -5 & 2 & 1 \end{array} \right]$$

$$R_1 \rightarrow R_1 + 8R_3 \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -40 & 16 & 9 \\ 0 & 1 & 0 & 13 & -5 & -3 \\ 0 & 0 & -1 & -5 & 2 & 1 \end{array} \right]$$

$$R_3 \rightarrow R_3 \times (-1) \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -40 & 16 & 9 \\ 0 & 1 & 0 & 13 & -5 & -3 \\ 0 & 0 & 1 & 5 & -2 & -1 \end{array} \right]$$

So, we get $A^{-1} = \begin{bmatrix} -40 & 16 & 9 \\ 13 & -5 & -3 \\ 5 & -2 & -1 \end{bmatrix}$

EXERCISE 1.3

1. Solve the following system of linear equations by matrix inversion method:

(i) $2x + 5y = -2, x + 2y = -3$ [June-2023]

(ii) $2x - y = 8, 3x + 2y = -2$ [PTA -3]

(iii) $2x + 3y - z = 9, x + y + z = 9, 3x - y - z = -1$

(iv) $x + y + z - 2 = 0, 6x - 4y + 5z - 31 = 0,$
 $5x + 2y + 2z = 13.$

Sol. (i) $2x + 5y = -2, x + 2y = -3$

The matrix form of the system is

$$= \begin{pmatrix} 2 & 5 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -2 \\ -3 \end{pmatrix}$$

$$\Rightarrow AX = B \text{ where}$$

$$A = \begin{pmatrix} 2 & 5 \\ 1 & 2 \end{pmatrix},$$

$$B = \begin{pmatrix} -2 \\ -3 \end{pmatrix},$$

$$X = \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\Rightarrow X = A^{-1}B$$

$$|A| = \begin{vmatrix} 2 & 5 \\ 1 & 2 \end{vmatrix} = 4 - 5 = -1 \neq 0.$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$= \frac{1}{-1} \begin{bmatrix} 2 & -5 \\ -1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} -2 & 5 \\ 1 & -2 \end{bmatrix}$$

$$\therefore X = A^{-1}B = \begin{bmatrix} -2 & 5 \\ 1 & -2 \end{bmatrix} \begin{bmatrix} -2 \\ -3 \end{bmatrix} = \begin{bmatrix} 4 - 15 \\ -2 + 6 \end{bmatrix} = \begin{bmatrix} -11 \\ 4 \end{bmatrix}$$

$$\therefore x = -11, y = 4.$$

(ii) $2x - y = 8, 3x + 2y = -2$

The matrix form of the system is

$$\begin{bmatrix} 2 & -1 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ -2 \end{bmatrix}$$

$$\Rightarrow AX = B \text{ where } A = \begin{bmatrix} 2 & -1 \\ 3 & 2 \end{bmatrix},$$

$$B = \begin{bmatrix} 8 \\ -2 \end{bmatrix}$$

$$\Rightarrow X = A^{-1}B.$$

$$\text{Now, } |A| = \begin{vmatrix} 2 & -1 \\ 3 & 2 \end{vmatrix} = 4 + 3 = 7$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$= \frac{1}{7} \begin{bmatrix} 2 & 1 \\ -3 & 2 \end{bmatrix}$$

$$\therefore X = A^{-1}B = \frac{1}{7} \begin{bmatrix} 2 & 1 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 8 \\ -2 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 16-2 \\ -24-4 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 14 \\ -28 \end{bmatrix} = \begin{bmatrix} \frac{14}{7} \\ \frac{-28}{7} \end{bmatrix} = \begin{bmatrix} 2 \\ -4 \end{bmatrix}$$

$$\therefore x=2, y=-4$$

(iii) $2x+3y-z=9, x+y+z=9, 3x-y-z=-1.$

The matrix form of the system is

$$\begin{bmatrix} 2 & 3 & -1 \\ 1 & 1 & 1 \\ 3 & -1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 9 \\ 9 \\ -1 \end{bmatrix}$$

$$\Rightarrow AX=B \text{ where } A = \begin{bmatrix} 2 & 3 & -1 \\ 1 & 1 & 1 \\ 3 & -1 & -1 \end{bmatrix},$$

$$X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \text{ and } B = \begin{bmatrix} 9 \\ 9 \\ -1 \end{bmatrix}$$

$$\Rightarrow X = A^{-1}B$$

$$|A| = \begin{vmatrix} 2 & 3 & -1 \\ 1 & 1 & 1 \\ 3 & -1 & -1 \end{vmatrix} = 2 \begin{vmatrix} 1 & 1 \\ -1 & -1 \end{vmatrix} - 3 \begin{vmatrix} 1 & 1 \\ 3 & -1 \end{vmatrix} - 1 \begin{vmatrix} 1 & 1 \\ 3 & -1 \end{vmatrix}$$

[Expanded along R_1]

$$= 2(-1+1) - 3(-1-3) - 1(-1-3)$$

$$= 0 - 3(-4) - 1(-4) = 12 + 4 = 16.$$

$$\text{adj } A = \begin{bmatrix} + \begin{vmatrix} 1 & 1 \\ -1 & -1 \end{vmatrix} & - \begin{vmatrix} 1 & 1 \\ 3 & -1 \end{vmatrix} & + \begin{vmatrix} 1 & 1 \\ 3 & -1 \end{vmatrix} \\ - \begin{vmatrix} 3 & -1 \\ -1 & -1 \end{vmatrix} & + \begin{vmatrix} 2 & -1 \\ 3 & -1 \end{vmatrix} & - \begin{vmatrix} 2 & 3 \\ 3 & -1 \end{vmatrix} \\ + \begin{vmatrix} 3 & -1 \\ 1 & 1 \end{vmatrix} & - \begin{vmatrix} 2 & -1 \\ 1 & 1 \end{vmatrix} & + \begin{vmatrix} 2 & 3 \\ 1 & 1 \end{vmatrix} \end{bmatrix}^T$$

$$= \begin{bmatrix} +(-1+1) & -(-1-3) & +(-1-3) \\ -(-3-1) & +(-2+3) & -(-2-9) \\ +(3+1) & -(2+1) & +(2-3) \end{bmatrix}^T$$

$$= \begin{bmatrix} 0 & 4 & -4 \\ 4 & 1 & 11 \\ 4 & -3 & -1 \end{bmatrix}^T = \begin{bmatrix} 0 & 4 & 4 \\ 4 & 1 & -3 \\ -4 & 11 & -1 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{16} \begin{bmatrix} 0 & 4 & 4 \\ 4 & 1 & -3 \\ -4 & 11 & -1 \end{bmatrix}$$

$$\therefore X = A^{-1}B = \frac{1}{16} \begin{bmatrix} 0 & 4 & 4 \\ 4 & 1 & -3 \\ -4 & 11 & -1 \end{bmatrix} \begin{bmatrix} 9 \\ 9 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{16} \begin{bmatrix} 0+36-4 \\ 36+9+3 \\ -36+99+1 \end{bmatrix} = \frac{1}{16} \begin{bmatrix} 32 \\ 48 \\ 64 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}$$

$$\therefore x=2, y=3, z=4$$

(iv) $x+y+z-2=0, 6x-4y+5z-31=0, 5x+2y+2z=13$

The matrix form of the system is

$$\begin{bmatrix} 1 & 1 & 1 \\ 6 & -4 & 5 \\ 5 & 2 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 31 \\ 13 \end{bmatrix}$$

$$AX=B \text{ where } A = \begin{bmatrix} 1 & 1 & 1 \\ 6 & -4 & 5 \\ 5 & 2 & 2 \end{bmatrix},$$

$$X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 2 \\ 31 \\ 13 \end{bmatrix}$$

$$\Rightarrow X = A^{-1}B$$

$$|A| = \begin{vmatrix} 1 & 1 & 1 \\ 6 & -4 & 5 \\ 5 & 2 & 2 \end{vmatrix} = 1 \begin{vmatrix} -4 & 5 \\ 2 & 2 \end{vmatrix} - 1 \begin{vmatrix} 6 & 5 \\ 5 & 2 \end{vmatrix} + 1 \begin{vmatrix} 6 & -4 \\ 5 & 2 \end{vmatrix}$$

$$= 1(-8-10) - 1(12-25) + 1(12+20)$$

$$= 1(-18) - 1(-13) + 1(32) = -18 + 13 + 32 = 27$$

$$\text{adj } A = \begin{bmatrix} + \begin{vmatrix} -4 & 5 \\ 2 & 2 \end{vmatrix} & - \begin{vmatrix} 6 & 5 \\ 5 & 2 \end{vmatrix} & + \begin{vmatrix} 6 & -4 \\ 5 & 2 \end{vmatrix} \\ - \begin{vmatrix} 1 & 1 \\ 2 & 2 \end{vmatrix} & + \begin{vmatrix} 1 & 1 \\ 5 & 2 \end{vmatrix} & - \begin{vmatrix} 1 & 1 \\ 5 & 2 \end{vmatrix} \\ + \begin{vmatrix} 1 & 1 \\ -4 & 5 \end{vmatrix} & - \begin{vmatrix} 1 & 1 \\ 6 & 5 \end{vmatrix} & + \begin{vmatrix} 1 & 1 \\ 6 & -4 \end{vmatrix} \end{bmatrix}^T$$

$$= \begin{bmatrix} +(-8-10) & -(12-25) & +(12+20) \\ -(2-2) & +(2-5) & -(2-5) \\ +(5+4) & -(5-6) & +(-4-6) \end{bmatrix}^T$$

$$= \begin{bmatrix} -18 & 13 & 32 \\ 0 & -3 & 3 \\ 9 & 1 & -10 \end{bmatrix}^T = \begin{bmatrix} -18 & 0 & 9 \\ 13 & -3 & 1 \\ 32 & 3 & -10 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{27} \begin{bmatrix} -18 & 0 & 9 \\ 13 & -3 & 1 \\ 32 & 3 & -10 \end{bmatrix}$$

$$\therefore X = A^{-1} B = \frac{1}{27} \begin{bmatrix} -18 & 0 & 9 \\ 13 & -3 & 1 \\ 32 & 3 & -10 \end{bmatrix} \begin{bmatrix} 2 \\ 31 \\ 13 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{27} \begin{bmatrix} -36+0+117 \\ 26-93+13 \\ 64+93-130 \end{bmatrix} = \frac{1}{27} \begin{bmatrix} 81 \\ -54 \\ 27 \end{bmatrix} = \begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix}$$

$$\therefore x=3, y=-2, z=1$$

2. If $A = \begin{bmatrix} -5 & 1 & 3 \\ 7 & 1 & -5 \\ 1 & -1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$, find

the products AB and BA and hence solve the system of equations $x+y+2z=1$, $3x+2y+z=7$, $2x+y+3z=2$. [Qy. - 2023]

Sol. Given $A = \begin{bmatrix} -5 & 1 & 3 \\ 7 & 1 & -5 \\ 1 & -1 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$

$$AB = \begin{bmatrix} -5 & 1 & 3 \\ 7 & 1 & -5 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} -5+3+6 & -5+2+3 & -10+1+9 \\ 7+3-10 & 7+2-5 & 14+1-15 \\ 1-3+2 & 1-2+1 & 2-1+3 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix} = 4 \cdot I_3$$

$$BA = \begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix} \begin{bmatrix} -5 & 1 & 3 \\ 7 & 1 & -5 \\ 1 & -1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} -5+7+2 & 1+1-2 & 3-5+2 \\ -15+14+1 & 3+2-1 & 9-10+1 \\ -10+7+3 & 2+1-3 & 6-5+3 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix} = 4 \cdot I_3$$

So, we get $AB = BA = 4 \cdot I_3$

$$\Rightarrow \left(\frac{1}{4}A\right) B = B \left(\frac{1}{4}A\right) = I$$

$$\Rightarrow B^{-1} = \frac{1}{4} A$$

Writing the given set of equations in matrix form we get,

$$\begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 7 \\ 2 \end{bmatrix}$$

$$\Rightarrow B \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 7 \\ 2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = B^{-1} \begin{bmatrix} 1 \\ 7 \\ 2 \end{bmatrix} = \left[\frac{1}{4}A\right] \begin{bmatrix} 1 \\ 7 \\ 2 \end{bmatrix}$$

$$= \frac{1}{4} \begin{bmatrix} -5 & 1 & 3 \\ 7 & 1 & -5 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 7 \\ 2 \end{bmatrix}$$

$$= \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{4} \begin{bmatrix} -5+7+6 \\ 7+7-10 \\ 1-7+2 \end{bmatrix}$$

$$= \frac{1}{4} \begin{bmatrix} 8 \\ 4 \\ -4 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix}$$

$$\therefore x=2, y=1, z=-1$$

3. A man is appointed in a job with a monthly salary of certain amount and a fixed amount of annual increment. If his salary was ₹ 19,800 per month at the end of the first month after 3 years of service and ₹ 23,400 per month at the end of the first month after 9 years of service, find his starting salary and his annual increment. (Use matrix inversion method to solve the problem.)

Sol. Let the man's starting salary be ₹ x and his annual increment be ₹ y .

By the given data $x+3y=19800$ and $x+9y=23,400$.

The matrix form of the given system of equations is

$$\begin{bmatrix} 1 & 3 \\ 1 & 9 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 19800 \\ 23400 \end{bmatrix}$$

$$\Rightarrow AX = B \text{ where } A = \begin{bmatrix} 1 & 3 \\ 1 & 9 \end{bmatrix} \text{ and } B = \begin{bmatrix} 19800 \\ 23400 \end{bmatrix}$$

$$\Rightarrow X = A^{-1} B$$

$$|A| = \begin{vmatrix} 1 & 3 \\ 1 & 9 \end{vmatrix}$$

$$= 9 - 3 = 6 \neq 0$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj} A = \frac{1}{6} \begin{bmatrix} 9 & -3 \\ -1 & 1 \end{bmatrix}$$

$$\therefore X = A^{-1} B$$

$$= \frac{1}{6} \begin{bmatrix} 9 & -3 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 19800 \\ 23400 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{6} \begin{bmatrix} 178200 - 70200 \\ -19800 + 23400 \end{bmatrix} = \frac{1}{6} \begin{bmatrix} 108000 \\ 3600 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 18000 \\ 600 \end{bmatrix}$$

∴ $x = 18000, y = 600$.

Hence the man's starting salary is ₹ 18000 and his annual increment is ₹ 600.

- 4. Four men and 4 women can finish a piece of work jointly in 3 days while 2 men and 5 women can finish the same work jointly in 4 days. Find the time taken by one man alone and that of one woman alone to finish the same work by using matrix inversion method.**

Sol. Let the time by one man alone be x days and one woman alone be y days

∴ By the given data,

$$\frac{4}{x} + \frac{4}{y} = \frac{1}{3} \text{ and } \frac{2}{x} + \frac{5}{y} = \frac{1}{4}$$

$$\text{put } \frac{1}{x} = s \text{ and } \frac{1}{y} = t$$

$$\therefore 4s + 4t = \frac{1}{3} \text{ and } 2s + 5t = \frac{1}{4}$$

The matrix form of the system of equation is

$$\begin{bmatrix} 4 & 4 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} s \\ t \end{bmatrix} = \begin{bmatrix} \frac{1}{3} \\ \frac{1}{4} \end{bmatrix} \Rightarrow AX = B \text{ where}$$

$$A = \begin{bmatrix} 4 & 4 \\ 2 & 5 \end{bmatrix} \text{ and } B = \begin{bmatrix} \frac{1}{3} \\ \frac{1}{4} \end{bmatrix}$$

$$X = A^{-1} B$$

$$\text{Now } |A| = \begin{vmatrix} 4 & 4 \\ 2 & 5 \end{vmatrix} = 20 - 8 = 12 \neq 0$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{12} \begin{bmatrix} 5 & -4 \\ -2 & 4 \end{bmatrix}$$

$$\therefore X = A^{-1} B = \frac{1}{12} \begin{bmatrix} 5 & -4 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} \frac{1}{3} \\ \frac{1}{4} \end{bmatrix} = \frac{1}{12} \begin{bmatrix} \frac{5}{3} - 1 \\ -\frac{2}{3} + 1 \end{bmatrix}$$

$$= \frac{1}{12} \begin{bmatrix} \frac{2}{3} \\ \frac{1}{3} \end{bmatrix} = \begin{bmatrix} \frac{2}{3} \times \frac{1}{12} \\ \frac{1}{3} \times \frac{1}{12} \end{bmatrix} = \begin{bmatrix} \frac{1}{18} \\ \frac{1}{36} \end{bmatrix}$$

$$\therefore s = \frac{1}{18}$$

$$\Rightarrow \frac{1}{x} = \frac{1}{18}$$

$$\Rightarrow x = 18$$

$$t = \frac{1}{36}$$

$$\Rightarrow \frac{1}{y} = \frac{1}{36}$$

$$\Rightarrow y = 36.$$

Hence, the time taken by 1 man alone is 18 days and the time taken by 1 woman alone is 36 days.

- 5. The prices of three commodities A, B and C are ₹ x , ₹ y and ₹ z per units respectively. A person P purchases 4 units of B and sells two units of A and 5 units of C. Person Q purchases 2 units of C and sells 3 units of A and one unit of B. Person R purchases one unit of A and sells 3 unit of B and one unit of C. In the process, P, Q and R earn ₹ 15,000, ₹ 1,000 and ₹ 4,000 respectively. Find the prices per unit of A, B and C. (Use matrix inversion method to solve the problem.)**

Sol. Let the prices per unit for the commodities A, B and C be ₹ x , ₹ y and ₹ z .

By the given data,

$$2x - 4y + 5z = 15000$$

$$3x + y - 2z = 1000$$

$$-x + 3y + z = 4000$$

The matrix form of the system of equations is

$$\begin{bmatrix} 2 & -4 & 5 \\ 3 & 1 & -2 \\ -1 & 3 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 15000 \\ 1000 \\ 4000 \end{bmatrix}$$

$$AX = B \text{ where } A = \begin{bmatrix} 2 & -4 & 5 \\ 3 & 1 & -2 \\ -1 & 3 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix},$$

$$\text{and } B = \begin{bmatrix} 15000 \\ 1000 \\ 4000 \end{bmatrix}$$

$$\Rightarrow X = A^{-1} B$$

$$|A| = \begin{vmatrix} 2 & -4 & 5 \\ 3 & 1 & -2 \\ -1 & 3 & 1 \end{vmatrix}$$

$$= 2 \begin{vmatrix} 1 & -2 \\ 3 & 1 \end{vmatrix} + 4 \begin{vmatrix} 3 & -2 \\ -1 & 1 \end{vmatrix} + 5 \begin{vmatrix} 3 & 1 \\ -1 & 3 \end{vmatrix}$$

$$= 2(1+6) + 4(3-2) + 5(9+1)$$

$$= 2(7) + 4(1) + 5(10) = 14 + 4 + 50 = 68.$$

$$\text{adj } A = \begin{bmatrix} + \begin{vmatrix} 1 & -2 \\ 3 & 1 \end{vmatrix} & - \begin{vmatrix} 3 & -2 \\ -1 & 1 \end{vmatrix} & + \begin{vmatrix} 3 & 1 \\ -1 & 3 \end{vmatrix} \\ - \begin{vmatrix} -4 & 5 \\ 3 & 1 \end{vmatrix} & + \begin{vmatrix} 2 & 5 \\ -1 & 1 \end{vmatrix} & - \begin{vmatrix} 2 & -4 \\ -1 & 3 \end{vmatrix} \\ + \begin{vmatrix} -4 & 5 \\ 1 & -2 \end{vmatrix} & - \begin{vmatrix} 2 & 5 \\ 3 & -2 \end{vmatrix} & + \begin{vmatrix} 2 & -4 \\ 3 & 1 \end{vmatrix} \end{bmatrix}^T$$

$$= \begin{bmatrix} +(1+6) & -(3-2) & +(9+1) \\ -(-4-15) & +(2+5) & -(6-4) \\ +(8-5) & -(-4-15) & +(2+12) \end{bmatrix}^T$$

$$= \begin{bmatrix} 7 & -1 & 10 \\ 19 & 7 & -2 \\ 3 & 19 & 14 \end{bmatrix}^T = \begin{bmatrix} 7 & 19 & 3 \\ -1 & 7 & 19 \\ 10 & -2 & 14 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{68} \begin{bmatrix} 7 & 19 & 3 \\ -1 & 7 & 19 \\ 10 & -2 & 14 \end{bmatrix}$$

$$\therefore X = A^{-1} B = \frac{1}{68} \begin{bmatrix} 7 & 19 & 3 \\ -1 & 7 & 19 \\ 10 & -2 & 14 \end{bmatrix} \begin{bmatrix} 15000 \\ 1000 \\ 4000 \end{bmatrix}$$

$$= \frac{1}{68} \begin{bmatrix} 105000 + 19000 + 12000 \\ -15000 + 7000 + 76000 \\ 150000 - 2000 + 56000 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{68} \begin{bmatrix} 136000 \\ 68000 \\ 204000 \end{bmatrix} = \begin{bmatrix} 2000 \\ 1000 \\ 3000 \end{bmatrix}$$

$\therefore x = 2000, y = 1000, z = 3000.$

Hence the prices per unit of the commodities A, B and C are ₹ 2000, ₹ 1000 and ₹ 3000 respectively.

EXERCISE 1.4

1. Solve the following systems of linear equations by Cramer's rule :

(i) $5x - 2y + 16 = 0, x + 3y - 7 = 0$

(ii) $\frac{3}{x} + 2y = 12, \frac{2}{x} + 3y = 13$ [Qy. - 2023]

(iii) $3x + 3y - z = 11, 2x - y + 2z = 9, 4x + 3y + 2z = 25$ [Hy - 2019; July - 2022]

(iv) $\frac{3}{x} - \frac{4}{y} - \frac{2}{z} - 1 = 0, \frac{1}{x} + \frac{2}{y} + \frac{1}{z} - 2 = 0,$

$\frac{2}{x} - \frac{5}{y} - \frac{4}{z} + 1 = 0$ [FRT - 2022; Mar. - 2024]

Sol. (i) Given $\Delta = \begin{vmatrix} 5 & -2 \\ 1 & 3 \end{vmatrix} = 15 + 2 = 17$

$$\Delta_1 = \begin{vmatrix} -16 & -2 \\ 7 & 3 \end{vmatrix}$$

$$= -48 + 14 = -34$$

$$\Delta_2 = \begin{vmatrix} 5 & -16 \\ 1 & 7 \end{vmatrix} = 35 + 16 = 51$$

$$\therefore x = \frac{\Delta_1}{\Delta} = \frac{-34}{17} = -2$$

$$y = \frac{\Delta_2}{\Delta} = \frac{51}{17} = 3$$

$$x = -2, y = 3.$$

(ii) Let $\frac{1}{x} = z$

$$\therefore 3z + 2y = 12, 2z + 3y = 13$$

$$\therefore \Delta = \begin{vmatrix} 3 & 2 \\ 2 & 3 \end{vmatrix} = 9 - 4 = 5$$

$$\Delta_1 = \begin{vmatrix} 12 & 2 \\ 13 & 3 \end{vmatrix} = 36 - 26 = 10$$

$$\Delta_2 = \begin{vmatrix} 3 & 12 \\ 2 & 13 \end{vmatrix} = 39 - 24 = 15$$

$$\therefore z = \frac{\Delta_1}{\Delta} = \frac{10}{5} = 2 \Rightarrow \frac{1}{x} = 2 \Rightarrow x = \frac{1}{2}$$

$$y = \frac{\Delta_2}{\Delta} = \frac{15}{5} = 3$$

$$\therefore x = \frac{1}{2}, y = 3$$

$$\begin{aligned}
 \text{(iii)} \quad \Delta &= \begin{vmatrix} 3 & 3 & -1 \\ 2 & -1 & 2 \\ 4 & 3 & 2 \end{vmatrix} \\
 &= 3 \begin{vmatrix} -1 & 2 \\ 3 & 2 \end{vmatrix} - 3 \begin{vmatrix} 2 & 2 \\ 4 & 2 \end{vmatrix} - 1 \begin{vmatrix} 2 & -1 \\ 4 & 3 \end{vmatrix} \\
 &= 3(-2-6) - 3(4-8) - 1(6+4) \\
 &= 3(-8) - 3(-4) - 1(10) \\
 &= -24 + 12 - 10 = -22 \\
 \Delta_x &= \begin{vmatrix} 11 & 3 & -1 \\ 9 & -1 & 2 \\ 25 & 3 & 2 \end{vmatrix} \\
 &= 11 \begin{vmatrix} -1 & 2 \\ 3 & 2 \end{vmatrix} - 3 \begin{vmatrix} 9 & 2 \\ 25 & 2 \end{vmatrix} - 1 \begin{vmatrix} 9 & -1 \\ 25 & 3 \end{vmatrix} \\
 &= 11(-2-6) - 3(18-50) - 1(27+25) \\
 &= 11(-8) - 3(-32) - 1(52) \\
 &= -88 + 96 - 52 = -44 \\
 \Delta_y &= \begin{vmatrix} 3 & 11 & -1 \\ 2 & 9 & 2 \\ 4 & 25 & 2 \end{vmatrix} \\
 &= 3 \begin{vmatrix} 9 & 2 \\ 25 & 2 \end{vmatrix} - 11 \begin{vmatrix} 2 & 2 \\ 4 & 2 \end{vmatrix} - 1 \begin{vmatrix} 2 & 9 \\ 4 & 25 \end{vmatrix} \\
 &= 3(18-50) - 11(4-8) - 1(50-36) \\
 &= 3(-32) - 11(-4) - 1(14) \\
 &= -96 + 44 - 14 = -66 \\
 \Delta_z &= \begin{vmatrix} 3 & 3 & 11 \\ 2 & -1 & 9 \\ 4 & 3 & 25 \end{vmatrix} \\
 &= 3 \begin{vmatrix} -1 & 9 \\ 3 & 25 \end{vmatrix} - 3 \begin{vmatrix} 2 & 9 \\ 4 & 25 \end{vmatrix} + 11 \begin{vmatrix} 2 & -1 \\ 4 & 3 \end{vmatrix} \\
 &= 3(-25-27) - 3(50-36) + 11(6+4) \\
 &= 3(-52) - 3(14) + 11(10) \\
 &= -156 - 42 + 110 = -88 \\
 \therefore x &= \frac{\Delta_1}{\Delta} = \frac{-44}{-22} = 2 \\
 y &= \frac{\Delta_2}{\Delta} = \frac{-66}{-22} = 3 \\
 z &= \frac{\Delta_3}{\Delta} = \frac{-88}{-22} = 4 \\
 \therefore x &= 2, y = 3, z = 4.
 \end{aligned}$$

$$\text{(iv)} \quad \text{Put } \frac{1}{x} = X, \frac{1}{y} = Y, \frac{1}{z} = Z$$

$$\text{We get } 3X - 4Y - 2Z = 1, X + 2Y + Z = 2, \\ 2X - 5Y - 4Z = -1$$

$$\begin{aligned}
 \therefore \Delta &= \begin{vmatrix} 3 & -4 & -2 \\ 1 & 2 & 1 \\ 2 & -5 & -4 \end{vmatrix} = 3 \begin{vmatrix} 2 & 1 \\ -5 & -4 \end{vmatrix} + 4 \begin{vmatrix} 1 & 1 \\ 2 & -4 \end{vmatrix} - 2 \begin{vmatrix} 1 & 2 \\ 2 & -5 \end{vmatrix} \\
 &= 3(-8+5) + 4(-4-2) - 2(-5-4) \\
 &= 3(-3) + 4(-6) - 2(-9) \\
 &= -9 - 24 + 18 = -15 \\
 \Delta_x &= \begin{vmatrix} 1 & -4 & -2 \\ 2 & 2 & 1 \\ -1 & -5 & -4 \end{vmatrix} \\
 &= 1 \begin{vmatrix} 2 & 1 \\ -5 & -4 \end{vmatrix} + 4 \begin{vmatrix} 2 & 1 \\ -1 & -4 \end{vmatrix} - 2 \begin{vmatrix} 2 & 2 \\ -1 & -5 \end{vmatrix} \\
 &= 1(-8+5) + 4(-8+1) - 2(-10+2) \\
 &= 1(-3) + 4(-7) - 2(-8) \\
 &= -3 - 28 + 16 = -15 \\
 \Delta_y &= \begin{vmatrix} 3 & 1 & -2 \\ 1 & 2 & 1 \\ 2 & -1 & -4 \end{vmatrix} \\
 &= 3 \begin{vmatrix} 2 & 1 \\ -1 & -4 \end{vmatrix} - 1 \begin{vmatrix} 1 & 1 \\ 2 & -4 \end{vmatrix} - 2 \begin{vmatrix} 1 & 2 \\ 2 & -1 \end{vmatrix} \\
 &= 3(-8+1) - 1(-4-2) - 2(-1-4) \\
 &= 3(-7) - 1(-6) - 2(-5) = -21 + 6 + 10 = -5 \\
 \Delta_z &= \begin{vmatrix} 3 & -4 & 1 \\ 1 & 2 & 2 \\ 2 & -5 & -1 \end{vmatrix} = 3 \begin{vmatrix} 2 & 2 \\ -5 & -1 \end{vmatrix} + 4 \begin{vmatrix} 1 & 2 \\ 2 & -1 \end{vmatrix} + 1 \begin{vmatrix} 1 & 2 \\ 2 & -5 \end{vmatrix} \\
 &= 3(-2+10) + 4(-1-4) + 1(-5-4) \\
 &= 3(8) + 4(-5) + 1(-9) \\
 &= 24 - 20 - 9 = -5
 \end{aligned}$$

$$\begin{aligned}
 \therefore X &= \frac{\Delta_x}{\Delta} = \frac{-15}{-15} = 1 \Rightarrow \frac{1}{x} = 1 \\
 \Rightarrow x &= 1 \\
 Y &= \frac{\Delta_y}{\Delta} = \frac{-5}{-15} = \frac{1}{3} \\
 \Rightarrow \frac{1}{y} &= \frac{1}{3} \Rightarrow y = 3 \\
 Z &= \frac{\Delta_z}{\Delta} = \frac{-5}{-15} = \frac{1}{3} \Rightarrow \frac{1}{z} = \frac{1}{3} \Rightarrow z = 3 \\
 \therefore x &= 1, y = 3, z = 3.
 \end{aligned}$$

2. In a competitive examination, one mark is awarded for every correct answer while $\frac{1}{4}$ mark is deducted for every wrong answer. A student answered 100 questions and got 80 marks. How many questions did he answer correctly? (Use Cramer's rule to solve the problem). [Qy - 2019]

Sol. Let x represent the number of question with correct answer and y represent the number of questions with wrong answers.

By the given data, $x + y = 100$ and ... (1)

$$1 \cdot x - \frac{1}{4} y = 80$$

Multiplying by 4 we get,

$$4x - y = 320 \quad \dots (2)$$

From (1) and (2)

$$\Delta = \begin{vmatrix} 1 & 1 \\ 4 & -1 \end{vmatrix} = -1 - 4 = -5$$

$$\Delta_1 = \begin{vmatrix} 100 & 1 \\ 320 & -1 \end{vmatrix} = -100 - 320 = -420$$

$$\Delta_2 = \begin{vmatrix} 1 & 100 \\ 4 & 320 \end{vmatrix} = 320 - 400 = -80$$

$$\therefore x = \frac{\Delta_1}{\Delta} = \frac{-420}{-5} = +84$$

$$\text{and } y = \frac{\Delta_2}{\Delta} = \frac{-80}{-5} = 16$$

Hence, the number of questions with correct answer is 84.

3. A chemist has one solution which is 50% acid and another solution which is 25% acid. How much each should be mixed to make 10 litres of a 40% acid solution ? (Use Cramer's rule to solve the problem). [Hy. - 2023]

Sol. Let the amount of 50% acid be x litres and the amount of 25% acid be y litres

By the given data, $x + y = 10$... (1)

$$\text{and } x \left(\frac{50}{100} \right) + y \left(\frac{25}{100} \right) = 10 \left(\frac{40}{100} \right)$$

$$\Rightarrow 50x + 25y = 400 \Rightarrow 2x + y = 16 \quad \dots (2)$$

$$\Delta = \begin{vmatrix} 1 & 1 \\ 2 & 1 \end{vmatrix} = -1$$

$$\Delta_x = \begin{vmatrix} 10 & 1 \\ 16 & 1 \end{vmatrix} = 10 - 16 = -6$$

$$\Rightarrow \Delta_y = \begin{vmatrix} 1 & 10 \\ 2 & 16 \end{vmatrix} = 16 - 20 = -4$$

$$x = \frac{\Delta_x}{\Delta} = \frac{-6}{-1} = 6$$

$$y = \frac{\Delta_y}{\Delta} = \frac{-4}{-1} = 4$$

i.e., 6 litres of 50% acid and 4 litres of 25% acid solution to be mixed to get 10 litres of 40% of acid solution.

4. A fish tank can be filled in 10 minutes using both pumps A and B simultaneously. However, pump B can pump water in or out at the same rate. If pump B is inadvertently run in reverse, then the tank will be filled in 30 minutes. How long would it take each pump to fill the tank by itself ? (Use Cramer's rule to solve the problem).

Sol. Let the pump A can fill the tank in x minutes, and the pump B can fill the tank in y minutes

In 1 minute A can fill $\frac{1}{x}$ units and in 1 minute B can fill $\frac{1}{y}$ units

$$\therefore \frac{1}{x} + \frac{1}{y} = \frac{1}{10}$$

$$\text{and } \frac{1}{x} - \frac{1}{y} = \frac{1}{30}$$

$$\text{Put } \frac{1}{x} = a \text{ and } \frac{1}{y} = b$$

$$\Rightarrow a + b = \frac{1}{10} \quad \dots (1)$$

$$\text{and } a - b = \frac{1}{30} \quad \dots (2)$$

$$\Delta = \begin{vmatrix} 1 & 1 \\ 1 & -1 \end{vmatrix} = -1 - 1 = -2$$

$$\Delta_1 = \begin{vmatrix} \frac{1}{10} & 1 \\ \frac{1}{30} & -1 \end{vmatrix} = \frac{-1}{10} - \frac{1}{30} = \frac{-3-1}{30} = \frac{-4}{30} = \frac{-2}{15}$$

$$\Delta_2 = \begin{vmatrix} 1 & \frac{1}{10} \\ 1 & \frac{1}{30} \end{vmatrix} = \frac{1}{30} - \frac{1}{10} = \frac{1-3}{30} = \frac{-2}{30} = \frac{-1}{15}$$

$$\therefore a = \frac{\Delta_1}{\Delta} = \frac{-2}{15} = \frac{1}{15} \Rightarrow \frac{1}{x} = \frac{1}{15} \Rightarrow x = 15$$

$$b = \frac{\Delta_2}{\Delta} = \frac{-1}{15} = \frac{1}{30} \Rightarrow \frac{1}{y} = \frac{1}{30} \Rightarrow y = 30$$

Hence the pump A can fill the tank in 15 minutes and the pump B can fill the tank in 30 minutes.

5. A family of 3 people went out for dinner in a restaurant. The cost of two dosai, three idlies and two vadais is ₹ 150. The cost of the two dosai, two idlies and four vadais is ₹ 200. The cost of five dosai, four idlies and two vadais is ₹ 250. The family has ₹ 350 in hand and they ate 3 dosai and six idlies and six vadais. Will they be able to manage to pay the bill within the amount they had ?

Sol. Let the cost of one dosa be ₹ x
 The cost of one idli be ₹ y
 and the cost of one vadai be ₹ z
 By the given data,

$$2x + 3y + 2z = 150$$

$$2x + 2y + 4z = 200$$

$$5x + 4y + 2z = 250$$

$$\therefore \Delta = \begin{vmatrix} 2 & 3 & 2 \\ 2 & 2 & 4 \\ 5 & 4 & 2 \end{vmatrix}$$

$$\begin{aligned} &= 2 \begin{vmatrix} 2 & 4 \\ 4 & 2 \end{vmatrix} - 3 \begin{vmatrix} 2 & 4 \\ 5 & 2 \end{vmatrix} + 2 \begin{vmatrix} 2 & 2 \\ 5 & 4 \end{vmatrix} \\ &= 2(4 - 16) - 3(4 - 20) + 2(8 - 10) \\ &= 2(-12) - 3(-16) + 2(-2) \\ &= -24 + 48 - 4 = 20 \end{aligned}$$

$$\Delta_1 = \begin{vmatrix} 150 & 3 & 2 \\ 200 & 2 & 4 \\ 250 & 4 & 2 \end{vmatrix}$$

Taking 50 common from C_1 and 2 common from C_3 we get,

$$\begin{aligned} &= 100 \begin{vmatrix} 3 & 3 & 1 \\ 4 & 2 & 2 \\ 5 & 4 & 1 \end{vmatrix} = 100 \left[3 \begin{vmatrix} 2 & 2 \\ 4 & 1 \end{vmatrix} - 3 \begin{vmatrix} 4 & 2 \\ 5 & 1 \end{vmatrix} + 1 \begin{vmatrix} 4 & 2 \\ 5 & 4 \end{vmatrix} \right] \\ &= 100[3(2 - 8) - 3(4 - 10) + 1(16 - 10)] \\ &= 100[3(-6) - 3(-6) + 6] \\ &= 100[-18 + 18 + 6] = 600. \end{aligned}$$

$$\Delta_2 = \begin{vmatrix} 2 & 150 & 2 \\ 2 & 200 & 4 \\ 5 & 250 & 2 \end{vmatrix} = 100 \begin{vmatrix} 2 & 3 & 1 \\ 2 & 4 & 2 \\ 5 & 5 & 1 \end{vmatrix}$$

$$\begin{aligned} &= 100 \left[2 \begin{vmatrix} 4 & 2 \\ 5 & 1 \end{vmatrix} - 3 \begin{vmatrix} 2 & 2 \\ 5 & 1 \end{vmatrix} + 1 \begin{vmatrix} 2 & 4 \\ 5 & 5 \end{vmatrix} \right] \\ &= 100[2(4 - 10) - 3(2 - 10) + 1(10 - 20)] \\ &= 100[2(-6) - 3(-8) + 1(-10)] \\ &= 100[-12 + 24 - 10] = 100[2] = 200. \end{aligned}$$

$$\begin{aligned} \Delta_3 &= \begin{vmatrix} 2 & 3 & 150 \\ 2 & 2 & 200 \\ 5 & 4 & 250 \end{vmatrix} = 50 \begin{vmatrix} 2 & 3 & 3 \\ 2 & 2 & 4 \\ 5 & 4 & 5 \end{vmatrix} \\ &= 50 \left[2 \begin{vmatrix} 2 & 4 \\ 4 & 5 \end{vmatrix} - 3 \begin{vmatrix} 2 & 4 \\ 5 & 5 \end{vmatrix} + 3 \begin{vmatrix} 2 & 2 \\ 5 & 4 \end{vmatrix} \right] \\ &= 50[2(10 - 16) - 3(10 - 20) + 3(8 - 10)] \\ &= 50[2(-6) - 3(-10) + 3(-2)] \\ &= 50[-12 + 30 - 6] = 50[12] = 600. \end{aligned}$$

$$\therefore x = \frac{\Delta_1}{\Delta} = \frac{600}{20} = 30$$

$$y = \frac{\Delta_2}{\Delta} = \frac{200}{20} = 10$$

$$z = \frac{\Delta_3}{\Delta} = \frac{600}{20} = 30.$$

Hence, the price of one dosa be ₹ 30, one idli be ₹ 10 and the price of 1 vadai be ₹ 30.

Also the cost of 3 dosai, six idlies and six vadais is
 $= 3x + 6y + 6z = 3(30) + 6(10) + 6(30)$
 $= 90 + 60 + 180 = ₹ 330$

Since the family had ₹ 350 in hand, they will be able to manage to pay the bill.

EXERCISE 1.5

1. Solve the following systems of linear equations by Gaussian elimination method :

(i) $2x - 2y + 3z = 2, x + 2y - z = 3, 3x - y + 2z = 1.$

(ii) $2x + 4y + 6z = 22, 3x + 8y + 5z = 27,$
 $-x + y + 2z = 2$

Sol. (i) Transforming the augmented matrix to echelon form, we get

$$\left[\begin{array}{ccc|c} 2 & -2 & 3 & 2 \\ 1 & 2 & -1 & 3 \\ 3 & -1 & 2 & 1 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2}$$

$$\left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 2 & -2 & 3 & 2 \\ 3 & -1 & 2 & 1 \end{array} \right] \xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 3R_1 \end{array}}$$

$$\left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -6 & 5 & -4 \\ 0 & -7 & 5 & -8 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -6 & 5 & -4 \\ 0 & -1 & 0 & -4 \end{array} \right] \xrightarrow{R_3 \rightarrow 6R_3 - 2R_2}$$

$$\left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -6 & 5 & -4 \\ 0 & 0 & -5 & -20 \end{array} \right]$$

Writing the equivalent equations from the row-echelon matrix, we get,

$$x + 2y - z = 3 \quad \dots (1)$$

$$-6y + 5z = -4 \quad \dots (2)$$

$$-5z = -20 \Rightarrow z = \frac{-20}{-5} = 4.$$

Substituting $z = 4$ in 2 we get,

$$-6y + 5(4) = -4$$

$$\Rightarrow -6y + 20 = -4 \Rightarrow -4 - 20 = -24$$

$$\Rightarrow y = \frac{-24}{-6} = 4.$$

Substituting $y = z = 4$ in (1) we get

$$x + 2(4) - 4 = 3 \Rightarrow x + 8 - 4 = 3$$

$$\Rightarrow x + 4 = 3$$

$$\Rightarrow x = 3 - 4 = -1.$$

$$\therefore x = -1, y = 4, z = 4.$$

$$(ii) \quad 2x + 4y + 6z = 22, 3x + 8y + 5z = 27,$$

$$-x + y + 2z = 2$$

Reducing the augmented matrix to an equivalent row echelon form by using elementary row operations, we get

$$\left[\begin{array}{ccc|c} 2 & 4 & 6 & 22 \\ 3 & 8 & 5 & 27 \\ -1 & 1 & 2 & 2 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_3} \left[\begin{array}{ccc|c} -1 & 1 & 2 & 2 \\ 3 & 8 & 5 & 27 \\ 2 & 4 & 6 & 22 \end{array} \right]$$

$$\begin{array}{l} R_2 \rightarrow R_2 + 3R_1 \\ R_3 \rightarrow R_3 + 2R_1 \end{array} \rightarrow \left[\begin{array}{ccc|c} -1 & 1 & 2 & 2 \\ 0 & 11 & 11 & 33 \\ 0 & 6 & 10 & 26 \end{array} \right]$$

$$\begin{array}{l} R_2 \rightarrow R_2 \div 11 \\ R_3 \rightarrow R_3 \div 2 \end{array} \rightarrow \left[\begin{array}{ccc|c} -1 & 1 & 2 & 2 \\ 0 & 1 & 1 & 3 \\ 0 & 3 & 5 & 13 \end{array} \right]$$

$$R_3 \rightarrow R_3 - 3R_2 \rightarrow \left[\begin{array}{ccc|c} -1 & 1 & 2 & 2 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 2 & 4 \end{array} \right]$$

$$R_3 \rightarrow R_3 \div 2 \rightarrow \left[\begin{array}{ccc|c} -1 & 1 & 2 & 2 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 1 & 2 \end{array} \right]$$

Writing the equivalent equations from the row echelon matrix we get,

$$-x + y + 2z = 2 \quad \dots (1)$$

$$y + z = 3 \quad \dots (2)$$

$$z = 2 \quad \dots (3)$$

Substituting (3) in (2) we get, $y + 2 = 3$

$$\Rightarrow y = 3 - 2 = 1$$

Substituting $y = 1$ and $z = 2$ in (1) we get,

$$-x + 1 + 2(2) = 2 \Rightarrow -x + 1 + 4 = 2$$

$$\Rightarrow -x + 5 = 2 \Rightarrow -x = 2 - 5$$

$$\Rightarrow -x = -3 \Rightarrow x = 3$$

$$\therefore x = 3, y = 1, z = 2.$$

2. If $ax^2 + bx + c$ is divided by $x + 3$, $x - 5$, and $x - 1$, the remainders are 21, 61 and 9 respectively. Find a , b and c . (Use Gaussian elimination method.) [PTA -3]

Sol.

$$\text{Let } P(x) = ax^2 + bx + c$$

$$\text{Given } P(-3) = 21$$

$$[\because P(x) \div x + 3, \text{ the remainder is } 21]$$

$$\Rightarrow a(-3)^2 + b(-3) + c = 21$$

$$\Rightarrow 9a - 3b + c = 21 \quad \dots (1)$$

$$\text{Also, } P(5) = 61$$

$$\Rightarrow a(5)^2 + b(5) + c = 61$$

[using remainder theorem]

$$\Rightarrow 25a + 5b + c = 61 \quad \dots (2)$$

$$\text{and } P(1) = 9$$

$$\Rightarrow a(1)^2 + b(1) + c = 9$$

$$\Rightarrow a + b + c = 9 \quad \dots (3)$$

Reducing the augment matrix to an equivalent row-echelon form using elementary row operations, we get

$$\left[\begin{array}{ccc|c} 9 & -3 & 1 & 21 \\ 25 & 5 & 1 & 61 \\ 1 & 1 & 1 & 9 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_3} \left[\begin{array}{ccc|c} 1 & 1 & 1 & 9 \\ 25 & 5 & 1 & 61 \\ 9 & -3 & 1 & 21 \end{array} \right]$$

$$\begin{array}{l} R_3 \rightarrow R_3 - 9R_1 \\ R_2 \rightarrow R_2 - 25R_1 \end{array} \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 1 & 9 \\ 0 & -20 & -24 & -164 \\ 0 & -12 & -8 & -60 \end{array} \right]$$

$$\begin{array}{l} R_2 \rightarrow R_2 \div 4 \\ R_3 \rightarrow R_3 \div 4 \end{array} \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 1 & 9 \\ 0 & -5 & -6 & -41 \\ 0 & -3 & -2 & -15 \end{array} \right]$$

$$R_3 \rightarrow R_3 - \frac{3}{5}R_2 \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 1 & 9 \\ 0 & -5 & -6 & -41 \\ 0 & 0 & \frac{8}{5} & \frac{48}{5} \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow 5R_3} \left[\begin{array}{ccc|c} 1 & 1 & 1 & 9 \\ 0 & -5 & -6 & -41 \\ 0 & 0 & 8 & 48 \end{array} \right]$$

Writing the equivalent equations from the row-echelon matrix we get,

$$a + b + c = 9 \quad \dots(1)$$

$$-5b - 6c = -41 \quad \dots(2)$$

$$8c = 48$$

$$\Rightarrow c = \frac{48}{8} = 6$$

Substituting $c = 6$ in (2) we get,

$$\Rightarrow -5b - 6(6) = -41$$

$$\Rightarrow -5b = 36 - 41$$

$$\Rightarrow -5b = -41 + 36 = -5$$

$$\Rightarrow b = \frac{-5}{-5} = 1$$

Substituting $b = 1, c = 6$ in (1) we get,

$$a + 1 + 6 = 9$$

$$\Rightarrow a + 7 = 9$$

$$\Rightarrow a = 9 - 7$$

$$\Rightarrow a = 2$$

$$\therefore a = 2, b = 1 \text{ and } c = 6$$

- 3. An amount of ₹ 65,000 is invested in three bonds at the rates of 6%, 8% and 9% per annum respectively. The total annual income is ₹ 4,800. The income from the third bond is ₹ 600 more than that from the second bond. Determine the price of each bond. (Use Gaussian elimination method.)**

Sol. Let the price of bond invested in 6%, 8% and 9% rates be ₹ x , ₹ y and ₹ z respectively

$$\therefore \text{By the given data, } x + y + z = 65,000 \quad \dots(1)$$

$$\frac{6 \times x \times 1}{100} + \frac{8 \times y \times 1}{100} + \frac{9 \times z \times 1}{100} = 4,800$$

$$[\because \text{Interest} = \frac{\text{PNR}}{100}]$$

$$\Rightarrow \frac{6x}{100} + \frac{8y}{100} + \frac{9z}{100} = 4,800$$

$$\Rightarrow 6x + 8y + 9z = 4,80,000 \quad \dots(2)$$

$$\text{Also, } \frac{9z}{100} = 600 + \frac{8y}{100}$$

$$\Rightarrow \frac{-8y}{100} + \frac{9z}{100} = 600$$

$$\Rightarrow -8y + 9z = 60,000 \quad \dots(3)$$

Reducing the augmented matrix to an equivalent row-echelon form by using elementary row operation, we get

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 65,000 \\ 6 & 8 & 9 & 4,80,000 \\ 0 & -8 & 9 & 60,000 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 - 6R_1} \left[\begin{array}{ccc|c} 1 & 1 & 1 & 65,000 \\ 0 & 2 & 3 & 90,000 \\ 0 & -8 & 9 & 60,000 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 + 4R_2} \left[\begin{array}{ccc|c} 1 & 1 & 1 & 65,000 \\ 0 & 2 & 3 & 90,000 \\ 0 & 0 & 21 & 4,20,000 \end{array} \right]$$

Writing the equivalent from the row echelon matrix we get,

$$x + y + z = 65,000 \quad \dots(1)$$

$$2y + 3z = 90,000 \quad \dots(2)$$

$$21z = 4,20,000$$

$$\Rightarrow z = \frac{4,20,000}{21} = 20,000$$

Substituting $z = 20,000$ in (2),

$$2y + 3(20,000) = 90,000$$

$$\Rightarrow 2y + 60,000 = 90,000$$

$$\Rightarrow 2y = 90,000 - 60,000$$

$$= 30,000$$

$$\Rightarrow y = \frac{30,000}{2} = 15,000$$

Substituting $y = 15,000$ and $z = 20,000$ in (1) we get,

$$x + 15,000 + 20,000 = 65,000$$

$$\Rightarrow x + 35,000 = 65,000$$

$$\Rightarrow x = 65,000 - 35,000$$

$$\Rightarrow x = 30,000$$

Thus the price of 6% bond is ₹ 30,000 the price of 8% bond is ₹ 15,000 and the price of 9% bond is ₹ 20,000.

- 4. A boy is walking along the path $y = ax^2 + bx + c$ through the points $(-6, 8)$, $(-2, 12)$ and $(3, 8)$. He wants to meet his friend at $P(7, 60)$. Will he meet his friend? (Use Gaussian elimination method.)** [Mar. - 2023]

Sol. Given $y = ax^2 + bx + c \quad \dots(1)$

$(-6, 8)$ lies on (1)

$$\Rightarrow 8 = a(-6)^2 + b(-6) + c$$

$$\Rightarrow 8 = 36a - 6b + c \quad \dots(2)$$

$(-2, -12)$ lies on (1)

$$\Rightarrow -12 = a(-2)^2 + b(-2) + c$$

$$\Rightarrow -12 = 4a - 2b + c \quad \dots(3)$$

Also $(3, 8)$ lies on (1)

$$\Rightarrow 8 = a(3)^2 + b(3) + c$$

$$\Rightarrow 8 = 9a + 3b + c \quad \dots(4)$$

Reducing the augment matrix to an equivalent row-echelon form by using elementary row operations, we get,

$$\left[\begin{array}{ccc|c} 36 & -6 & 1 & 8 \\ 4 & -2 & 1 & -12 \\ 9 & 3 & 1 & 8 \end{array} \right] \xrightarrow{\substack{R_2 \rightarrow 9R_2 - R_1 \\ R_3 \rightarrow 4R_3 - R_1}} \left[\begin{array}{ccc|c} 36 & -6 & 1 & 8 \\ 0 & -12 & 8 & -116 \\ 0 & 18 & 3 & 24 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 \div 4} \left[\begin{array}{ccc|c} 36 & -6 & 1 & 8 \\ 0 & -3 & 2 & -29 \\ 0 & 6 & 1 & 8 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 + 2R_2} \left[\begin{array}{ccc|c} 36 & -6 & 1 & 8 \\ 0 & -3 & 2 & -29 \\ 0 & 0 & 5 & -50 \end{array} \right]$$

Writing the equivalent equation from the row echelon matrix, we get $36a - 6b + c = 8 \quad \dots(1)$

$$-3b + 2c = -29 \quad \dots(2)$$

$$5c = -50$$

$$\Rightarrow c = \frac{-50}{5} = -10$$

Substituting $c = -10$ in (2) we get,

$$-3b + 2(-10) = -29$$

$$\Rightarrow -3b - 20 = -29$$

$$\Rightarrow -3b = -29 + 20$$

$$\Rightarrow -3b = -9$$

$$\Rightarrow b = \frac{-9}{-3} = 3$$

Substituting $b = 3$ and $c = -10$ in (1) we get,

$$36a - 6(3) - 10 = 8$$

$$\Rightarrow 36a - 18 - 10 = 8$$

$$\Rightarrow 36a - 28 = 8$$

$$\Rightarrow 36a = 8 + 28 = 36$$

$$\Rightarrow a = \frac{36}{36} = 1$$

$$\therefore a = 1, b = 3, c = -10$$

Hence the path of the boy is

$$y = 1(x^2) + 3(x) - 10$$

$$\Rightarrow y = x^2 + 3x - 10$$

Since his friend is at $P(7, 60)$,

$$60 = (7)^2 + 3(7) - 10$$

$$\Rightarrow 60 = 49 + 21 - 10$$

$$\Rightarrow 60 = 70 - 10 = 60$$

$$\Rightarrow 60 = 60$$

Since $(7, 60)$ satisfies his path, he can meet his friend who is at $P(7, 60)$

EXERCISE 1.6

1. Test for consistency and if possible, solve the following systems of equations by rank method.

(i) $x - y + 2z = 2, 2x + y + 4z = 7,$
 $4x - y + z = 4$

(ii) $3x + y + z = 2, x - 3y + 2z = 1,$
 $7x - y + 4z = 5.$

(iii) $2x + 2y + z = 5, x - y + z = 1,$
 $3x + y + 2z = 4$ [Sep. - 2020; Hy. - 2023]

(iv) $2x - y + z = 2, 6x - 3y + 3z = 6,$
 $4x - 2y + 2z = 4.$ [PTA - 5; Hy - 2019]

Sol. (i) $x - y + 2z = 2, 2x + y + 4z = 7, 4x - y + z = 4$

The matrix form of the system is $AX = B$

where $A = \begin{bmatrix} 1 & -1 & 2 \\ 2 & 1 & 4 \\ 4 & -1 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ and $B = \begin{bmatrix} 2 \\ 7 \\ 4 \end{bmatrix}$

Applying elementary row operations on the augment matrix $[A|B]$ we get,

$$[A|B] = \left[\begin{array}{ccc|c} 1 & -1 & 2 & 2 \\ 2 & 1 & 4 & 7 \\ 4 & -1 & 1 & 4 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 - 2R_1, R_3 \rightarrow R_3 - 4R_1} \left[\begin{array}{ccc|c} 1 & -1 & 2 & 2 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & -7 & -4 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{ccc|c} 1 & -1 & 2 & 2 \\ 0 & 3 & 0 & 3 \\ 0 & 0 & -7 & -7 \end{array} \right]$$

Here $\rho(A) = 3$ and $\rho[A|B] = 3$

$\therefore \rho(A) = \rho[A|B] = 3 = \text{number of unknowns}$

Hence the system is consistent with unique solution.

Writing the equivalent equations from the row-echelon matrix, we get

$$x - y + 2z = 2 \quad \dots(1)$$

$$3y = 3 \Rightarrow y = 1 \quad \dots(2)$$

$$-7z = -7 \Rightarrow z = \frac{-7}{-7} = 1 \quad \dots(3)$$

Solutions $y = 1$ and $z = 1$ in (1) we get,

$$x - 1 + 2(1) = 2$$

$$\Rightarrow x - 1 + 2 = 2$$

$$\Rightarrow x + 1 = 2$$

$$\Rightarrow x = 2 - 1 = 1$$

$$\Rightarrow x = 1$$

$$\therefore x = 1, y = 1, z = 1$$

(ii) $3x + y + z = 2, x - 3y + 2z = 1,$
 $7x - y + 4z = 5.$

Then matrix form of the system is $AX = B$

$$\text{where } A = \begin{bmatrix} 3 & 1 & 1 \\ 1 & -3 & 2 \\ 7 & -1 & 4 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 2 \\ 1 \\ 5 \end{bmatrix}$$

Applying elementary row operations on the augment matrix $[A|B]$ we get,

$$[A|B] = \left[\begin{array}{ccc|c} 3 & 1 & 1 & 2 \\ 1 & -3 & 2 & 1 \\ 7 & -1 & 4 & 5 \end{array} \right] \xrightarrow{R_2 \leftrightarrow R_1} \left[\begin{array}{ccc|c} 1 & -3 & 2 & 1 \\ 3 & 1 & 1 & 2 \\ 7 & -1 & 4 & 5 \end{array} \right]$$

$$\xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - 3R_1 \\ R_3 \rightarrow R_3 - 7R_1 \end{array}} \left[\begin{array}{ccc|c} 1 & -3 & 2 & 1 \\ 0 & 10 & -5 & -1 \\ 0 & 20 & -10 & -2 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - 2R_2} \left[\begin{array}{ccc|c} 1 & -3 & 2 & 1 \\ 0 & 10 & -5 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Here $\rho(A) = 2$, and $\rho[A|B] = 2$ [since there only two non-zero rows]. So, $\rho(A) = \rho[A|B] = 2 < 3$, the given system is consistent with one parameter family of solutions. So, put $z = t$, $x \in \mathbb{R}$. Writing the equivalent equations from the row echelon matrix we get,

$$x - 3y + 2z = 1 \quad \dots(1)$$

$$10y - 5z = -1 \quad \dots(2)$$

$$z = t \quad \dots(3)$$

$$(2) \text{ becomes } 10y - 5t = -1$$

$$\Rightarrow 10y = 5t - 1$$

$$\Rightarrow y = \frac{1}{10} [5t - 1]$$

$$\text{Also, from (1), } x - \frac{3}{10} [5t - 1] + 2t = 1$$

$$\Rightarrow x = \frac{3}{10} [5t - 1] - 2t + 1 = \frac{15t - 3 - 20t + 10}{10}$$

$$\Rightarrow x = \frac{1}{10} [-5t + 7]$$

Hence the solution set is

$$x = \frac{1}{10} (7 - 5t), y = \frac{1}{10} (5t - 1) \text{ and } z = t \text{ where}$$

$$t \in \mathbb{R}.$$

(iii) $2x + 2y + z = 5, x - y + z = 1, 3x + y + 2z = 4$

The matrix form of the given system is $AX = B$ where

$$A = \begin{bmatrix} 2 & 2 & 1 \\ 1 & -1 & 1 \\ 3 & 1 & 2 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 5 \\ 1 \\ 4 \end{bmatrix}$$

Applying elementary row operations on the augmented matrix $[A|B]$ we get,

$$[A|B] = \left[\begin{array}{ccc|c} 2 & 2 & 1 & 5 \\ 1 & -1 & 1 & 1 \\ 3 & 1 & 2 & 4 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccc|c} 1 & -1 & 1 & 1 \\ 2 & 2 & 1 & 5 \\ 3 & 1 & 2 & 4 \end{array} \right]$$

$$\xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 3R_1 \end{array}} \left[\begin{array}{ccc|c} 1 & -1 & 1 & 1 \\ 0 & 4 & -1 & 3 \\ 0 & 4 & -1 & 1 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{ccc|c} 1 & -1 & 1 & 1 \\ 0 & 4 & -1 & 3 \\ 0 & 0 & 0 & -2 \end{array} \right]$$

Here $\rho(A) = 2$ [$\because$ There are 2 non-Zero rows]

and $\rho[A|B] = 3$ [$\because$ There are 3 non-zero rows]

Here, $\rho(A) \neq \rho[A|B]$

Hence, the given system is inconsistent and has no solution.

(iv) $2x - y + z = 2, 6x - 3y + 3z = 6,$
 $4x - 2y + 2z = 4.$

The matrix form of the given system is $AX = B$ where

$$A = \begin{bmatrix} 2 & -1 & 1 \\ 6 & -3 & 3 \\ 4 & -2 & 2 \end{bmatrix} \quad X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \quad B = \begin{bmatrix} 2 \\ 6 \\ 4 \end{bmatrix}$$

Applying elementary row operations on the augment matrix $[A|B]$, we get,

$$[A|B] = \left[\begin{array}{ccc|c} 2 & -1 & 1 & 2 \\ 6 & -3 & 3 & 6 \\ 4 & -2 & 2 & 4 \end{array} \right] \xrightarrow{\substack{R_2 \rightarrow R_2 - 3R_1 \\ R_3 \rightarrow R_3 - 2R_1}} \left[\begin{array}{ccc|c} 2 & -1 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Here $\rho(A) = 1$ [$\because$ only one non zero row]

and $\rho[A|B] = 1$ [$\because$ only one-zero row]

$\therefore \rho(A) = \rho[A|B] = 1 < 3$, the given system is consistent and has two parameter family of solutions.

So, $z = t$ and $y = s$ where $s, t \in \mathbb{R}$.

Writing the equivalent equations from the row-echelon matrix, we get

$$2x - y + z = 2 \quad \dots(1)$$

$$y = s \quad \dots(2)$$

$$z = t \quad \dots(3)$$

Substituting (2) and (3) in (1) we get,

$$2x - s + t = 2$$

$$\Rightarrow 2x = s - t + 2$$

$$\Rightarrow x = \frac{1}{2} [s - t + 2]$$

$\therefore$ Solution set is $x = \frac{1}{2} (s - t + 2)$, $y = s$, $z = t$ where $s, t \in \mathbb{R}$

2. Find the value of k for which the equations

$$kx - 2y + z = 1, x - 2ky + z = -2,$$

$$x - 2y + kz = 1 \text{ have} \quad [\text{Qy. - 2019 \& 2023}]$$

(i) no solution

(ii) unique solution

(iii) infinitely many solution

Sol. $kx - 2y + z = 1, x - 2ky + z = -2, x - 2y + kz = 1$

The matrix form of the system is $AX = B$ where

$$A = \begin{bmatrix} k & -2 & 1 \\ 1 & -2k & 1 \\ 1 & -2 & k \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$$

Applying elementary row operation on the augment matrix $[A|B]$ we get,

$$[A|B] = \left[\begin{array}{ccc|c} k & -2 & 1 & 1 \\ 1 & -2k & 1 & -2 \\ 1 & -2 & k & 1 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_3} \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 1 & -2k & 1 & -2 \\ k & -2 & 1 & 1 \end{array} \right]$$

$$\xrightarrow{\substack{R_2 \rightarrow R_2 - R_1 \\ R_3 \rightarrow R_3 - kR_1}} \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 0 & -2k + 2 & 1 - k & -3 \\ 0 & -2 + 2k & 1 - k^2 & 1 - k \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 + R_2} \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 0 & -2k + 2 & 1 - k & -3 \\ 0 & 0 & 1 - k^2 + 1 - k & -k - 2 \end{array} \right]$$

$$\rightarrow \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 0 & -2k + 2 & 1 - k & -3 \\ 0 & 0 & -k^2 - k + 2 & -k - 2 \end{array} \right]$$

$$\rightarrow \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 0 & -2k + 2 & 1 - k & -3 \\ 0 & 0 & (k + 2)(1 - k) & -k - 2 \end{array} \right] \quad \dots(1)$$

Case (i) when $k = 1$

$$[A|B] \rightarrow \left[\begin{array}{ccc|c} 1 & -2 & 1 & 1 \\ 0 & 0 & 0 & -3 \\ 0 & 0 & 0 & -3 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 1 \\ 0 & 0 & 0 & -3 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Here $\rho(A) = 1$ and $\rho[A|B] = 2$

So, $\rho(A) \neq \rho[A|B] \Rightarrow$ The system has no solution.

Case (ii) When $k \neq 1, k \neq -2$

$$[A|B] \rightarrow \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 0 & -2k + 2 & 1 - k & -3 \\ 0 & 0 & \text{not zero} & \text{not zero} \end{array} \right]$$

$\Rightarrow \rho(A) = 3$ and $\rho[A|B] = 3$

so, $\rho(A) = \rho[A|B] = 3 =$ the number of unknowns

Hence, the system has unique solution.

Case (iii) when $k = -2$

$$\rho[A|B] \rightarrow \left[\begin{array}{ccc|c} 1 & -2 & -2 & 1 \\ 1 & 6 & 3 & -3 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Here $\rho(A) = 2$ and $\rho[A|B] = 2$

$\therefore \rho(A) = \rho[A|B] = 2 < 3$, the number of unknowns

so the system is consistent with infinitely many solutions.

3. Investigate the values of λ and μ the system of linear equations $2x + 3y + 5z = 9, 7x + 3y - 5z = 8, 2x + 3y + \lambda z = \mu$, have

(i) no solution

(ii) a unique solution

(iii) an infinite number of solutions.

Sol. $2x + 3y + 5z = 9, 7x + 3y - 5z = 8, 2x + 3y + \lambda z = \mu$

The matrix form of the system is $AX = B$ where

$$A = \begin{bmatrix} 2 & 3 & 5 \\ 7 & 3 & -5 \\ 2 & 3 & \lambda \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 9 \\ 8 \\ \mu \end{bmatrix}$$

Applying elementary row operations on the augmented matrix $[A|B]$ we get,

$$[A|B] = \left[\begin{array}{ccc|c} 2 & 3 & 5 & 9 \\ 7 & 3 & -5 & 8 \\ 2 & 3 & \lambda & \mu \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccc|c} 7 & 3 & -5 & 8 \\ 2 & 3 & 5 & 9 \\ 2 & 3 & \lambda & \mu \end{array} \right]$$

$$\xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - \frac{2}{7}R_1 \\ R_3 \rightarrow R_3 - R_2 \end{array}} \left[\begin{array}{ccc|c} 7 & 3 & -5 & -8 \\ 0 & \frac{15}{7} & \frac{45}{7} & \frac{45}{7} \\ 0 & 0 & \lambda - 5 & \mu - 9 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 \times 7} \left[\begin{array}{ccc|c} 7 & 3 & -5 & -8 \\ 0 & 15 & 45 & 47 \\ 0 & 0 & \lambda - 5 & \mu - 9 \end{array} \right]$$

Case (i) When $\lambda = 5$

$$[A|B] = \left[\begin{array}{ccc|c} 7 & 3 & -5 & -8 \\ 0 & 15 & 45 & 47 \\ 0 & 0 & 0 & \mu - 9 \end{array} \right]$$

Here $\rho(A) = 2$ and $\rho[A|B] = 3$

So, $\rho(A) \neq \rho[A|B]$

Hence the system is inconsistent and has no solution

Case (ii) When $\lambda \neq 5$, $\mu \neq 9$

$$[A|B] = \left[\begin{array}{ccc|c} 7 & 3 & -5 & -8 \\ 0 & 15 & 45 & 47 \\ 0 & 0 & \text{not zero} & \text{not zero} \end{array} \right]$$

Here $\rho(A) = 3$ and $\rho[A|B] = 3$

$\therefore \rho(A) = \rho[A|B] = 3 = \text{number of unknowns}$

Hence, the system is consistent with unique solution

Case (iii) When $\lambda = 5$ and $\mu = 9$

$$[A|B] = \left[\begin{array}{ccc|c} 7 & 3 & -5 & -8 \\ 0 & 15 & 45 & 47 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Here $\rho(A) = 2$, $\rho[A|B] = 2$

$\therefore \rho(A) = \rho[A|B] = 2 < \text{number of unknowns}$

$\therefore$ The system is consistent and has infinite number of solutions.

EXERCISE 1.7

1. Solve the following system of homogeneous equations.

(i) $3x + 2y + 7z = 0, 4x - 3y - 2z = 0,$
 $5x + 9y + 23z = 0$

(ii) $2x + 3y - z = 0, x - y - 2z = 0, 3x + y + 3z = 0.$

Sol. (i) $3x + 2y + 7z = 0, 4x - 3y - 2z = 0,$
 $5x + 9y + 23z = 0$

Here the number of equations is equal to the number of unknowns.

Transforming into echelon form, the augmented matrix becomes

$$\left[\begin{array}{ccc|c} 3 & 2 & 7 & 0 \\ 4 & -3 & -2 & 0 \\ 5 & 9 & 23 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 3 & 2 & 7 & 0 \\ 0 & -17 & -34 & 0 \\ 0 & 17 & 34 & 0 \end{array} \right] \begin{array}{l} R_2 = 3R_2 - 4R_1 \\ R_3 = 3R_3 - 5R_1 \end{array}$$

$$\left[\begin{array}{ccc|c} 3 & 2 & 7 & 0 \\ 0 & -17 & -34 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \begin{array}{l} \\ R_3 = R_3 + R_2 \\ \end{array}$$

So, $\rho(AB) = \rho(A) = 2$ and $n = 3$.

Hence, the system has a one parameter family of solutions. Writing the equations using the echelon form, we get

$$3x + 2y + 7z = 0 \text{ and } -17y - 34z = 0$$

To solve the equations let $z = t$,

$$\text{then } -17y - 34t = 0$$

$$-17y = 34t$$

$$y = \frac{34t}{17}$$

$$y = -2t$$

Substituting $z = t$ and $y = -2t$ in

$$3x + 2y + 7z = 0$$

$$\text{We get } 3x + 2(-2t) + 7t = 0$$

$$3x - 4t + 7t = 0$$

$$3x + 3t = 0$$

$$3x = -3t$$

$$x = -\frac{3t}{3}$$

$$x = -t,$$

So the solution is

$$x = -t, y = -2t \text{ and } z = t \text{ where } t \in \mathbb{R}$$

(ii) $2x + 3y - z = 0, x - y - 2z = 0, 3x + y + 3z = 0$.

Here the number of equations is equal to the number of unknowns. Transforming into echelon form, the augmented matrix becomes

$$\begin{bmatrix} 2 & 3 & -1 & 0 \\ 1 & -1 & -2 & 0 \\ 3 & 1 & 3 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -1 & -2 & 0 \\ 2 & 3 & -1 & 0 \\ 3 & 1 & 3 & 0 \end{bmatrix} R_1 \leftrightarrow R_2$$

$$\begin{bmatrix} 1 & -1 & -2 & 0 \\ 0 & 5 & 3 & 0 \\ 0 & 4 & 9 & 0 \end{bmatrix} \begin{matrix} R_2 = R_2 - 2R_1 \\ R_3 = R_3 - 3R_1 \end{matrix}$$

$$\rightarrow \begin{bmatrix} 1 & -1 & -2 & 0 \\ 0 & 5 & 3 & 0 \\ 0 & 0 & 33 & 0 \end{bmatrix} R_3 = 5R_2 - 4R_2$$

So, $\rho(AB) = \rho(A) = 3$ and $n = 3$

Hence, the system has a unique solution.

Since $x = 0, y = 0, z = 0$, is always a solution of the homogeneous system, the only solution is the trivial solution

$$x = 0, y = 0, z = 0$$

2. Determine the values of λ for which the following system of equations

$$x + y + 3z = 0, 4x + 3y + \lambda z = 0, 2x + y + 2z = 0 \text{ has}$$

(i) a unique solution

(ii) a non-trivial solution.

Sol. $x + y + 3z = 0, 4x + 3y + \lambda z = 0, 2x + y + 2z = 0$

Reducing the augmented matrix to row – echelon form we get,

$$[A|0] = \begin{bmatrix} 1 & 1 & 3 & 0 \\ 4 & 3 & \lambda & 0 \\ 2 & 1 & 2 & 0 \end{bmatrix} \xrightarrow{R_1 \leftrightarrow R_3} \begin{bmatrix} 1 & 1 & 3 & 0 \\ 2 & 1 & 2 & 0 \\ 4 & 3 & \lambda & 0 \end{bmatrix}$$

$$\xrightarrow{\begin{matrix} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 4R_1 \end{matrix}} \begin{bmatrix} 1 & 1 & 3 & 0 \\ 0 & -1 & -4 & 0 \\ 0 & -1 & \lambda - 12 & 0 \end{bmatrix}$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \begin{bmatrix} 1 & 1 & 3 & 0 \\ 0 & -1 & -4 & 0 \\ 0 & 0 & \lambda - 8 & 0 \end{bmatrix}$$

Case (i) when $\lambda \neq 8$

$$[A|0] = \begin{bmatrix} 1 & 1 & 3 & 0 \\ 0 & -1 & -4 & 0 \\ 0 & 0 & \text{not zero} & 0 \end{bmatrix}$$

Here $\rho(A) = 3, \rho([A|0]) = 3$

$\therefore \rho(A) = \rho([A|0]) = 3 =$ the number of unknowns
 $\therefore$ The given system is consistent and has unique solution.

Case (ii) when $\lambda = 8$

$$[A|0] = \begin{bmatrix} 1 & 1 & 3 & 0 \\ 0 & -1 & -4 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Here $\rho(A) = 2, \rho([A|0]) = 2$

$\therefore \rho(A) = \rho([A|0]) = 2 < 3$, the number of unknowns,


$\therefore$ The system is consistent and has non-trivial solutions.

3. By using Gaussian elimination method, balance the chemical reaction equation:



Sol. Given $C_2H_6 + O_2 \rightarrow H_2O + CO_2$

We have to find positive integers x_1, x_2, x_3 and x_4 such that



The number of carbon atoms on the LHS of (1) should be equal to the number of carbon atoms on the RHS of (1).

$$\therefore 2x_1 = 1x_4$$

$$\Rightarrow 2x_1 - x_4 = 0 \quad \dots(2)$$

Considering hydrogen atoms we get,

$$6x_1 = 2x_3 \Rightarrow 6x_1 - 2x_3 = 0$$

$$\Rightarrow 3x_1 - x_3 = 0 \quad \dots(3)$$

Also, considering oxygen atoms we get,

$$2x_2 = 1x_3 + 2x_4$$

$$\Rightarrow 2x_2 - x_3 - 2x_4 = 0 \quad \dots(4)$$

Equations (2), (3) and (4) form a homogeneous system of linear equations in 4 unknowns

$\therefore$ The augmented matrix $[A|0]$ is

$$\begin{bmatrix} 2 & 0 & 0 & -1 & 0 \\ 3 & 0 & -1 & 0 & 0 \\ 0 & 2 & -1 & -2 & 0 \end{bmatrix}$$

By Gaussian elimination method, we get,

$$\xrightarrow{R_1 \rightarrow R_1 \div 2} \begin{bmatrix} 1 & 0 & 0 & -\frac{1}{2} & 0 \\ 3 & 0 & -1 & 0 & 0 \\ 0 & 2 & -1 & -2 & 0 \end{bmatrix}$$

$$\xrightarrow{R_2 \rightarrow R_2 - 3R_1} \left[\begin{array}{cccc|c} 1 & 0 & 0 & -\frac{1}{2} & 0 \\ 0 & 0 & -1 & \frac{3}{2} & 0 \\ 0 & 2 & -1 & -2 & 0 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{cccc|c} 1 & 0 & 0 & -\frac{1}{2} & 0 \\ 0 & 0 & -1 & \frac{3}{2} & 0 \\ 0 & 2 & 0 & -\frac{7}{2} & 0 \end{array} \right]$$

$$\xrightarrow{R_2 \leftrightarrow R_3} \left[\begin{array}{cccc|c} 1 & 0 & 0 & -\frac{1}{2} & 0 \\ 0 & 2 & 0 & -\frac{7}{2} & 0 \\ 0 & 0 & -1 & \frac{3}{2} & 0 \end{array} \right]$$

$$\begin{array}{l} R_1 \rightarrow 2R_1 \\ R_2 \rightarrow 2R_2 \\ R_2 \rightarrow 2R_3 \end{array} \rightarrow \left[\begin{array}{cccc|c} 2 & 0 & 0 & -1 & 0 \\ 0 & 4 & 0 & -7 & 0 \\ 0 & 0 & -2 & 3 & 0 \end{array} \right]$$

Here $\rho(A) = \rho([A|B]) = 3 < 4$, the number of unknowns

∴ The system is consistent with one parameter family of solutions, so let $x_4 = t$

Writing the equations, from the row-echelon form we get (from equation (2))

$$\begin{aligned} 2x_1 - x_4 &= 0 \\ \Rightarrow 2x_1 &= x_4 \\ \Rightarrow 2x_1 &= t \Rightarrow x_1 = \frac{t}{2} \end{aligned}$$

From equation (3)

$$\begin{aligned} \Rightarrow 3x_1 - x_3 &= 0 \\ \Rightarrow x_3 &= 3x_1 \Rightarrow x_3 = \frac{3t}{2} \end{aligned}$$

From equation (4)

$$\begin{aligned} 2x_2 - x_3 - 2x_4 &= 0 \Rightarrow 2x_2 - \frac{3t}{2} - 2t = 0 \\ \Rightarrow 2x_2 - \frac{7t}{4} &= 0 \Rightarrow x_2 = \frac{7t}{4} \end{aligned}$$


Since x_1, x_2, x_3 and x_4 are positive integers, let us choose $t = 4$

$$\therefore x_1 = \frac{4}{2} = 2$$

$$\therefore x_2 = \frac{7(4)}{4} = 7$$

$$\therefore x_3 = \frac{3(4)}{2} = 6 \text{ and } x_4 = t = 4$$

So, the balanced equation is



EXERCISE 1.8

Choose the Correct or the most suitable answer from the given four alternatives :

1. If $|\text{adj}(\text{adj} A)| = |A|^9$, then the order of the square matrix A is [FRT -2022]

- (1) 3 (2) 4 (3) 2 (4) 5

[Ans. (2) 4]

Hint : $|\text{adj}(\text{adj} A)| = |A|^{(n-1)^2}$

$$\begin{aligned} \therefore (n-1)^2 &= 9 \Rightarrow (n-1) = 3 \\ \Rightarrow n-1 &= 3 \Rightarrow n = 4 \end{aligned}$$

2. If A is a 3×3 non-singular matrix such that $AA^T = A^T A$ and $B = A^{-1}A^T$, then $BB^T =$

- (1) A (2) B (3) I_3 (4) B^T

[Ans. (3) I_3]

Hint : $BB^T = (A^{-1}A^T)(A^{-1}A^T)^T$

$$\begin{aligned} &= (A^{-1}A^T)(A^T)^T \cdot (A^{-1})^T \\ &= (A^{-1}A^T)A(A^{-1})^T = A^{-1}(A \cdot A^T)(A^{-1})^T \\ &= (A^{-1}A) \cdot A^T(A^T)^{-1} [\because (A^{-1})^T = (A^T)^{-1}] \\ &= I \cdot I = I \quad [\because A^T \cdot (A^T)^{-1} = I] \end{aligned}$$

3. If $A = \begin{bmatrix} 3 & 5 \\ 1 & 2 \end{bmatrix}$, $B = \text{adj} A$ and $C = 3A$, then $\frac{|\text{adj} B|}{|C|} =$

[Govt. MQP-2019]

- (1) $\frac{1}{3}$ (2) $\frac{1}{9}$ (3) $\frac{1}{4}$ (4) 1

[Ans. (2) $\frac{1}{9}$]

$$\begin{aligned} \text{Hint : } \frac{|\text{adj} B|}{|C|} &= \frac{|\text{adj}(\text{adj} A)|}{|3A|} = \frac{|A|^{(n-1)^2}}{|3|^2 \cdot |A|} \\ &= \frac{|A|^2}{9 \cdot |A|} = \frac{|A|}{9} = \frac{1}{9} \end{aligned}$$

4. If $A \begin{bmatrix} 1 & -2 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} 6 & 0 \\ 0 & 6 \end{bmatrix}$, then $A =$

(1) $\begin{bmatrix} 1 & -2 \\ 1 & 4 \end{bmatrix}$ (2) $\begin{bmatrix} 1 & 2 \\ -1 & 4 \end{bmatrix}$

(3) $\begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$ (4) $\begin{bmatrix} 4 & -1 \\ 2 & 1 \end{bmatrix}$
[Ans. (3) $\begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$]

Hint : $AX = B$

$\Rightarrow A = BX^{-1}$ where

$X = \begin{bmatrix} 1 & -2 \\ 1 & 4 \end{bmatrix}, B = \begin{bmatrix} 6 & 0 \\ 0 & 6 \end{bmatrix}$

$A = \begin{bmatrix} 6 & 0 \\ 0 & 6 \end{bmatrix} \cdot \frac{1}{|X|} \cdot \text{adj}(X)$
 $= \cancel{6} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \cdot \frac{1}{\cancel{6}} \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$
 $= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix} = \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$

5. If $A = \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}$, then $9I_2 - A =$ [PTA - 2; FRT - 2022]

(1) A^{-1} (2) $\frac{A^{-1}}{2}$ (3) $3A^{-1}$ (4) $2A^{-1}$

[Ans. (4) $2A^{-1}$]

Hint : $9I - A = \begin{bmatrix} 9 & 0 \\ 0 & 9 \end{bmatrix} - \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}$

$= \begin{bmatrix} 9-7 & 0-3 \\ 0-4 & 9-2 \end{bmatrix}$
 $= \begin{bmatrix} 2 & -3 \\ -4 & 7 \end{bmatrix} = \text{adj } A$

But $A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{2} \text{adj } A$

$\Rightarrow \text{adj } A = 2A^{-1}$

6. If $A = \begin{bmatrix} 2 & 0 \\ 1 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 4 \\ 2 & 0 \end{bmatrix}$ then $|\text{adj}(AB)| =$
 [Qy. - 2023]

(1) -40 (2) -80 (3) -60 (4) -20

[Ans. (2) -80]

Hint : $AB = \begin{bmatrix} 2+0 & 8+0 \\ 1+10 & 4+0 \end{bmatrix} = \begin{bmatrix} 2 & 8 \\ 11 & 4 \end{bmatrix}$

$\text{adj}(AB) = \begin{bmatrix} 4 & -8 \\ -11 & 2 \end{bmatrix}$

$|\text{adj}(AB)| = 8 - 88 = -80$

7. If $P = \begin{bmatrix} 1 & x & 0 \\ 1 & 3 & 0 \\ 2 & 4 & -2 \end{bmatrix}$ is the adjoint of 3×3 matrix

A and $|A| = 4$, then x is

(1) 15 (2) 12 (3) 14 (4) 11

[Ans. (4) 11]

Hint : $|\text{adj } A| = |A|^{n-1}$

$1 \begin{vmatrix} 3 & 0 \\ 4 & -2 \end{vmatrix} - x \begin{vmatrix} 1 & 0 \\ 2 & -2 \end{vmatrix} + 0 = 4^{3-1}$

$\Rightarrow -6 - x(-2) = 4^2$

$\Rightarrow -6 + 2x = 16$

$\Rightarrow 2x = 22$

$\Rightarrow x = 11$

8. If $A = \begin{bmatrix} 3 & 1 & -1 \\ 2 & -2 & 0 \\ 1 & 2 & -1 \end{bmatrix}$ and $A^{-1} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$

then the value of a_{23} is

[PTA - 5]

(1) 0 (2) -2 (3) -3 (4) -1

[Ans. (4) -1]

Hint : $|A| = 3 \begin{vmatrix} -2 & 0 \\ 2 & -1 \end{vmatrix} - 1 \begin{vmatrix} 2 & 0 \\ 1 & -1 \end{vmatrix} - 1 \begin{vmatrix} 2 & -2 \\ 1 & 2 \end{vmatrix}$

$= 3(2) - 1(-2) - 1(4 + 2)$

$= 6 + 2 - 6 = 2$

$a_{23} = \frac{1}{|A|} \text{co-factor of } a_{32} = \frac{1}{2} \times - \begin{vmatrix} 3 & -1 \\ 2 & 0 \end{vmatrix}$

$= -\frac{1}{2}(0 + 2) = \frac{-2}{2} = -1$

9. If A, B and C are invertible matrices of some order, then which one of the following is not true?
 [Aug. - 2021; Mar. - 2024]

(1) $\text{adj } A = |A|A^{-1}$

(2) $\text{adj}(AB) = (\text{adj } A)(\text{adj } B)$

(3) $\det A^{-1} = (\det A)^{-1}$

(4) $(ABC)^{-1} = C^{-1}B^{-1}A^{-1}$

[Ans. (2) $\text{adj}(AB) = (\text{adj } A)(\text{adj } B)$]

10. If $(AB)^{-1} = \begin{bmatrix} 12 & -17 \\ -19 & 27 \end{bmatrix}$ and $A^{-1} = \begin{bmatrix} 1 & -1 \\ -2 & 3 \end{bmatrix}$,

then $B^{-1} =$ [Mar. - 2020; Hy. - 2023]

- (1) $\begin{bmatrix} 2 & -5 \\ -3 & 8 \end{bmatrix}$ (2) $\begin{bmatrix} 8 & 5 \\ 3 & 2 \end{bmatrix}$
 (3) $\begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix}$ (4) $\begin{bmatrix} 8 & -5 \\ -3 & 2 \end{bmatrix}$

[Ans. (1) $\begin{bmatrix} 2 & -5 \\ -3 & 8 \end{bmatrix}$]

Hint : Since $(AB)^{-1} = B^{-1}A^{-1}$, we get,

$$\begin{bmatrix} 12 & -17 \\ -19 & 27 \end{bmatrix} = B^{-1} \begin{bmatrix} 1 & -1 \\ -2 & 3 \end{bmatrix}$$

Let $X = B^{-1}Y$

$$B^{-1} = XY^{-1} = \begin{bmatrix} 12 & -17 \\ -19 & 27 \end{bmatrix} \cdot \frac{1}{|Y|} \cdot (\text{adj } Y)$$

$$= \begin{bmatrix} 12 & -17 \\ -19 & 27 \end{bmatrix} \cdot \frac{1}{1} \begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 12 & -17 \\ -19 & 27 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 36-34 & 12-17 \\ -57+54 & -19+27 \end{bmatrix} = \begin{bmatrix} 2 & -5 \\ -3 & 8 \end{bmatrix}$$

11. If $A^T \cdot A^{-1}$ is symmetric, then $A^2 =$

[Sep. - 2020; FRT-2022; June & Qy. - 2023]

- (1) A^{-1} (2) $(A^T)^2$
 (3) A^T (4) $(A^{-1})^2$

[Ans : (2) $(A^T)^2$]

Hint : $\Rightarrow A^T A^{-1} = (A^T A^{-1})^T$

$$= (A^{-1})^T (A^T)^T = (A^{-1})^T A$$

$$\Rightarrow A^T A^{-1} = (A^{-1})^T A$$

$$\Rightarrow A^T A^{-1} = (A^T)^{-1} \cdot A$$

[Premultiplying by A^T and post multiplying by A on both sides, we get]

$$A^T (A^T A^{-1}) A = A^T [(A^T)^{-1} \cdot A] A$$

$$\Rightarrow (A^T)^2 (A^{-1} A) = A^T (A^T)^{-1} \cdot A^2$$

$$\Rightarrow (A^T)^2 (I) = I A^2$$

$$\Rightarrow (A^T)^2 = A^2$$

12. If A is a non-singular matrix such that

$$A^{-1} = \begin{bmatrix} 5 & 3 \\ -2 & -1 \end{bmatrix}, \text{ then } (A^T)^{-1} =$$

[PTA -1 ; Hy - 2019; July - 2022; Mar. - 2024]

- (1) $\begin{bmatrix} -5 & 3 \\ 2 & 1 \end{bmatrix}$ (2) $\begin{bmatrix} 5 & 3 \\ -2 & -1 \end{bmatrix}$
 (3) $\begin{bmatrix} -1 & -3 \\ 2 & 5 \end{bmatrix}$ (4) $\begin{bmatrix} 5 & -2 \\ 3 & -1 \end{bmatrix}$

Hint : $(A^T)^{-1} = (A^{-1})^T = \begin{bmatrix} 5 & -2 \\ 3 & -1 \end{bmatrix}$ [Ans : (4) $\begin{bmatrix} 5 & -2 \\ 3 & -1 \end{bmatrix}$]

13. If $A = \begin{bmatrix} 3 & 4 \\ 5 & 5 \\ x & 3 \\ 5 & 5 \end{bmatrix}$ and $A^T = A^{-1}$, then the value of x is

[July - 2022]

- (1) $\frac{-4}{5}$ (2) $\frac{-3}{5}$ (3) $\frac{3}{5}$ (4) $\frac{4}{5}$

[Ans: (1) $\frac{-4}{5}$]

Hint : Since $A^T = A^{-1}$, $AA^T = A^T A = I$ [$\therefore$ they are orthogonal]

$$\Rightarrow \begin{bmatrix} 3 & 4 \\ 5 & 5 \\ x & 3 \\ 5 & 5 \end{bmatrix} \begin{bmatrix} 3 & x \\ 4 & 3 \\ 5 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} \frac{9}{25} + \frac{16}{25} & \frac{3x}{5} + \frac{12}{25} \\ \frac{3x}{5} + \frac{12}{25} & x^2 + \frac{9}{25} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow \frac{3x}{5} + \frac{12}{25} = 0 \quad [\text{Equating } a_{12} \text{ both sides}]$$

$$\Rightarrow \frac{3x}{5} = \frac{-12}{25} \Rightarrow x = \frac{-12}{25} \times \frac{5}{3} = \frac{-4}{5}$$

14. If $A = \begin{bmatrix} 1 & \tan \frac{\theta}{2} \\ -\tan \frac{\theta}{2} & 1 \end{bmatrix}$ and $AB = I_2$, then $B =$ [PTA -3]

- (1) $\left(\cos^2 \frac{\theta}{2}\right) A$ (2) $\left(\cos^2 \frac{\theta}{2}\right) A^T$
 (3) $(\cos^2 \theta) I$ (4) $\left(\sin^2 \frac{\theta}{2}\right) A$

[Ans: (2) $\left(\cos^2 \frac{\theta}{2}\right) A^T$]

Hint : $B = A^{-1} = \frac{1}{|A|} \text{adj } A$

$$= \frac{1}{1 + \tan^2 \frac{\theta}{2}} \begin{bmatrix} 1 & -\tan \frac{\theta}{2} \\ \tan \frac{\theta}{2} & 1 \end{bmatrix} = \frac{1}{\sec^2 \frac{\theta}{2}} \cdot A^T = \left(\cos^2 \frac{\theta}{2} \right) A^T$$

15. If $A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ and $A(\text{adj } A) = \begin{bmatrix} k & 0 \\ 0 & k \end{bmatrix}$, then $k =$ [FRT - 2022]

- (1) 0 (2) $\sin \theta$ (3) $\cos \theta$ (4) 1

[Ans: (4) 1]

Hint : We know $A(\text{adj } A) = (\text{adj } A)A = |A| I$
 $\Rightarrow |A| = k$

$$\therefore k = \begin{vmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{vmatrix} = \cos^2 \theta + \sin^2 \theta = 1$$

16. If $A = \begin{bmatrix} 2 & 3 \\ 5 & -2 \end{bmatrix}$ be such that $\lambda A^{-1} = A$, then λ is [May - 2022]

- (1) 17 (2) 14 (3) 19 (4) 21

[Ans: (3) 19]

Hint : $\lambda \cdot \frac{1}{|A|} \text{adj } A = A$

$$\Rightarrow \lambda \cdot \frac{1}{(-4-15)} \begin{bmatrix} -2 & -3 \\ -5 & 2 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 5 & -2 \end{bmatrix}$$

$$\Rightarrow \frac{-\lambda}{19} \begin{bmatrix} -2 & -3 \\ -5 & 2 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 5 & -2 \end{bmatrix}$$

$$\Rightarrow \frac{\lambda}{19} \begin{bmatrix} 2 & 3 \\ 5 & -2 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 5 & -2 \end{bmatrix}$$

$$\Rightarrow \frac{\lambda}{19} = 1 \Rightarrow \lambda = 19$$

17. If $\text{adj } A = \begin{bmatrix} 2 & 3 \\ 4 & -1 \end{bmatrix}$ and $\text{adj } B = \begin{bmatrix} 1 & -2 \\ -3 & 1 \end{bmatrix}$ then $\text{adj } (AB)$ is [Hy - 2019]

- (1) $\begin{bmatrix} -7 & -1 \\ 7 & -9 \end{bmatrix}$ (2) $\begin{bmatrix} -6 & 5 \\ -2 & -10 \end{bmatrix}$
 (3) $\begin{bmatrix} -7 & 7 \\ -1 & -9 \end{bmatrix}$ (4) $\begin{bmatrix} -6 & -2 \\ 5 & -10 \end{bmatrix}$

[Ans: (2) $\begin{bmatrix} -6 & 5 \\ -2 & -10 \end{bmatrix}$]

Hint : $\text{adj } (AB) = (\text{adj } B)(\text{adj } A)$

$$= \begin{bmatrix} 1 & -2 \\ -3 & 1 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 4 & -1 \end{bmatrix} = \begin{bmatrix} 2-8 & 3+2 \\ -6+4 & -9-1 \end{bmatrix} = \begin{bmatrix} -6 & 5 \\ -2 & -10 \end{bmatrix}$$

18. The rank of the matrix $\begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 4 & 6 & 8 \\ -1 & -2 & -3 & -4 \end{bmatrix}$ is [Qy-2019; June-2023]

- (1) 1 (2) 2 (3) 4 (4) 3

[Ans: (1) 1]

Hint : $\begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 4 & 6 & 8 \\ -1 & -2 & -3 & -4 \end{bmatrix} \xrightarrow{\substack{R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 + R_1}} \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$

$\therefore$ Rank is 1 [$\because$ only one non-zero row]

19. If $x^a y^b = e^m, x^c y^d = e^n, \Delta_1 = \begin{bmatrix} m & b \\ n & d \end{bmatrix}, \Delta_2 = \begin{bmatrix} a & m \\ c & n \end{bmatrix}, \Delta_3 = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then the values of x and y are respectively, [PTA -3; Sep. - 2020]

- (1) $e^{(\Delta_2/\Delta_1)}, e^{(\Delta_3/\Delta_1)}$
 (2) $\log(\Delta_1/\Delta_3), \log(\Delta_2/\Delta_3)$
 (3) $\log(\Delta_2/\Delta_1), \log(\Delta_3/\Delta_1)$
 (4) $e^{(\Delta_1/\Delta_3)}, e^{(\Delta_2/\Delta_3)}$ [Ans: (4) $e^{(\Delta_1/\Delta_3)}, e^{(\Delta_2/\Delta_3)}$]

Hint :

$$x^a y^b = e^m \Rightarrow a \log x + b \log y = m$$

[Taking log both sides]

$$x^c y^d = e^n \Rightarrow c \log x + d \log y = n$$

$$\text{put } \Delta_3 = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$\Delta_1 = \begin{bmatrix} m & b \\ n & d \end{bmatrix}, \Delta_2 = \begin{bmatrix} a & m \\ c & n \end{bmatrix}$$

$$\log x = \frac{\Delta_1}{\Delta_3} \text{ and } \log y = \frac{\Delta_2}{\Delta_3}$$

$$\therefore x = e^{\Delta_1/\Delta_3} \text{ and } y = e^{\Delta_2/\Delta_3}$$

20. Which of the following is/are correct?

- (i) Adjoint of a symmetric matrix is also a symmetric matrix.
 (ii) Adjoint of a diagonal matrix is also a diagonal matrix.
 (iii) If A is a square matrix of order n and λ is a scalar, then $\text{adj } (\lambda A) = \lambda^n \text{adj } (A)$.
 (iv) $A(\text{adj } A) = (\text{adj } A)A = |A|I$

- (1) Only (i) (2) (ii) and (iii)

- (3) (iii) and (iv) (4) (i), (ii) and (iv)

[Ans: (4) (i) (ii) and (iv)]

21. If $\rho(A) = \rho([A|B])$, then the system $AX = B$ of linear equations is [PTA-6; Mar. - 2020]

- (1) consistent and has a unique solution
- (2) consistent
- (3) consistent and has infinitely many solution
- (4) inconsistent [Ans: (2) Consistent]

22. If $0 \leq \theta \leq \pi$ and the system of equations $x + (\sin \theta)y - (\cos \theta)z = 0$, $(\cos \theta)x - y + z = 0$, $(\sin \theta)x + y - z = 0$ has a non-trivial solution then θ is [Qy-2019]

- (1) $\frac{2\pi}{3}$
- (2) $\frac{3\pi}{4}$
- (3) $\frac{5\pi}{6}$
- (4) $\frac{\pi}{4}$

[Ans: (4) $\frac{\pi}{4}$]

Hint : $A = \begin{bmatrix} 1 & \sin \theta & -\cos \theta \\ \cos \theta & -1 & 1 \\ \sin \theta & 1 & -1 \end{bmatrix}$

The system has non-trivial solution if $|A| = 0$

$$\begin{vmatrix} 1 & \sin \theta & -\cos \theta \\ \cos \theta & -1 & 1 \\ \sin \theta & 1 & -1 \end{vmatrix} = 0$$

$$\Rightarrow 1 \begin{vmatrix} -1 & 1 \\ 1 & -1 \end{vmatrix} - \sin \theta \begin{vmatrix} \cos \theta & 1 \\ \sin \theta & -1 \end{vmatrix} - \cos \theta \begin{vmatrix} \cos \theta & -1 \\ \sin \theta & 1 \end{vmatrix} = 0$$

$$\Rightarrow 1(1-1) - \sin \theta (-\cos \theta - \sin \theta) - \cos \theta (\cos \theta + \sin \theta) = 0$$

$$\Rightarrow \sin \theta \cos \theta + \sin^2 \theta - \cos^2 \theta - \cos \theta \sin \theta = 0$$

$$\Rightarrow \cos^2 \theta - \sin^2 \theta = 0 \Rightarrow \cos 2\theta = 0$$

$$\Rightarrow \cos 2\theta = \cos \frac{\pi}{2} \quad \left[\because \cos \frac{\pi}{2} = 0 \right]$$

$$\Rightarrow 2\theta = \frac{\pi}{2} \Rightarrow \theta = \frac{\pi}{4}$$

23. The augmented matrix of a system of linear

equations is $\begin{bmatrix} 1 & 2 & 7 & 3 \\ 0 & 1 & 4 & 6 \\ 0 & 0 & \lambda - 7 & \mu + 5 \end{bmatrix}$. The system

has infinitely many solutions if

- (1) $\lambda = 7, \mu \neq -5$
- (2) $\lambda = -7, \mu = 5$
- (3) $\lambda \neq 7, \mu \neq -5$
- (4) $\lambda = 7, \mu = -5$

[Ans: (4) $\lambda = 7, \mu = -5$]

Hint : When $\lambda = 7$ and $\mu = -5$,

$$[A|B] = \begin{bmatrix} 1 & 2 & 7 & 3 \\ 0 & 1 & 4 & 6 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\rho(A) = \rho([A|B]) = 2 < 3$, the number of unknowns.

∴ The system is consistent and has infinitely many solutions.

24. Let $A = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$ and $4B = \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & x \\ -1 & 1 & 3 \end{bmatrix}$.

If B is the inverse of A , then the value of x is

- (1) 2
- (2) 4
- (3) 3
- (4) 1

[Ans: (4) 1]

Hint : $A = B^{-1} \Rightarrow A \cdot B = B^{-1} \cdot B \Rightarrow AB = I$

$$\therefore \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix} \frac{1}{4} \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & x \\ -1 & 1 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\frac{1}{4} a_{13} = 0 \Rightarrow \frac{1}{4} \cdot [-2 - x + 3] = 0$$

$$\Rightarrow -x + 1 = 0 \times 4 = 0 \Rightarrow x = 1$$

25. If $A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$, then $\text{adj}(\text{adj } A)$ is [PTA - 2]

$$(1) \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix} \quad (2) \begin{bmatrix} 6 & -6 & 8 \\ 4 & -6 & 8 \\ 0 & -2 & 2 \end{bmatrix}$$

$$(3) \begin{bmatrix} -3 & 3 & -4 \\ -2 & 3 & -4 \\ 0 & 1 & -1 \end{bmatrix} \quad (4) \begin{bmatrix} 3 & -3 & 4 \\ 0 & -1 & 1 \\ 2 & -3 & 4 \end{bmatrix}$$

[Ans: (1) $\begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$]

Hint : $\text{adj}(\text{adj } A) = |A|^{n-2} A = |A| \cdot A$

$$|A| = \begin{vmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{vmatrix}$$

$$= 3 \begin{vmatrix} -3 & 4 \\ -1 & 1 \end{vmatrix} + 3 \begin{vmatrix} 2 & 4 \\ 0 & 1 \end{vmatrix} + 4 \begin{vmatrix} 2 & -3 \\ 0 & -1 \end{vmatrix}$$

$$= 3(-3+4) + 3(2-0) + 4(-2-0)$$

$$= 3(1) + 6 - 8 = 1$$

$$\therefore \text{adj}(\text{adj } A) = 1 \cdot A = A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$$

PTA QUESTION & ANSWERS

1 MARK

1. If A and B are orthogonal, then $(AB)^T (AB)$ is [PTA - 1]

(1) A (2) B (3) I (4) A^T

Hint : $AA^T = A^T A = I$ [Ans: (3) I]

$$BB^T = B^T B = I$$

$$\begin{aligned}(AB)^T (AB) &= B^T A^T (AB) = B^T (A^T A) B \\ &= B^T (IB) = I\end{aligned}$$

2. The adjoint of 3×3 matrix P is $\begin{bmatrix} -1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$, then the possible value(s) of the determinant P is (are) [PTA - 4]

(1) 3 (2) -3 (3) ± 3 (4) $\pm \sqrt{3}$

[Ans: (3) ± 3]

Hint : $\begin{vmatrix} -1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \end{vmatrix} = -1[1-4] - 2[1-4] + 2[2-2]$
 $= -1(-3) - 2(-3) + 0 = 3 + 6 = 9$
 $|P| = \pm \sqrt{9} = \pm 3$

3. If A is a 3×3 matrix such that $|3 \text{adj } A| = 3$ then $|A|$ is equal to [PTA - 5]

(1) $\frac{1}{3}$ (2) $-\frac{1}{3}$ (3) $\pm \frac{1}{3}$ (4) ± 3

Hint : $|3 \text{adj } A| = 27 |\text{adj } A|$ [Ans: (3) $\pm \frac{1}{3}$]

$$3 = 27|A|^2$$

$$|A|^2 = \frac{1}{9}$$

$$|A| = \pm \frac{1}{3}$$

4. Let A be a non-singular matrix then which one of the following is false [PTA - 6]

(1) $(\text{adj } A)^{-1} = \frac{A}{|A|}$
 (2) I is an orthogonal matrix
 (3) $\text{adj}(\text{adj } A) = |A|^n A$
 (4) If A is symmetric then $\text{adj } A$ is symmetric

[Ans: (3) $\text{adj}(\text{adj } A) = |A|^n A$]

Hint : $\text{adj}(\text{adj } A) = |A|^{n-2} A$

2 MARKS

1. Prove that $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ is orthogonal.

[PTA - 1; FRT -2022; Mar. & Qy. - 2023]

Sol. Let $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$
 Then, $A^T = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}^T = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$
 $= \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$
 $= \begin{bmatrix} \cos^2 \theta + \sin^2 \theta & \cos \theta \sin \theta - \sin \theta \cos \theta \\ \sin \theta \cos \theta - \cos \theta \sin \theta & \sin^2 \theta + \cos^2 \theta \end{bmatrix}$
 $= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2.$

Similarly, we get $A^T A = I_2.$

Hence $AA^T = A^T A = I_2 \Rightarrow A$ is orthogonal.

2. If $A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 0 \\ 2 & 5 \end{bmatrix}$, find $\text{adj}(AB)$. [PTA - 3]

Sol. $AB = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 2 & 5 \end{bmatrix} = \begin{bmatrix} 8+6 & 0+15 \\ 4+4 & 0+10 \end{bmatrix} = \begin{bmatrix} 14 & 15 \\ 8 & 10 \end{bmatrix}$
 $\text{adj}(AB) = \begin{bmatrix} 10 & -15 \\ -8 & 14 \end{bmatrix}$

3 MARKS

1. If $A = \begin{bmatrix} 3 & -2 \\ \lambda & -2 \end{bmatrix}$, find the value of λ so that

$$A^2 = \lambda A - 2I. \quad [\text{PTA} - 2]$$

Sol. $A^2 = \begin{bmatrix} 3 & -2 \\ \lambda & -2 \end{bmatrix} \begin{bmatrix} 3 & -2 \\ \lambda & -2 \end{bmatrix}$
 $= \begin{bmatrix} 9-2\lambda & -6+4 \\ 3\lambda-2\lambda & -2\lambda+4 \end{bmatrix} = \begin{bmatrix} 9-2\lambda & -2 \\ \lambda & -2\lambda+4 \end{bmatrix}$
 $\lambda A - 2I = \lambda \begin{bmatrix} 3 & -2 \\ \lambda & -2 \end{bmatrix} - 2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
 $= \begin{bmatrix} 3\lambda & -2\lambda \\ \lambda^2 & -2\lambda \end{bmatrix} - \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} 3\lambda-2 & -2\lambda \\ \lambda^2 & -2\lambda-2 \end{bmatrix}$
 $\therefore -2\lambda = -2$
 $\lambda = 1$

2. Find the rank of the matrix $\begin{bmatrix} 4 & -2 & 6 & 8 \\ 1 & 1 & -3 & -1 \\ 15 & -3 & 9 & 21 \end{bmatrix}$ [PTA - 4]

Sol. $\begin{bmatrix} 4 & -2 & 6 & 8 \\ 1 & 1 & -3 & -1 \\ 15 & -3 & 9 & 21 \end{bmatrix} \xrightarrow{R_1 \leftrightarrow R_2} \begin{bmatrix} 1 & 1 & -3 & -1 \\ 4 & -2 & 6 & 8 \\ 15 & -3 & 9 & 21 \end{bmatrix}$
 $R_2 \rightarrow R_2 + (4)R_1$ $\begin{bmatrix} 1 & 1 & -3 & -1 \\ 0 & -6 & 18 & 12 \\ 0 & -18 & 54 & 36 \end{bmatrix}$
 $R_3 \rightarrow R_3 + (-15)R_1$ $\begin{bmatrix} 1 & 1 & -3 & -1 \\ 0 & -6 & 18 & 12 \\ 0 & -18 & 54 & 36 \end{bmatrix}$
 $R_2 \rightarrow \frac{R_2}{-6}$ $\begin{bmatrix} 1 & 1 & -3 & -1 \\ 0 & 1 & -3 & -2 \\ 0 & -18 & 54 & 36 \end{bmatrix}$
 $R_3 \rightarrow \frac{R_3}{-18}$ $\begin{bmatrix} 1 & 1 & -3 & -1 \\ 0 & 1 & -3 & -2 \\ 0 & 1 & -3 & -2 \end{bmatrix} \xrightarrow{R_3 \rightarrow R_3 - R_2} \begin{bmatrix} 1 & 1 & -3 & -1 \\ 0 & 1 & -3 & -2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$

3. Solve by matrix inversion method :
 $5x + 2y = 4, 7x + 3y = 5.$

Sol. Matrix form $AX = B$ [PTA - 5]
 $A = \begin{bmatrix} 5 & 2 \\ 7 & 3 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 4 \\ 5 \end{bmatrix}$
 $|A| = \begin{vmatrix} 5 & 2 \\ 7 & 3 \end{vmatrix} = 15 - 14 = 1$
 $A^{-1} = \begin{bmatrix} 3 & -2 \\ -7 & 5 \end{bmatrix}$
 $X = A^{-1}B$
 $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 & -2 \\ -7 & 5 \end{bmatrix} \begin{bmatrix} 4 \\ 5 \end{bmatrix}$
 $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 12 & -10 \\ -28 & 25 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \end{bmatrix}$
 $x = 2, y = -3$

4. Find the adjoint of the matrix $A = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$
 and verify that $A(adjA) = (adjA)A = |A|I$.

Sol. $A = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$ [PTA - 6]
 $adj A = \begin{bmatrix} -5 & -3 \\ -2 & 1 \end{bmatrix}$
 $A(adjA) = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix} \begin{bmatrix} -5 & -3 \\ -2 & 1 \end{bmatrix}$
 $= \begin{bmatrix} -5-6 & -3+3 \\ -10+10 & -6-5 \end{bmatrix} = \begin{bmatrix} -11 & 0 \\ 0 & -11 \end{bmatrix} \dots(1)$
 $(adj A)A = \begin{bmatrix} -5 & -3 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$

$$= \begin{bmatrix} -5-6 & -15+15 \\ -2+2 & -6-5 \end{bmatrix} = \begin{bmatrix} -11 & 0 \\ 0 & -11 \end{bmatrix} \dots(2)$$

$$|A| = \begin{vmatrix} 1 & 3 \\ 2 & -5 \end{vmatrix} = -5 - 6 = -11$$

$$|A|I = -11 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} -11 & 0 \\ 0 & -11 \end{bmatrix} \dots(3)$$

From (1), (2) and (3)

$$A(adjA) = (adj A)A = |A|I$$

5 MARKS

1. Examine the consistency of the system of equation $4x + 3y + 6z = 25, x + 5y + 7z = 13.$
 $2x + 9y + z = 1.$ It is consistent then solve.

[PTA - 1]

Sol. Transforming the augmented matrix to echelon form, we get

$$\begin{bmatrix} 4 & 3 & 6 & 25 \\ 1 & 5 & 7 & 13 \\ 2 & 9 & 1 & 1 \end{bmatrix} \xrightarrow{R_1 \rightarrow R_2} \begin{bmatrix} 1 & 5 & 7 & 13 \\ 4 & 3 & 6 & 25 \\ 2 & 9 & 1 & 1 \end{bmatrix}$$

$$\xrightarrow{\substack{R_2 \rightarrow R_2 - 4R_1, \\ R_3 \rightarrow R_3 - 2R_1}} \begin{bmatrix} 1 & 5 & 7 & 13 \\ 0 & -17 & -22 & -27 \\ 0 & -1 & -13 & -25 \end{bmatrix}$$

$$\xrightarrow{\substack{R_2 \rightarrow R_2 \div (-1), \\ R_3 \rightarrow R_3 \div (-1)}} \begin{bmatrix} 1 & 5 & 7 & 13 \\ 0 & 17 & 22 & 27 \\ 0 & 1 & 13 & 25 \end{bmatrix}$$

$$\xrightarrow{R_3 \rightarrow 17R_3 - R_2} \begin{bmatrix} 1 & 5 & 7 & 13 \\ 0 & 17 & 22 & 27 \\ 0 & 0 & 199 & 398 \end{bmatrix}$$

The equivalent system is written by using the echelon form:

$$x + 5y + 7z = 13, \dots (1)$$

$$17y + 22z = 27, \dots (2)$$

$$199z = 398. \dots (3)$$

$$\text{From (3), we get } z = \frac{398}{199} = 2.$$

$$\text{Substituting } z = 2 \text{ in (2), we get } \frac{27 - 22 \times 2}{17} = \frac{-17}{17} = -1$$

$$\text{Substituting } z = 2, y = -1 \text{ in (1), we get}$$

$$x = 13 - 5 \times (-1) - 7 \times 2 = 4.$$

So, the solution is $(x = 4, y = -1, z = 2).$

Note. The above method of going from the last equation to the first equation is called the method of back substitution.

2. Find the inverse of the non-singular matrix

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 3 & 2 & 1 \\ 2 & 1 & 2 \end{bmatrix}, \text{ by elementary transformations.}$$

[PTA - 6]

Sol. Applying Gauss-Jordan method, we get

$$[A|I_3] = \left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 3 & 2 & 1 & 0 & 1 & 0 \\ 2 & 1 & 2 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow \frac{1}{2}R_1} \left[\begin{array}{ccc|ccc} 1 & \left(\frac{1}{2}\right) & \left(\frac{1}{2}\right) & \left(\frac{1}{2}\right) & 0 & 0 \\ 3 & 2 & 1 & 0 & 1 & 0 \\ 2 & 1 & 2 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - 3R_1 \\ R_3 \rightarrow R_3 - 2R_1 \end{array}} \left[\begin{array}{ccc|ccc} 1 & \left(\frac{1}{2}\right) & \left(\frac{1}{2}\right) & \left(\frac{1}{2}\right) & 0 & 0 \\ 3 & 2 & 1 & -\left(\frac{3}{2}\right) & 1 & 0 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow 2R_2} \left[\begin{array}{ccc|ccc} 1 & \left(\frac{1}{2}\right) & \left(\frac{1}{2}\right) & \left(\frac{1}{2}\right) & 0 & 0 \\ 0 & 1 & -1 & -1 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_1 - \frac{1}{2}R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 2 & -1 & 0 \\ 0 & 1 & -1 & -3 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{\begin{array}{l} R_1 \rightarrow R_1 - R_3 \\ R_2 \rightarrow R_2 + R_3 \end{array}} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 3 & -1 & -1 \\ 0 & 1 & 0 & -4 & 2 & 1 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right]$$

$$\text{So, } A^{-1} = \begin{bmatrix} 3 & -1 & -1 \\ -4 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$$

3. Investigate for what values of λ and μ the system of linear equations $x + 2y + z = 7$, $x + y + \lambda z = \mu$ has (i) no solution (ii) a unique solution (iii) an infinite number of solutions.

[PTA - 2]

Sol. Here the number of unknowns is 3.The matrix form of the system is $AX = B$, where

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 1 & \lambda \\ 1 & 3 & -5 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 7 \\ \mu \\ 5 \end{bmatrix}$$

Applying elementary row operations on the augmented matrix $[A|B]$, we get

$$[A|B] = \left[\begin{array}{ccc|c} 1 & 2 & 1 & 7 \\ 1 & 1 & \lambda & \mu \\ 1 & 3 & -5 & 5 \end{array} \right] \xrightarrow{R_2 \leftrightarrow R_3} \left[\begin{array}{ccc|c} 1 & 2 & 1 & 7 \\ 1 & 3 & -5 & 5 \\ 1 & 1 & \lambda & \mu \end{array} \right]$$

$$\xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - R_1, \\ R_3 \rightarrow R_3 - R_1, \end{array}} \left[\begin{array}{ccc|c} 1 & 2 & 1 & 7 \\ 0 & 1 & -6 & -2 \\ 0 & -1 & \lambda - 1 & \mu - 7 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 + R_2} \left[\begin{array}{ccc|c} 1 & 2 & 1 & 7 \\ 0 & 1 & -6 & -2 \\ 0 & 0 & \lambda - 7 & \mu - 9 \end{array} \right]$$

(i) If $\lambda = 7$ and $\mu \neq 9$, then $\rho(A) = 2$ and $([A|B]) = 3$ and So $\rho(A) \neq 2([A|B])$. Hence the given system is inconsistent and has no solution.(ii) If $\lambda = 7$ and μ is any real number, then $\rho(A) = 3$ and $([A|B]) = 3$.So $\rho(A) = ([A|B]) = 3 =$ Number of unknowns Hence the given system is consistent and has a unique solution.(iii) If $\lambda = 7$ and $\mu = 9$, then $\rho(A) = 2$ and $\rho([A|B]) = 2$.So, $\rho(A) = ([A|B]) = 2$ Number of unknowns. Hence the given system is consistent and has infinite number of solutions.**GOVT. EXAM QUESTION & ANSWERS****1 MARK****I. Choose the Correct or the most suitable answer from the given four alternatives :****1. If the inverse of the matrix $\begin{bmatrix} 1 & 2 \\ 3 & -5 \end{bmatrix}^n$ is $\frac{1}{11} \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then the ascending order of a, b, c, d is**

[Govt. MQP-2019]

- (1) a, b, c, d (2) d, b, c, a
 (3) c, a, b, d (4) b, a, c, d

[Ans: (2) d, b, c, a]**Hint :** Inverse matrix = $\frac{1}{-5-6} \begin{bmatrix} -5 & -2 \\ -3 & 1 \end{bmatrix}$

$$= \frac{-1}{11} \begin{bmatrix} -5 & -2 \\ -3 & 1 \end{bmatrix} = \begin{bmatrix} \frac{5}{11} & \frac{2}{11} \\ \frac{3}{11} & \frac{-1}{11} \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

2. If A is an orthogonal matrix, then |A| is [Qy - 2019]

- (1) 1 (2) -1 (3) ± 1 (4) 0

[Ans: (3) ± 1]

Hint : The determinant of an orthogonal matrix is equal to 1 or -1.

3. The system of linear equations $x + y + z = 2$, $2x + y - z = 3$, $3x + 2y + kz = 4$ has a unique solution if [Qy - 2019]

- (1) $k \neq 0$ (2) $-1 < k < 1$
(3) $-2 < k < 2$ (4) $k = 0$ [Ans: (1) $k \neq 0$]

Hint : $\begin{vmatrix} 1 & 1 & 1 \\ 2 & 1 & -1 \\ 3 & 2 & k \end{vmatrix} \neq 0$

$$\Rightarrow 1[k+2] - 1[2k+3] + 1[4-3] \neq 0$$

$$\Rightarrow k+2-2k-3+1 \neq 0 \Rightarrow -k+0 \neq 0 \Rightarrow k \neq 0$$

4. The inverse of $\begin{bmatrix} 3 & 1 \\ 5 & 2 \end{bmatrix}$ is : [Aug. - 2021]

- (1) $\begin{bmatrix} 3 & -1 \\ -5 & -3 \end{bmatrix}$ (2) $\begin{bmatrix} 2 & -1 \\ -5 & 3 \end{bmatrix}$
(3) $\begin{bmatrix} -3 & 5 \\ 1 & -2 \end{bmatrix}$ (4) $\begin{bmatrix} -2 & 5 \\ 1 & -3 \end{bmatrix}$
[Ans: (2) $\begin{bmatrix} 2 & -1 \\ -5 & 3 \end{bmatrix}$]

Hint : $A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{1} \begin{bmatrix} 2 & -1 \\ -5 & 3 \end{bmatrix} = \begin{bmatrix} 2 & -1 \\ -5 & 3 \end{bmatrix}$

5. The adjoint of $\begin{bmatrix} -2 & 1 \\ -1 & 4 \end{bmatrix}$ is [FRT - 2022]

- (1) $\begin{bmatrix} 4 & -1 \\ 1 & -2 \end{bmatrix}$ (2) $\begin{bmatrix} -2 & -1 \\ 1 & 4 \end{bmatrix}$
(3) $\begin{bmatrix} -4 & 1 \\ -1 & 2 \end{bmatrix}$ (4) $\begin{bmatrix} 4 & 1 \\ -1 & -2 \end{bmatrix}$
[Ans: (1) $\begin{bmatrix} 4 & -1 \\ 1 & -2 \end{bmatrix}$]

6. Which one of the following is incorrect?

[May - 2022]

- (1) If A is a square matrix of order n , and λ is a scalar, then $\text{Adj}(\lambda A) = \lambda^n (\text{Adj } A)$.
(2) Adjoint of a symmetric matrix is also a symmetric matrix.
(3) $A(\text{Adj } A) = (\text{Adj } A)A = |A|I$.
(4) Adjoint of a diagonal matrix is also a diagonal matrix.

[Ans: (1) If A is a square matrix of order n , and λ is a scalar, then $\text{Adj}(\lambda A) = \lambda^n (\text{Adj } A)$.]

7. A square matrix A of order n has inverse if and only if : [Mar - 2023]

- (1) $\rho(A) > n$ (2) $\rho(A) = n$
(3) $\rho(A) \neq n$ (4) $\rho(A) < n$

[Ans: (2) $\rho(A) = n$]

Hint : A square matrix A of order n has inverse if and only if $p(A) = n$.

8. $|\text{adj}(\text{adj}A)| = |A|^{16}$, then the order of the square matrix A is : [Mar - 2023]

- (1) 2 (2) 3 (3) 5 (4) 4

[Ans: (3) 5]

Hint :

$$|\text{adj}(\text{adj}A)| = |A|^{(n-1)^2}$$

$$(n-1)^2 = 16 \Rightarrow (n-1)^2 = 4^2$$

$$n-1 = 4 \Rightarrow n = 5$$

9. The value of $\int_{-1}^2 |x| dx$ is [June & Hy. - 2023]

- (1) $\frac{1}{2}$ (2) $\frac{3}{2}$ (3) $\frac{5}{2}$ (4) $\frac{7}{2}$

[Ans: (3) $\frac{5}{2}$]

2 MARKS

1. Solve the following system of linear equations by Cramer's rule $2x - y = 3$, $x + 2y = -1$.

[Govt. MQP-2019]

Sol.

$$\begin{aligned} 2x - y &= 3 \\ x + 2y &= -1 \end{aligned}$$

$$\Delta = \begin{vmatrix} 2 & -1 \\ 1 & 2 \end{vmatrix} = 2 \times 2 - (-1)(-1) = 4 - 1 = 3$$

$$\Delta_1 = \begin{vmatrix} 3 & -1 \\ -1 & 2 \end{vmatrix} = 3 \times 2 - (-1) \times (-1) = 6 - 1 = 5$$

$$\Delta_2 = \begin{vmatrix} 2 & 3 \\ 1 & -1 \end{vmatrix} = -2 - 3 = -5$$

$$x = \frac{\Delta_1}{\Delta} = \frac{5}{5} = 1$$

$$y = \frac{\Delta_2}{\Delta} = \frac{-5}{5} = -1$$

2. If $\text{adj } A = \begin{bmatrix} -1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$, find A^{-1} . [Qy - 2019]

Sol. We compute $|\text{adj } A| = 9$
So, we get $A^{-1} = \pm \frac{1}{\sqrt{|\text{adj } A|}} \text{adj } (A)$

$$= \pm \frac{1}{\sqrt{9}} \begin{bmatrix} -1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} = \pm \frac{1}{3} \begin{bmatrix} -1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

3. If A is symmetric, prove that $\text{adj } A$ is also symmetric. [FRT - 2022]

Sol. Refer Text Book Example 1.7

3 MARKS

1. Verify $(AB)^{-1} = B^{-1} A^{-1}$ with $A = \begin{bmatrix} 0 & -3 \\ 1 & 4 \end{bmatrix}$, $B = \begin{bmatrix} -2 & -3 \\ 0 & -1 \end{bmatrix}$ [Sep.- 2020; July - 2022]

Sol. We get $AB = \begin{bmatrix} 0 & -3 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} -2 & -3 \\ 0 & -1 \end{bmatrix}$

$$= \begin{bmatrix} 0+0 & 0+3 \\ -2+0 & -3-4 \end{bmatrix} = \begin{bmatrix} 0 & 3 \\ -2 & -7 \end{bmatrix}$$

$$(AB)^{-1} = \frac{1}{(0+6)} \begin{bmatrix} -7 & -3 \\ 2 & 0 \end{bmatrix} = \frac{1}{6} \begin{bmatrix} -7 & -3 \\ 2 & 0 \end{bmatrix} \dots(1)$$

$$A^{-1} = \frac{1}{(0+3)} \begin{bmatrix} 4 & 3 \\ -1 & 0 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 4 & 3 \\ -1 & 0 \end{bmatrix}$$

$$B^{-1} = \frac{1}{(2-0)} \begin{bmatrix} -1 & 3 \\ 0 & -2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} -1 & 3 \\ 0 & -2 \end{bmatrix}$$

$$B^{-1}A^{-1} = \frac{1}{2} \begin{bmatrix} -1 & 3 \\ 0 & -2 \end{bmatrix} \frac{1}{3} \begin{bmatrix} 4 & 3 \\ -1 & 0 \end{bmatrix}$$

$$= \frac{1}{6} \begin{bmatrix} -4-3 & -3+0 \\ 0+2 & 0+0 \end{bmatrix} = \frac{1}{6} \begin{bmatrix} -7 & -3 \\ 2 & 0 \end{bmatrix} \dots(2)$$

As the matrices in (1) and (2) are same,
 $(AB)^{-1} = B^{-1}A^{-1}$ is verified.

2. Solve the following system of linear equations, using matrix inversion method : $5x + 2y = 3$, $3x + 2y = 5$. [May - 2022]

Sol. The matrix form of the system is $AX = B$, where

$$A = \begin{bmatrix} 5 & 2 \\ 3 & 2 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 3 \\ 5 \end{bmatrix}$$

$$\text{We find } |A| = \begin{vmatrix} 5 & 2 \\ 3 & 2 \end{vmatrix} = 10 - 6 = 4 \neq 0. \text{ So } A^{-1}$$

$$\text{exists and } A^{-1} = \frac{1}{4} \begin{bmatrix} 2 & -2 \\ -3 & 5 \end{bmatrix}$$

Then, applying the formula $X = A^{-1} B$, we get

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 2 & -2 \\ -3 & 5 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 6 & -10 \\ -9 & +25 \end{bmatrix}$$

$$= \frac{1}{4} \begin{bmatrix} -4 \\ -4 \\ 16 \\ 4 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \\ 4 \\ 1 \end{bmatrix}$$

So the solution is $x = -1, y = 4$.

3. If the rank of the matrix

$$\begin{bmatrix} \lambda & -1 & 0 \\ 0 & \lambda & -1 \\ -1 & 0 & \lambda \end{bmatrix} \text{ is 2, then find } \lambda. \quad [\text{Hy. - 2023}]$$

Sol. Given rank of $\begin{bmatrix} \lambda & -1 & 0 \\ 0 & \lambda & -1 \\ -1 & 0 & \lambda \end{bmatrix}$ is 2

⇒ The value of the third order determinant is zero

$$\Rightarrow \begin{vmatrix} \lambda & -1 & 0 \\ 0 & \lambda & -1 \\ -1 & 0 & \lambda \end{vmatrix} = 0$$

$$\Rightarrow \lambda \begin{vmatrix} \lambda & -1 \\ 0 & \lambda \end{vmatrix} + 1 \begin{vmatrix} 0 & -1 \\ -1 & \lambda \end{vmatrix} + 0 = 0$$

$$\Rightarrow \lambda(\lambda^2 - 0) + 1(0 - 1) = 0$$

$$\Rightarrow \lambda^3 - 1 = 0$$

$$\Rightarrow \lambda^3 = 1$$

$$\Rightarrow \lambda = 1$$

$$\therefore \lambda = 1$$

4. Find the inverse of the matrix $\begin{bmatrix} 2 & -1 & 3 \\ -5 & 3 & 1 \\ -3 & 2 & 3 \end{bmatrix}$.

Sol. Refer Text Book Example 1.3 [FRT - 2022]

5. If $A = \begin{pmatrix} 5 & 2 \\ 7 & 3 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix}$ then verify

$$(AB)^{-1} = B^{-1} A^{-1}. \quad [\text{FRT} - 2022]$$

Sol.

$$A = \begin{pmatrix} 5 & 2 \\ 7 & 3 \end{pmatrix} \quad B = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix}$$

$$AB = \begin{pmatrix} 5 & 2 \\ 7 & 3 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 10-2 & -5+2 \\ 14-3 & -7+3 \end{pmatrix} = \begin{pmatrix} 8 & -3 \\ 11 & -4 \end{pmatrix}$$

$$|AB| = \begin{vmatrix} 8 & -3 \\ 11 & -4 \end{vmatrix} = -32 + 33 = 1$$

$$(AB)^{-1} = \frac{1}{|AB|} \text{adj } AB = \frac{1}{1} \begin{pmatrix} -4 & 3 \\ -11 & 8 \end{pmatrix} \quad \dots (1)$$

$$B^{-1} = \frac{1}{|B|} \text{adj } B = \frac{1}{(2-1)} \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$A^{-1} = \frac{1}{15-14} \begin{pmatrix} 3 & -2 \\ -7 & 5 \end{pmatrix} = \begin{pmatrix} 3 & -2 \\ -7 & 5 \end{pmatrix}$$

$$B^{-1} A^{-1} = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 3 & -2 \\ -7 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} 3-7 & -2+5 \\ 3-14 & -2+10 \end{bmatrix} = \begin{bmatrix} -4 & 3 \\ -11 & 8 \end{bmatrix} \quad \dots (2)$$

From (1) and (2) $(AB)^{-1} = B^{-1} A^{-1}$.

Hence proved.

6. If $A = \begin{bmatrix} 2 & -1 & 3 \\ -5 & 3 & 1 \\ -3 & 2 & 3 \end{bmatrix}$, then find $|\text{adj}(\text{adj } A)|$. [Mar. - 2024]

Sol. Given $\text{adj } A = \begin{bmatrix} 2 & -1 & 3 \\ -5 & 3 & 1 \\ -3 & 2 & 3 \end{bmatrix}$.

We know that $|\text{adj}(\text{adj } A)| = |A|^{(n-1)^2}$

Here $n = 3$, $|\text{adj}(\text{adj } A)| = |A|^{(3-1)^2} = |A|^4$

$$|A| = \begin{vmatrix} 2 & -1 & 3 \\ -5 & 3 & 1 \\ -3 & 2 & 3 \end{vmatrix}$$

$$= 2(9-2) + 1(-15+3) + 3(-10+9)$$

$$= 2(7) + 1(-12) + 3(-1) = 14 - 12 - 3$$

$$= 14 - 15 = -1$$

$$|\text{adj}(\text{adj } A)| = -1$$

5 MARKS

1. By using Gaussian elimination method, balance the chemical reaction equation:



Sol. We are searching for positive integers x_1, x_2, x_3 and x_4 such that

$$x_1 \text{C}_5\text{H}_8 + x_2 \text{O}_2 = x_3 \text{CO}_2 + x_4 \text{H}_2\text{O} \quad \dots (1)$$

The number of carbon atoms on the left-hand side of (1) should be equal to the number of carbon atoms on the right-hand side of (1). So we get a linear homogeneous equation

$$5x_1 = x_3$$

$$\Rightarrow 5x_1 - x_3 = 0 \quad \dots (2)$$

Similarly, considering hydrogen and oxygen atoms, we get respectively,

$$8x_1 = 2x_4$$

$$\Rightarrow 4x_1 - x_4 = 0, \quad \dots (3)$$

$$2x_2 = 2x_3 + x_4$$

$$\Rightarrow 2x_2 - 2x_3 - x_4 = 0. \quad \dots (4)$$

Equations (2), (3), and (4) constitute a homogeneous system of linear equations in four unknowns.

The augmented matrix is $[A|B]$

$$= \left[\begin{array}{cccc|c} 5 & 0 & -1 & 0 & 0 \\ 4 & 0 & 0 & -1 & 0 \\ 0 & 2 & -2 & -1 & 0 \end{array} \right]$$

By Gaussian elimination method, we get

$$[A|B] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{cccc|c} 4 & 0 & 0 & -1 & 0 \\ 5 & 0 & -1 & 0 & 0 \\ 0 & 2 & -2 & -1 & 0 \end{array} \right]$$

$$\xrightarrow{R_2 \leftrightarrow R_3} \left[\begin{array}{cccc|c} 4 & 0 & 0 & -1 & 0 \\ 0 & 2 & -2 & -1 & 0 \\ 5 & 0 & -1 & 0 & 0 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow 4R_3 - 5R_1} \left[\begin{array}{cccc|c} 4 & 0 & 0 & -1 & 0 \\ 0 & 2 & -2 & -1 & 0 \\ 0 & 0 & -4 & 5 & 0 \end{array} \right]$$

Therefore, $\rho(A) = \rho([A|B]) = 3 < 4 =$ Number of unknowns.

The system is consistent and has infinite number of solutions.

Writing the equations using the echelon form, we get $4x_1 - x_4 = 0$, $2x_2 - 2x_3 - x_4 = 0$, $-4x + 5x_4 = 0$.


So, one of the unknowns should be chosen arbitrarily as a non-zero real number.

Let us choose $x_4 = t$. Then, by back substitution, we $x_3 = \frac{5t}{4}$, $x_2 = \frac{7t}{4}$, $x_1 = \frac{t}{4}$

Since x_1, x_2, x_3 and x_4 are positive integers, let us choose $t = 4$.

Then, we get $x_1 = 1, x_2 = 7, x_3 = 5$ and $x_4 = 4$.

So, the balanced equation is



2. Test the consistency of the following system of linear equations by rank method.

$$x - y + z = -9; \quad 2x - y + z = 4$$

$$3x - y + z = 6; \quad 4x - y + 2z = 7 \quad [\text{Mar. - 2020}]$$

Sol. $[A/B] = \left[\begin{array}{ccc|c} 1 & -1 & 1 & -9 \\ 2 & -1 & 1 & 4 \\ 3 & -1 & 1 & 6 \\ 4 & -1 & 2 & 7 \end{array} \right]$

$$\sim \left[\begin{array}{ccc|c} 1 & -1 & 1 & -9 \\ 0 & 1 & -1 & 22 \\ 0 & 2 & -2 & 33 \\ 0 & 3 & -2 & 43 \end{array} \right] \begin{array}{l} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 3R_1 \\ R_4 \rightarrow R_4 - 4R_1 \end{array}$$

$$\sim \left[\begin{array}{ccc|c} 1 & -1 & 1 & -9 \\ 0 & 1 & -1 & 22 \\ 0 & 0 & 0 & -11 \\ 0 & 0 & 1 & -23 \end{array} \right] \begin{array}{l} R_3 \rightarrow R_3 - 2R_2 \\ R_4 \rightarrow R_4 - 3R_2 \end{array}$$

$$R_3 \leftrightarrow R_4$$

$\rho(A) \neq \rho(A/B)$ Inconsistent and no solution.

3. Solve the system of equations $x - y + 2z = 2$, $2x + y + 4z = 7$, $4x - y + z = 4$ by Cramer's rule.

Sol. First we evaluate the determinants [Aug. - 2021]

$$\Delta = \begin{vmatrix} 1 & -1 & 2 \\ 2 & 1 & 4 \\ 4 & -1 & 1 \end{vmatrix}$$

$$= 1(1 + 4) + 1(2 - 16) + 2(-2 - 4)$$

$$= 1(5) + 1(-14) + 2(-6) = 5 - 14 - 12$$

$$\Delta = -21$$

$$\Delta_1 = \begin{vmatrix} 2 & -1 & 2 \\ 7 & 1 & 4 \\ 4 & -1 & 1 \end{vmatrix}$$

$$= 2(1 + 4) + 1(7 - 16) + 2(-7 - 4)$$

$$= 2(5) + 1(-9) + 2(-11) = 10 - 9 - 22$$

$$\Delta_1 = -21$$

$$\Delta_2 = \begin{vmatrix} 1 & 2 & 2 \\ 2 & 7 & 4 \\ 4 & 4 & 1 \end{vmatrix}$$

$$= 1(7 - 16) - 2(2 - 16) + 2(8 - 28)$$

$$= 1(-9) - 2(-14) + 2(-20) = -9 + 28 - 40$$

$$\Delta_2 = -21$$

$$\Delta_3 = \begin{vmatrix} 1 & -1 & 2 \\ 2 & 1 & 7 \\ 4 & -1 & 4 \end{vmatrix}$$

$$= 1(4 + 7) + 1(8 - 28) + 2(-2 - 4)$$

$$= 1(11) + 1(-20) + 2(-6) = 11 - 20 - 12$$

$$\Delta_3 = -21$$

$$x = \frac{\Delta_1}{\Delta} = \frac{-21}{-21} = 1$$

$$y = \frac{\Delta_2}{\Delta} = \frac{-21}{-21} = 1$$

$$z = \frac{\Delta_3}{\Delta} = \frac{-21}{-21} = 1 \Rightarrow x = 1, y = 1, z = 1$$

4. Cramer's rule is not applicable to solve the system $3x + y + z = 2$, $x - 3y + 2z = 1$, $7x - y + 4z = 5$. Why? [May - 2022]

Sol. When the coefficient matrix is a square matrix and non-singular, then the matrix can be solved by Cramer's rule.

Coefficient matrix

$$\begin{vmatrix} 3 & 1 & 1 \\ 1 & -3 & 2 \\ 7 & -1 & 4 \end{vmatrix} = 3(-12 + 2) - 1(4 - 14) + 1(-1 + 21)$$

$$= 3(-10) - 1(-10) + 1(20)$$

$$= -30 + 10 + 20 = 0$$

$\therefore$ Coefficient matrix is singular

$\therefore$ Cramer's rule is not applicable.

5. If $A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$, verify that $A (\text{adj } A)$

$$= (\text{adj } A) A = |A| I_3, \quad [\text{FRT - 2022}]$$

Sol. Refer Text Book Example 1.1

6. Solve the system of linear equations by Cramer's Rule. $x_1 - x_2 = 3$, $2x_1 + 3x_2 + 4x_3 = 17$, $x_2 + 2x_3 = 7$. [June - 2023]

Sol. First we evaluate the determinants

$$\Delta = \begin{vmatrix} 1 & -1 & 0 \\ 2 & 3 & 4 \\ 0 & 1 & 2 \end{vmatrix} = 6 \neq 0,$$

$$\Delta_1 = \begin{vmatrix} 3 & -1 & 0 \\ 17 & 3 & 4 \\ 7 & 1 & 2 \end{vmatrix} = 12,$$

$$\Delta_2 = \begin{vmatrix} 1 & 3 & 0 \\ 2 & 17 & 4 \\ 0 & 7 & 2 \end{vmatrix} = -6,$$

$$\Delta_3 = \begin{vmatrix} 1 & -1 & 3 \\ 2 & 3 & 17 \\ 0 & 1 & 7 \end{vmatrix} = 24$$

By Cramer's rule, we get $x_1 = \frac{\Delta_1}{\Delta} = \frac{12}{6} = 2$,

$$x_2 = \frac{\Delta_2}{\Delta} = \frac{-6}{6} = -1,$$

$$x_3 = \frac{\Delta_3}{\Delta} = \frac{24}{6} = 4.$$

So, the solution is $(x_1 = 2, x_2 = -1, x_3 = 4)$.

7. If the system of equations $px + by + cz = 0$, $ax + qy + cz = 0$, $ax + by + rz = 9$ has a non-trivial solution and $p \neq a, q \neq b, r \neq c$, prove that

$$\frac{p}{p-a} + \frac{q}{q-b} + \frac{r}{r-c} = 2 \quad [\text{Hy. - 2023}]$$

Sol. Refer Text Book Example 1.40

ADDITIONAL QUESTION & ANSWERS

1 MARK

1. Choose the Correct or the most suitable answer from the given four alternatives :

1. If the system of equations $x = cy + bz$, $y = az + cx$ and $z = bx + ay$ has a non-trivial solution then

- (1) $a^2 + b^2 + c^2 = 1$ (2) $abc \neq 1$
(3) $a + b + c = 0$ (4) $a^2 + b^2 + c^2 + 2abc = 1$

[Ans: (4) $a^2 + b^2 + c^2 + 2abc = 1$]

Hint : $\begin{vmatrix} 1 & -c & -b \\ c & -1 & a \\ b & a & -1 \end{vmatrix} = 0$

$$\Rightarrow 1[1 - a^2] + c[-c - ab] - b[ac + b] = 0$$

$$\Rightarrow 1 - a^2 - c^2 - abc - abc - b^2 = 0$$

2. Let A be a 3×3 matrix and B its adjoint matrix. If $|B| = 64$, then $|A| =$

- (1) ± 2 (2) ± 4 (3) ± 8 (4) ± 12

Hint : $A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{|A|} B$ [Ans: (3) ± 8]

$$\Rightarrow |A| |A^{-1}| = |B| \Rightarrow |A|^3 |A^{-1}| = |B| = 64$$

$$\Rightarrow |A|^2 = 64 \Rightarrow |A| = \pm 8$$

3. If A^T is the transpose of a square matrix A, then

- (1) $|A| \neq |A^T|$ (2) $|A| = |A^T|$
(3) $|A| + |A^T| = 0$
(4) $|A| = |A^T|$ only when A is symmetric

[Ans: (2) $|A| = |A^T|$]

4. If A is a square matrix that $|A| = 2$, then for any positive integer n, $|A^n| =$

- (1) 0 (2) $2n$ (3) 2^n (4) n^2

[Ans: (3) 2^n]

Hint : $|A^n| = |A|^n = |A| |A| \dots \text{times}$
 $= (2)(2) \dots \text{times} = 2^n$

5. If A is a square matrix of order n, then $|\text{adj } A| =$

- (1) $|A|^{n-1}$ (2) $|A|^{n-2}$ (3) $|A|^n$ (4) None

[Ans: (1) $|A|^{n-1}$]

Hint : $\text{adj } A = |A| A^{-1}$

$$\Rightarrow |\text{adj } A| = |A|^n |A^{-1}| = |A|^{n-1}$$

6. If $A = \begin{pmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{pmatrix}$ and $A (\text{adj } A) = \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ then λ is

- (1) $\sin x \cos x$ (2) 1
(3) 2 (4) none [Ans: (2) 1]

Hint : $A (\text{adj } A) = \begin{pmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{pmatrix} \begin{pmatrix} \cos x & -\sin x \\ \sin x & \cos x \end{pmatrix}$
 $= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

7. If A is a matrix of order $m \times n$, then $\rho(A)$ is

- (1) m (2) n
(3) $\leq \min(m, n)$ (4) $\geq \min(m, n)$
[Ans: (3) $\leq \min(m, n)$]

8. If $\rho(A) = r$ then which of the following is correct?

- (1) all the minors of order n which do not vanish
(2) 'A' has at least one minor of order r which does not vanish and all higher order minors vanish
(3) 'A' has at least one $(r + 1)$ order minor which vanish
(4) all $(r + 1)$ and higher order minors should not vanish

[Ans: (2) 'A' has at least one minor of order r which does not vanish and all higher order minors vanish]

9. Which of the following is not an elementary transformation?

- (1) $R_i \leftrightarrow R_j$ (2) $R_i \rightarrow 2R_i + R_j$
(3) $C_j \rightarrow C_j + C_i$ (4) $R_i \rightarrow R_i + C_j$

[Ans: (4) $R_i \rightarrow R_i + C_j$]

II. Fill in the blanks :

1. Every homogeneous system.....

- (1) Is always consistent
(2) Has only trivial solution
(3) Has infinitely many solution
(4) Need not be consistent

[Ans: (1) Is always consistent]

2. If $\rho(A) \neq \rho([A|B])$, then the system is.....

- (1) consistent and has infinitely many solutions
(2) consistent and has a unique solution
(3) consistent
(4) inconsistent [Ans: (4) inconsistent]

3. In the non – homogeneous system of equations with 3 unknowns if $\rho(A) = \rho([A|B]) = 2$, then the system has.....

- (1) unique solution
(2) one parameter family of solution
(3) two parameter family of solutions
(4) inconsistent
[Ans: (2) one parameter family of solution]

4. In a homogeneous system if $\rho(A) = \rho([A|0]) <$ the number of unknowns then the system has.....

- (1) trivial solution
(2) only non – trivial solution
(3) no solution
(4) trivial solution and infinitely many non – trivial solutions
[Ans: (4) trivial solution and infinitely many non-trivial solutions]

5. In the system of equations with 3 unknowns, if $\Delta = 0$, and one of $\Delta_x, \Delta_y, \Delta_z$ is non zero then the system is.....

- (1) Consistent (2) inconsistent
(3) consistent with one parameter family of solutions
(4) consistent with two parameter family of solutions [Ans: (2) inconsistent]

6. In the system of linear equations with 3 unknowns if $\rho(A) = \rho([A|B]) = 1$, the system has.....

- (1) unique solution (2) inconsistent
(3) consistent with 2 parameter family of solution
(4) consistent with one parameter family of solution.
[Ans: (3) consistent with 2 parameter family of solution]

7. Cramer's rule is applicable only when.....

- (1) $\Delta \neq 0$ (2) $\Delta = 0$
 (3) $\Delta = 0, \Delta_x = 0$ (4) $\Delta_x = \Delta_y = \Delta_z = 0$

[Ans: (1) $\Delta \neq 0$]**8. If $A = [2 \ 0 \ 1]$ then the rank of AA^T is**

- (1) 1 (2) 2 (3) 3 (4) 0

[Ans: (1) 1]

Hint : $AA^T = [2 \ 0 \ 1] \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix} = [5]$
 $P(AA^T) = 1$

9. In a square matrix the minor M_{ij} and the co-factor A_{ij} of and element a_{ij} are related by

- (1) $A_{ij} = -M_{ij}$ (2) $A_{ij} = M_{ij}$
 (3) $A_{ij} = (-1)^{i+j} M_{ij}$ (4) $A_{ij} = (-1)^{i-j} M_{ij}$

[Ans: (3) $A_{ij} = (-1)^{i+j} m_{ij}$]**III. Match the following :****1. If A is a non - singular matrix of order n , then**

	List - I		List - II
i.	$ \text{adj } A $	a)	$ A ^{(n-1)^2}$
ii.	$(\text{adj } A)^T$	b)	$ A ^{n-1}$
iii.	$\text{adj } (\text{adj } A)$	c)	$\text{adj } (A^T)$
iv.	$ \text{adj } (\text{adj } A) $	d)	$ A ^{n-2} \cdot A$

The Correct match is

- (i) (ii) (iii) (iv)
 (1) a b c d
 (2) c d a b
 (3) d a b c
 (4) b c d a

[Ans : (4) i - b ii - c iii - d iv - a]**2. If A is a non - singular matrix of order n , then**

	List - I		List - II
i.	$(\text{adj } A)^{-1}$	a)	$(\text{adj } B) (\text{adj } A)$
ii.	$(\lambda A)^{-1}$	b)	$\lambda^{n-1} \text{adj } (A)$
iii.	$\text{adj } (\lambda A)$	c)	$\text{adj } (A^{-1})$
iv.	$\text{adj } (AB)$	d)	$\frac{1}{\lambda} A^{-1}$

The Correct match is

- (i) (ii) (iii) (iv)
 (1) c d b a
 (2) a b c d
 (3) b c d a
 (4) c a b d

[Ans : (1) i - c ii - d iii - b 4 - a]**3.**

	List - I		List - II
i.	$(A^T)^{-1}$	(a)	A
ii.	$A (\text{adj } A)$	(b)	$(A^{-1})^T$
iii.	$(AB)^{-1}$	(c)	$ A \cdot I_n$
iv.	$(A^{-1})^{-1}$	(d)	$B^{-1} A^{-1}$

The Correct match is

- (i) (ii) (iii) (iv)
 (1) b c d a
 (2) a b c d
 (3) c d a b
 (4) b a c d

[Ans : (1) i - b ii - c iii - d iv - a]**IV. Choose the odd man out :****1. The rank of any 3×4 matrix is**

- (1) May be 1 (2) May be 2
 (3) May be 3 (4) May be 4

[Ans: (4) May be 4]**Hint:** $[\rho(A) \leq \min(3, 4) \Rightarrow \rho(A) \leq 3 \text{ and it cannot be 4}]$ **2. If A is symmetric then**

- (1) $A^T = A$
 (2) $\text{adj } A$ is symmetric
 (3) $\text{adj } (A^T) = (\text{adj } A)^T$
 (4) A is orthogonal

Hint: [1, 2, 3 are properties of symmetric matrix]**[Ans: (4) A is orthogonal]****3. If A is a non-singular matrix of odd order then**

- (1) Order of A is $2m + 1$
 (2) Order of A is $2m + 2$
 (3) $|\text{adj } A|$ is positive (4) $|A| \neq 0$

[Ans: (2) Order of A is $2m + 2$]**Hint:** $(2m + 2)$ is even

4. A matrix which is obtained from an identity matrix by applying only one elementary transformation is

- (1) Identity matrix
- (2) Elementary matrix
- (3) Square matrix
- (4) Equivalent to identity matrix

[Ans: (1) Identity matrix]

V. Choose the incorrect answer :

1. In an echelon form which of the following is incorrect ?

- (1) Every row of A which has all its entries 0 occurs below every row which has a non-zero entry.
- (2) The first non-zero entry in each non-zero row is 1
- (3) The number of zeros before the first non-zero element in a row is less than the number of such zeros in the next row
- (4) Two row can have same number of zeros before the first non-zero entry

[Ans: (4) Two row can have same number of zeros before the first non-zero entry]

2. If A is an invertible matrix, then which of the following is not true.

- (1) $(A^2)^{-1} = (A^{-1})^2$
- (2) $|A^{-1}| = |A|^{-1}$
- (3) $(A^T)^{-1} = (A^{-1})^T$
- (4) $|A| \neq 0$

[Ans: (1) $(A^2)^{-1} = (A^{-1})^2$]

3. The matrix $\begin{bmatrix} 5 & 10 & 3 \\ -2 & -4 & 6 \\ -1 & -2 & x \end{bmatrix}$ is a singular matrix if the value of x is

- (1) 3
- (2) non- existent
- (3) All values of x
- (4) None

[Ans: (3) All values of x]

2 MARKS

1. Find the inverse of the non-singular matrix

$$A = \begin{bmatrix} 0 & 5 \\ -1 & 6 \end{bmatrix}, \text{ by Gauss Jordan method.}$$

Sol. Applying Gauss – Jordan method, we get

$$[A|I_2] = \left[\begin{array}{cc|cc} 0 & 5 & 1 & 0 \\ -1 & 6 & 0 & 1 \end{array} \right] \xrightarrow{R_1 \rightarrow R_2} \left[\begin{array}{cc|cc} -1 & 6 & 0 & 1 \\ 0 & 5 & 1 & 0 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow (-1)R_1} \left[\begin{array}{cc|cc} 1 & -6 & 0 & -1 \\ 0 & 5 & 1 & 0 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow \frac{1}{5} \times R_2} \left[\begin{array}{cc|cc} 1 & -6 & 0 & -1 \\ 0 & 1 & \frac{1}{5} & 2 \end{array} \right]$$

$$\xrightarrow{R_1 \rightarrow R_1 + 6R_2} \left[\begin{array}{cc|cc} 1 & 0 & \frac{6}{5} & -1 \\ 0 & 1 & \frac{1}{5} & 2 \end{array} \right]$$

$$\text{So, we get } A^{-1} = \xrightarrow{R_1 \rightarrow R_1 + 6R_2} \left[\begin{array}{cc|cc} \frac{6}{5} & -1 & & \\ \frac{1}{5} & 2 & & \end{array} \right] = \frac{1}{5} \begin{bmatrix} 6 & -5 \\ 1 & 0 \end{bmatrix}$$

2. For any 2×2 matrix, if $A (\text{adj } A) = \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix}$ then find $|A|$.

Sol. Given $A (\text{adj } A) = \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix} = 10 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \dots (1)$

We know $A (\text{adj } A) = (\text{adj } A) A = |A| \cdot I_2 \dots (2)$
Comparing (1) and (2), we get $|A| = 10$.

3. Show that the equations $3x + y + 9z = 0$, $3x + 2y + 12z = 0$ and $2x + y + 7z = 0$ have non-trivial solutions also.

Sol. The matrix form of the system is

$$\begin{bmatrix} 3 & 1 & 9 \\ 3 & 2 & 12 \\ 2 & 1 & 7 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$AX = B$ where

$$A = \begin{bmatrix} 3 & 1 & 9 \\ 3 & 2 & 12 \\ 2 & 1 & 7 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{aligned} |A| &= \begin{vmatrix} 3 & 1 & 9 \\ 3 & 2 & 12 \\ 2 & 1 & 7 \end{vmatrix} = 3 \begin{vmatrix} 2 & 12 \\ 1 & 7 \end{vmatrix} - 1 \begin{vmatrix} 3 & 12 \\ 2 & 7 \end{vmatrix} + 9 \begin{vmatrix} 3 & 2 \\ 2 & 1 \end{vmatrix} \\ &= 3(14 - 12) - 1(21 - 24) + 9(3 - 4) \\ &= 3(2) - 1(-3) + 9(-1) \\ &= 6 + 3 - 9 = 9 - 9 = 0 \end{aligned}$$

Since $|A| = 0$, the homogeneous system of equations have non-trivial solutions also.

4. For the matrix A, if $A^3 = I$, then find A^{-1} .

Sol. Given $A^3 = I$

Pre multiply by A^{-1} we get,

$$A^{-1} \cdot A^3 = A^{-1} \cdot I$$

$$\begin{aligned}\Rightarrow (A^{-1} \cdot A) A^2 &= A^{-1} & [\because A^{-1} I = A^{-1}] \\ \Rightarrow I \cdot A^2 &= A^{-1} & [\because A^{-1} A = I] \\ \Rightarrow A^2 &= A^{-1} & [\because I \cdot A^2 = A^2] \\ \therefore A^{-1} &= A^2\end{aligned}$$

5. Show that the system of equations is inconsistent. $2x + 5y = 7$, $6x + 15y = 13$.

Sol. Augmented matrix

$$[A|B] = \left[\begin{array}{cc|c} 2 & 5 & 7 \\ 6 & 15 & 13 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 - 3R_1} \left[\begin{array}{cc|c} 2 & 5 & 7 \\ 0 & 0 & -8 \end{array} \right]$$

Here $\rho(A) = 1$ and $\rho([A|B]) = 2$

$\therefore \rho(A) \neq \rho([A|B])$

Hence the system is inconsistent,

6. Find k if the equations $x + 2y + 2z = 0$, $x - 3y - 3z = 0$, $2x + y + kz = 0$ have only the trivial solution.

Sol. Matrix form of the given system of equations is

$$\begin{bmatrix} 1 & 2 & 2 \\ 1 & -3 & -3 \\ 2 & 1 & k \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$AX = B \text{ where } A = \begin{bmatrix} 1 & 2 & 2 \\ 1 & -3 & -3 \\ 2 & 1 & k \end{bmatrix}$$

Homogeneous system of equations has trivial solution only if $|A| \neq 0$.

$$\therefore \begin{vmatrix} 1 & 2 & 2 \\ 1 & -3 & -3 \\ 2 & 1 & k \end{vmatrix} \neq 0.$$

Expanding along R_1 ,

$$1 \begin{vmatrix} -3 & -3 \\ 1 & k \end{vmatrix} - 2 \begin{vmatrix} 1 & -3 \\ 2 & k \end{vmatrix} + 2 \begin{vmatrix} 1 & -3 \\ 2 & 1 \end{vmatrix} \neq 0.$$

$$\Rightarrow 1(-3k + 3) - 2(k + 6) + 2(1 + 6) \neq 0$$

$$\Rightarrow -3k + 3 - 2k - 12 + 14 \neq 0$$

$$\Rightarrow -5k + 5 \neq 0 \Rightarrow -5k \neq -5 \Rightarrow k \neq \frac{-5}{-5} = 1$$

$$\Rightarrow k \neq 1$$

3 MARKS

1. For what value of t will the system $tx + 3y - z = 1$, $x + 2y + z = 2$, $-tx + y + 2z = -1$ fail to have unique solution?

Sol. $\Delta = \begin{vmatrix} t & 3 & -1 \\ 1 & 2 & 1 \\ -t & 1 & 2 \end{vmatrix} = t \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} - 3 \begin{vmatrix} 1 & 1 \\ -t & 2 \end{vmatrix} - \begin{vmatrix} 1 & 2 \\ -t & 1 \end{vmatrix}$

$$= t(4 - 1) - 3(2 + t) - 1(1 + 2t)$$

$$= 3t - 6 - 3t - 1 - 2t = -7 - 2t$$

The system will fail to have unique solution if

$$\Delta = 0 \Rightarrow -7 - 2t = 0 \Rightarrow -2t = 7 \Rightarrow t = \frac{-7}{2}$$

$$\therefore t = \frac{-7}{2}.$$

2. Under what conditions will the rank of the

matrix $\begin{bmatrix} 1 & 0 & 0 \\ 0 & h-2 & 2 \\ 0 & 0 & h+2 \\ 0 & 0 & 3 \end{bmatrix}$ be less than 3?

Sol. Let $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & h-2 & 2 \\ 0 & 0 & h+2 \\ 0 & 0 & 3 \end{bmatrix}$

The rank of A will be less than 3 if every minor of order 3 vanishes

$$\therefore \begin{vmatrix} 1 & 0 & 0 \\ 0 & h-2 & 0 \\ 0 & 0 & 3 \end{vmatrix} = 0$$

$$\Rightarrow 1 \begin{vmatrix} h-2 & 0 \\ 0 & 3 \end{vmatrix} + 0 + 0 = 0 \Rightarrow 3(h-2) = 0$$

$$\Rightarrow h-2 = 0 \Rightarrow h = 2.$$

5 MARKS

1. Using determinants, find the quadratic defined by $f(x) = ax^2 + bx + c$, if $f(1) = 0$, $f(2) = -2$ and $f(3) = -6$.

Sol.

$$\text{Given } f(x) = ax^2 + bx + c$$

$$f(1) = 0 \Rightarrow a(1)^2 + b(1) + c = 0$$

$$\Rightarrow a + b + c = 0 \quad \dots(1)$$

$$f(2) = -2 \Rightarrow a(2^2) + b(2) + c = -2$$

$$\Rightarrow 4a + 2b + c = -2 \quad \dots(2)$$

$$f(3) = -6$$

$$\Rightarrow a(3^2) + b(3) + c = -6$$

$$\Rightarrow 9a + 3b + c = -6 \quad \dots(3)$$

$$\Delta = \begin{vmatrix} 1 & 1 & 1 \\ 4 & 2 & 1 \\ 9 & 3 & 1 \end{vmatrix} = 1 \begin{vmatrix} 2 & 1 \\ 3 & 1 \end{vmatrix} - 1 \begin{vmatrix} 4 & 1 \\ 9 & 1 \end{vmatrix} + 1 \begin{vmatrix} 4 & 2 \\ 9 & 3 \end{vmatrix}$$

$$= 1(2 - 3) - 1(4 - 9) + 1(12 - 18)$$

$$= -1 + 5 - 6 = -2 \neq 0$$

$$\begin{aligned}\Delta_1 &= \begin{vmatrix} 0 & 1 & 1 \\ -2 & 2 & 1 \\ -6 & 3 & 1 \end{vmatrix} = 0 - 1 \begin{vmatrix} -2 & 1 \\ -6 & 1 \end{vmatrix} + 1 \begin{vmatrix} -2 & 2 \\ -6 & 3 \end{vmatrix} \\ &= -1(-2+6) + 1(-6+12) \\ &= -1(4) + 1(6) = 2 \\ \Delta_2 &= \begin{vmatrix} 1 & 0 & 1 \\ 4 & -2 & 1 \\ 9 & -6 & 1 \end{vmatrix} = 1 \begin{vmatrix} -2 & 1 \\ -6 & 1 \end{vmatrix} + 0 + 1 \begin{vmatrix} 4 & -2 \\ 9 & -6 \end{vmatrix} \\ &= +1(-2+6) + 1(-24+18) = 4-6 = -2 \\ \Delta_3 &= \begin{vmatrix} 1 & 1 & 0 \\ 4 & 2 & -2 \\ 9 & 3 & -6 \end{vmatrix} = 1 \begin{vmatrix} 2 & -2 \\ 3 & -6 \end{vmatrix} - 1 \begin{vmatrix} 4 & -2 \\ 9 & -6 \end{vmatrix} \\ &= 1(-12+6) - 1(-24+18) = -6+6 = 0\end{aligned}$$

∴ By Cramer's rule,

$$a = \frac{\Delta_1}{\Delta} = \frac{2}{-2} = -1$$

$$b = \frac{\Delta_2}{\Delta} = \frac{-2}{-2} = 1$$

$$c = \frac{\Delta_3}{\Delta} = \frac{0}{-2} = 0$$

$$\therefore f(x) = (-1)x^2 + 1x + 0$$

$$\Rightarrow f(x) = -x^2 + x$$

2. The sum of three numbers is 20. If we multiply the third number by 2 and add the first number to the result we get 23. By adding second and third numbers to 3 times the first number we get 46. Find the numbers using Cramer's rule.

Sol. Let the required numbers be x, y and z

By the given data,

$$x + y + z = 20 \quad \dots(1)$$

$$2z + x = 23 \quad \Rightarrow x + 2z = 23 \quad \dots(2)$$

$$y + z + 3x = 46 \quad \Rightarrow 3x + y + z = 46 \quad \dots(3)$$

$$\Delta = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 0 & 2 \\ 3 & 1 & 1 \end{vmatrix} = 1 \begin{vmatrix} 0 & 2 \\ 1 & 1 \end{vmatrix} - 1 \begin{vmatrix} 1 & 2 \\ 3 & 1 \end{vmatrix} + 1 \begin{vmatrix} 1 & 0 \\ 3 & 1 \end{vmatrix} \\ = -2 + 5 + 1 = 4$$

$$\Delta_1 = \begin{vmatrix} 20 & 1 & 1 \\ 23 & 0 & 2 \\ 46 & 1 & 1 \end{vmatrix} = 20 \begin{vmatrix} 0 & 2 \\ 1 & 1 \end{vmatrix} - 1 \begin{vmatrix} 23 & 2 \\ 46 & 1 \end{vmatrix} + 1 \begin{vmatrix} 23 & 0 \\ 46 & 1 \end{vmatrix} \\ = -40 + 69 + 23 = 52$$

$$\Delta_2 = \begin{vmatrix} 1 & 20 & 1 \\ 1 & 23 & 2 \\ 3 & 46 & 1 \end{vmatrix} = 1 \begin{vmatrix} 23 & 2 \\ 46 & 1 \end{vmatrix} - 20 \begin{vmatrix} 1 & 2 \\ 3 & 1 \end{vmatrix} + 1 \begin{vmatrix} 1 & 23 \\ 3 & 46 \end{vmatrix} \\ = -69 + 100 - 23 = 8$$

$$\Delta_3 = \begin{vmatrix} 1 & 1 & 20 \\ 1 & 0 & 23 \\ 3 & 1 & 46 \end{vmatrix} = 1 \begin{vmatrix} 0 & 23 \\ 1 & 46 \end{vmatrix} - 1 \begin{vmatrix} 1 & 23 \\ 3 & 46 \end{vmatrix} + 20 \begin{vmatrix} 1 & 0 \\ 3 & 1 \end{vmatrix} \\ = -23 + 23 + 20 = 20$$

$$\therefore x = \frac{\Delta_1}{\Delta} = \frac{52}{4} = 13$$

$$y = \frac{\Delta_2}{\Delta} = \frac{8}{4} = 2 \text{ and } z = \frac{\Delta_3}{\Delta} = \frac{20}{4} = 5.$$

Hence the required numbers are 13, 2 and 5.

3. Show that the equations $-2x + y + z = a$, $x - 2y + z = b$, $x + y - 2z = c$ are consistent only if $a + b + c = 0$.

Sol. Augmented matrix $[A|B]$ is $\begin{bmatrix} -2 & 1 & 1 & a \\ 1 & -2 & 1 & b \\ 1 & 1 & -2 & c \end{bmatrix}$

$$\begin{aligned}[A|B] &\xrightarrow{R_1 \leftrightarrow R_2} \begin{bmatrix} 1 & -2 & 1 & b \\ -2 & 1 & 1 & a \\ 1 & 1 & -2 & c \end{bmatrix} \\ &\xrightarrow{\substack{R_2 \rightarrow R_2 + 2R_1 \\ R_3 \rightarrow R_3 - R_1}} \begin{bmatrix} 1 & -2 & 1 & b \\ 0 & -3 & 3 & a + 2b \\ 0 & 3 & -3 & c - b \end{bmatrix} \\ &\xrightarrow{R_3 \rightarrow R_3 + R_2} \begin{bmatrix} 1 & -2 & 1 & b \\ 0 & -3 & 3 & c + 2b \\ 0 & 0 & 0 & a + b + c \end{bmatrix}\end{aligned}$$

Here $\rho(A) = 2$

The given system is consistent only when

$\rho([A|B]) = 2$ $\rho([A|B]) = 2$ only if $a + b + c = 0$

Hence proved.

