



Mathematics

12th Standard

VOLUME - I & II

Based on the Updated New Textbook

FREE
Practice Workbook
with
Practice Model
Question Papers

Salient Features

- Answers for **All Textual Questions** are given.
- Exhaustive Additional Questions & Answers in all chapters.
- Model Question Papers 1 to 6 (PTA) : Questions are incorporated in the appropriate sections.
- Govt. Model Question Paper-2019 [**Govt. MQP-2019**], Quarterly Exam - 2019 ; 2023 & 2024 [**Qy. - 2019, 2023 & 2024**], Half yearly Exam - 2019; 2023 & 2024 [**Hy. - 2019; 2023 & 2024**] , Public Exam March - 2020; 2023; 2024 & 2025 May - 2022 [**May - 2022 ; Mar. -2020; 2023 ; 2024 & 2025**], Govt. Supplementary Exam September - 2020, August - 2021, July - 2022 & 2024, June - 2023 & 2025 [**Sep -2020, Aug. - 2021; June - 2023 & 2025 ; July - 2022 & 2024**] and First & Second Revision Test - 2022 [**FRT & SRT - 2022**] questions are incorporated in the appropriate sections.
- Instant Supplementary Exam - 2025 Question Paper is given with answers.
- Quarterly Exam - 2025 Question Paper is given with answers.



SURA PUBLICATIONS

Chennai

For Orders Contact



80562 94222 / 81242 01000 / 81243 01000
96001 75757 / 98409 26027

All rights reserved © SURA Publications.

No part of this book may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, digitally, electronically, mechanically, photocopying, recorded or otherwise, without the written permission of the publishers. Strict action will be taken.

Code No : SG 322

Authors :

Mr. G.Selvaraj, M.Sc., M.Ed., M.Phil.,
Chennai

Edited by :

Mr. S. Sathish M.Sc., M.Phil.,

Reviewed by :

Mr.S. Niranjana, B.Tech., (NITT)PGDM (IIM)
Chennai

Our Guides for XI, XII Standard

- ❖ சுராவின் தமிழ் உரைநூல்
- ❖ Sura's Smart English
- ❖ Sura's Mathematics (EM/TM)
- ❖ Sura's Physics (EM/TM)
- ❖ Sura's Botany (EM/TM)
- ❖ Sura's Chemistry (EM/TM)
- ❖ Sura's Bio-Botany & Botany (EM/TM)
(Short Version & Long Version)
- ❖ Sura's Bio-Zoology & Zoology (EM/TM)
(Short Version & Long Version)
- ❖ Sura's Computer Science (EM/TM)
- ❖ Sura's Computer Applications (EM/TM)
- ❖ Sura's Commerce (EM/TM)
- ❖ Sura's Economics (EM/TM)
- ❖ Sura's Accountancy (EM/TM)
- ❖ Sura's Business Maths (EM)

PREFACE

Sir,

*An equation has no meaning, for me
unless it expresses a thought of GOD - Ramanujam*
[Statement to a friend]

Respected Principals, Correspondents, Head Masters / Head
Mistresses, Teachers,

From the bottom of our heart, we at SURA Publications
sincerely thank you for the support and patronage that you have
extended to us for more than a decade.

It is in our sincerest effort we take the pride of releasing **Sura's
Mathematics Guide Volume I and Volume II** for +2 Standard.
This guide has been authored and edited by qualified teachers
having teaching experience for over a decade in their respective
subject fields. This Guide has been reviewed by reputed Professors
who are currently serving as Head of the Department in esteemed
Universities and Colleges.

With due respect to Teachers, I would like to mention that
this guide will serve as a teaching companion to qualified teachers.
Also, this guide will be an excellent learning companion to students
with exhaustive exercises and in-text questions in addition to precise
answers for textual questions.

In complete cognizance of the dedicated role of Teachers, I
completely believe that our students will learn the subject effectively
with this guide and prove their excellence in Board Examinations.

I once again sincerely thank the Teachers, Parents and Students
for supporting and valuing our efforts.

God Bless all.

Subash Raj, B.E., M.S.
- Publisher
Sura Publications

All the Best

Head Office:

Sura Publications

1620, 'J' Block, 16th Main Road,
Anna Nagar, Chennai - 600 040.

Phones : 044 - 4862 9977, 044 - 4862 7755.

e-mail : orders@surabooks.com

website : www.surabooks.com

For Orders Contact



80562 94222

81242 01000

81243 01000

96001 75757

98409 26027

19/11/2025

CONTENT

Chapter No	Title	Page No
Volume - I		
1.	Applications of Matrices and Determinants	1-46
2.	Complex Numbers	47-80
3.	Theory of Equations	81-102
4.	Inverse Trigonometric Functions	103-130
5.	Two Dimensional Analytical Geometry-II	131-170
6.	Applications of Vector Algebra	171-220
Volume - II		
7.	Applications of Differential Calculus	221-274
8.	Differentials and Partial Derivatives	275-304
9.	Applications of Integration	305-336
10.	Ordinary Differential Equations	337-374
11.	Probability Distributions	375-404
12.	Discrete Mathematics	405-420
Instant Supplementary Exam - 2025 Question Paper with Answers		421-428
Quarterly Exam - 2025 Question Paper with Answers		429-436

SYLLABUS

Chapter No	Title	Month
1.	Applications of Matrices and Determinants (1.1 - 1.5)	June
2.	Complex Numbers (2.1 - 2.8)	July
3.	Theory of Equations (3.1 - 3.9)	
4.	Inverse Trigonometric Functions (4.1 - 4.5)	
I MID TERM TEST (June, July)		
4.	Inverse Trigonometric Functions (4.6 - 4.10)	August
5.	Two Dimensional Analytical Ge- ometry-II (5.1 - 5.7)	
6.	Applications of Vector Algebra (6.1 - 6.5)	
6.	Applications of Vector Algebra (6.6 - 6.10)	Sep.
QUARTERLY EXAMINATION (June to September)		

Chapter No	Title	Month
7.	Applications of Differential Calculus (7.1 - 7.10)	October
8.	Differentials and Partial Derivatives (8.1 - 8.3)	
8.	Differentials and Partial Derivatives (8.4 - 8.6)	Nov.
9.	Applications of Integration (9.1-9.5)	
9.	Applications of Integration (9.6-9.9)	Dec.
10.	Ordinary Differential Equations (10.1 - 10.8)	
HALFYEARELY EXAMINATION (June to December)		
11.	Probability Distributions (11.1-11.6)	January
12.	Discrete Mathematics (12.1 - 12.3)	
ANNUAL EXAMINATION (Full Portions)		

TO ORDER WITH US

SCHOOLS and TEACHERS:

We are grateful for your support and patronage to '**SURA PUBLICATIONS**'

Kindly prepare your order in your School letterhead and send it to us.

For Orders contact: 81242 01000 / 81243 01000

DIRECT DEPOSIT

A/c Name : **Sura Publications**
Our A/c No. : **36550290536**
Bank Name : **STATE BANK OF INDIA**
Bank Branch : Padi
IFSC : SBIN0005083

A/c Name : **Sura Publications**
Our A/c No. : **21000210001240**
Bank Name : **UCO BANK**
Bank Branch : Anna Nagar West
IFSC : UCBA0002100

A/c Name : **Sura Publications**
Our A/c No. : **6502699356**
Bank Name : **INDIAN BANK**
Bank Branch : Asiad Colony
IFSC : IDIB000A098

A/c Name : **Sura Publications**
Our A/c No. : **1154135000017684**
Bank Name : **KVB BANK**
Bank Branch : Anna Nagar
IFSC : KVBL0001154

A/c Name : **Sura Publications**
Our A/c No. : **13240200032412**
Bank Name : **FEDERAL BANK**
Bank Branch : Anna Nagar
IFSC : FDRL0001324

A/c Name : **Sura Publications**
Our A/c No. : **50200031530945**
Bank Name : **HDFC BANK**
Bank Branch : Cenotaph Road, Teynampet
IFSC : HDFC0001216

After Deposit, please send challan and order to our address.

email to : orders@surabooks.com / Whatsapp : 81242 01000.



For Google Pay :
98409 26027



For PhonePe :
98409 26027



DEMAND DRAFT / CHEQUE

Please send Demand Draft / cheque in favour of '**SURA PUBLICATIONS**' payable at **Chennai**. The Demand Draft / cheque should be sent with your order in School letterhead.

STUDENTS :

Order via Money Order (M/O) to



SURA PUBLICATIONS

1620, 'J' Block, 16th Main Road, Anna Nagar, Chennai - 600 040.

Phones : 044-4862 9977, 044-4862 7755.

Mobile : 96001 75757 / 81242 01000 / 81243 01000.

email : orders@surabooks.com Website : www.surabooks.com

Volume - I

MATHEMATICS

CONTENTS

1. Applications of Matrices and Determinants	3 - 46
2. Complex Numbers.....	47 - 80
3. Theory of Equations	81 - 102
4. Inverse Trigonometric Functions.....	103 - 130
5. Two Dimensional Analytical Geometry-II	131 - 170
6. Applications of Vector Algebra	171 - 220

CHAPTER

1

APPLICATIONS OF MATRICES AND DETERMINANTS

MUST KNOW DEFINITIONS

- ✦ If $|A| \neq 0$, then A is a non-singular matrix and if $|A| = 0$, then A is a singular matrix.
- ✦ The adjoint matrix of A is defined as the transpose of the matrix of co-factors of A.
- ✦ If $AB = BA = I_n$, then the matrix B is called the inverse of A.
- ✦ If a square matrix has an inverse, then it is unique.
- ✦ A^{-1} exists if and only if A is non-singular.
- ✦ Singular matrix has no inverse.
- ✦ If A is non – singular and $AB = AC$, then $B = C$ (left cancellation law).
- ✦ If A is non – singular and $BA = CA$ then $B = C$ (Right cancellation law).
- ✦ If A and B are any two non-singular square matrices of order n , then $\text{adj} (AB) = (\text{adj} B) (\text{adj} A)$
- ✦ A square matrix A is called orthogonal if $AA^T = A^T A = I$
- ✦ Two matrices A and B of same order are said to be **equivalent** if one can be obtained from the other by the applications of elementary transformations (A~B).
- ✦ A non – zero matrix is in a **row - echelon** form if all zero rows occur as bottom rows of the matrix and if the first non – zero element in any lower row occurs to the right of the first non – zero entry in the higher row.
- ✦ The **rank** of a matrix A is defined as the order of a highest order non – vanishing minor of the matrix A [$\rho(A)$].
- ✦ The **rank** of a non – zero matrix is equal to the number of non – zero rows in a row – echelon form of the matrix.
- ✦ An **elementary matrix** is a matrix which is obtained from an identity matrix by applying only one elementary transformation. Every non-singular matrix can be transformed to an identity matrix by a sequence of elementary row operations.
- ✦ A system of linear equations having atleast one solution is said to be **consistent**.
- ✦ A system of linear equations having no solutions is said to be **inconsistent**.

IMPORTANT FORMULAE TO REMEMBER

- ✦ Co – factor of a_{ij} is $A_{ij} = (-1)^{i+j} M_{ij}$, where M_{ij} is the minor of a_{ij}
- ✦ For every square matrix A of order n , $A (\text{adj } A) = (\text{adj } A)A = |A| I_n$
 $AA^{-1} = A^{-1}A = I_n$
- ✦ If A is non – Singular then
 - (i) $|A^{-1}| = \frac{1}{|A|}$ (ii) $(A^T)^{-1} = (A^{-1})^T$
 - (iii) $(\lambda A^{-1}) = \frac{1}{\lambda} A^{-1}$ where λ is a non – zero scalar.

Reversal law for inverses :

- ✦ $(AB)^{-1} = B^{-1}A^{-1}$ where A, B are non – singular matrices of same order.

Law of double inverse :

- ✦ If A is non - singular, A^{-1} is also non – singular and $(A^{-1})^{-1} = A$.
- ✦ If A is a non - singular square matrix of order n , then
 - (i) $(\text{adj } A)^{-1} = \text{adj } (A^{-1}) = \frac{1}{|A|} \cdot A$
 - (ii) $|\text{adj } A| = |A|^{n-1}$
 - (iii) $\text{adj } (\text{adj } A) = |A|^{n-2}A$
 - (iv) $\text{adj } (\lambda A) = \lambda^{n-1} \text{adj } (A)$ where λ is a non – zero scalar
 - (v) $|\text{adj } (\text{adj } A)| = |A|^{(n-1)^2}$
 - (vi) $(\text{adj } A)^T = \text{adj } (A^T)$
- ✦ If a matrix contains at least one non – zero element, then $\rho(A) \geq 1$.
- ✦ The rank of identity matrix I_n is n . If A is an $m \times n$ matrix then $\rho(A) \leq \min \{ m, n \}$.
- ✦ A square matrix A of order n is invertible if and only if $\rho(A) = n$.
- ✦ Transforming a non-singular matrix A to the form I_n , by applying row operations is called Gauss – Jordan method.

Matrix – Inversion method :

- ✦ The solution for $AX = B$ is $X = A^{-1}B$ where A and B are square matrices of same order and non – singular

Cramer's Rule :

- ✦ If $\Delta = 0$, Cramer's rule cannot be applied $x_1 = \frac{\Delta_1}{\Delta}, x_2 = \frac{\Delta_2}{\Delta}, x_3 = \frac{\Delta_3}{\Delta}$

Gaussian Elimination method :

Transform the augmented matrix of the system of linear equations into row – echelon form and then solve by back substitution method.

Rouches capelli Theorem :

A system of equations $AX = B$ is consistent if and if $\rho(A) = \rho([A|B])$

- (i) If $\rho(A) = \rho([A|B]) = n$, the number of unknowns, then the system is consistent and has a unique solution.
- (ii) If $\rho(A) = \rho([A|B]) = n - k, k \neq 0$ then the system is consistent and has infinitely many solutions.
- (iii) If $\rho(A) \neq \rho([A|B])$, then the system is inconsistent and has no solution.

Homogeneous system of linear equations :

(i) If $\rho(A) = \rho([A|B]) = n$, then the system has a unique solution which is the trivial solution for trivial solution, $|A| \neq 0$

(ii) If $\rho(A) = \rho([A|B]) < n$, the system has a non – trivial solution.

For non – trivial solution, $|A| = 0$.

$$\star \quad A^{-1} = \pm \frac{1}{\sqrt{|adj A|}} \cdot adj A \quad \star \quad A = \pm \frac{1}{\sqrt{|adj A|}} \cdot adj (adj A)$$

EXERCISE 1.1**1. Find the adjoint of the following :**

(i) $\begin{bmatrix} -3 & 4 \\ 6 & 2 \end{bmatrix}$ (ii) $\begin{bmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{bmatrix}$ (iii) $\frac{1}{3} \begin{bmatrix} 2 & 2 & 1 \\ -2 & 1 & 2 \\ 1 & -2 & 2 \end{bmatrix}$

Sol. (i) Let $A = \begin{pmatrix} -3 & 4 \\ 6 & 2 \end{pmatrix}$
 $adj A = \begin{pmatrix} 2 & -4 \\ -6 & -3 \end{pmatrix}$

[Interchange the elements in the leading diagonal and change the sign of the elements in off diagonal]

(ii) Let $A = \begin{pmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{pmatrix}$
 $adj A = \begin{pmatrix} + \begin{vmatrix} 4 & 1 \\ 7 & 2 \end{vmatrix} - \begin{vmatrix} 3 & 1 \\ 3 & 2 \end{vmatrix} + \begin{vmatrix} 3 & 4 \\ 3 & 7 \end{vmatrix} \\ - \begin{vmatrix} 3 & 1 \\ 2 & 1 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 3 & 2 \end{vmatrix} - \begin{vmatrix} 2 & 3 \\ 3 & 7 \end{vmatrix} \\ + \begin{vmatrix} 3 & 1 \\ 4 & 1 \end{vmatrix} - \begin{vmatrix} 2 & 1 \\ 3 & 1 \end{vmatrix} + \begin{vmatrix} 2 & 3 \\ 3 & 4 \end{vmatrix} \end{pmatrix}^T$

$$= \begin{bmatrix} + (8-7) - (6-3) + (21-12) \\ - (6-7) + (4-3) - (14-9) \\ + (3-4) - (2-3) + (8-9) \end{bmatrix}^T = \begin{bmatrix} 1 & -3 & 9 \\ 1 & 1 & -5 \\ -1 & 1 & -1 \end{bmatrix}^T$$

$$adj A = \begin{bmatrix} 1 & 1 & -1 \\ -3 & 1 & 1 \\ 9 & -5 & -1 \end{bmatrix}$$

(iii) Let $A = \begin{bmatrix} 2 & 2 & 1 \\ -2 & 1 & 2 \\ 1 & -2 & 2 \end{bmatrix}$ and $\lambda = \frac{1}{3}$

Since $adj (\lambda A) = \lambda^{n-1} (adj A)$

we get $adj \left(\frac{1}{3} \begin{bmatrix} 2 & 2 & 1 \\ -2 & 1 & 2 \\ 1 & -2 & 2 \end{bmatrix} \right) = \left(\frac{1}{3} \right)^2$

$$adj \begin{pmatrix} 2 & 2 & 1 \\ -2 & 1 & 2 \\ 1 & -2 & 2 \end{pmatrix}$$

∴ Required adjoint matrix

$$\begin{aligned} & \begin{bmatrix} + \begin{vmatrix} 1 & 2 \\ -2 & 2 \end{vmatrix} - \begin{vmatrix} -2 & 2 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} -2 & 1 \\ 1 & -2 \end{vmatrix} \\ - \begin{vmatrix} 2 & 1 \\ -2 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} - \begin{vmatrix} 2 & 2 \\ 1 & -2 \end{vmatrix} \\ + \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} - \begin{vmatrix} 2 & 1 \\ -2 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 2 \\ -2 & 1 \end{vmatrix} \end{bmatrix}^T \\ &= \frac{1}{9} \begin{bmatrix} (2+4) - (-4-2) + (4-1) \\ - (4+2) + (4-1) - (-4-2) \\ + (4-1) - (4+2) + (2+4) \end{bmatrix}^T \\ &= \frac{1}{9} \begin{bmatrix} 6 & 6 & 3 \\ -6 & 3 & 6 \\ 3 & -6 & 6 \end{bmatrix}^T = \frac{1}{9} \begin{bmatrix} 6 & -6 & 3 \\ 6 & 3 & -6 \\ 3 & 6 & 6 \end{bmatrix} \\ &= \frac{3}{9} \begin{bmatrix} 2 & -2 & 1 \\ 2 & 1 & -2 \\ 1 & 2 & 2 \end{bmatrix} \end{aligned}$$

[Taking 3 common from each entry]

$$= \frac{1}{3} \begin{bmatrix} 2 & -2 & 1 \\ 2 & 1 & -2 \\ 1 & 2 & 2 \end{bmatrix}$$

2. Find the inverse (if it exists) of the following :

(i) $\begin{bmatrix} -2 & 4 \\ 1 & -3 \end{bmatrix}$ (ii) $\begin{bmatrix} 5 & 1 & 1 \\ 1 & 5 & 1 \\ 1 & 1 & 5 \end{bmatrix}$ (iii) $\begin{bmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{bmatrix}$

Sol. (i) Let $A = \begin{bmatrix} -2 & 4 \\ 1 & -3 \end{bmatrix}$

$$|A| = \begin{vmatrix} -2 & 4 \\ 1 & -3 \end{vmatrix} = 6 - 4 = 2 \neq 0$$

Since A is non-singular, A^{-1} exists

$$A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$\text{Now, adj } A = \begin{bmatrix} -3 & -4 \\ -1 & -2 \end{bmatrix}$$

[Inter change the entries in leading diagonal and change the sign of elements in the off diagonal]

$$\therefore A^{-1} = \frac{1}{2} \begin{bmatrix} -3 & -4 \\ -1 & -2 \end{bmatrix}$$

$$(ii) \text{ Let } A = \begin{bmatrix} 5 & 1 & 1 \\ 1 & 5 & 1 \\ 1 & 1 & 5 \end{bmatrix}$$

Expanding along R_1 ,

$$\begin{aligned} |A| &= 5 \begin{vmatrix} 5 & 1 \\ 1 & 5 \end{vmatrix} - 1 \begin{vmatrix} 1 & 1 \\ 1 & 5 \end{vmatrix} + 1 \begin{vmatrix} 1 & 5 \\ 1 & 1 \end{vmatrix} \\ &= 5(25 - 1) - 1(5 - 1) + 1(1 - 5) \\ &= 5(24) - 1(4) + 1(-4) \\ &= 120 - 4 - 4 = 120 - 8 = 112 \neq 0 \end{aligned}$$

Since A is non singular, A^{-1} exists.

$$\begin{aligned} \text{adj } A &= \begin{bmatrix} + \begin{vmatrix} 5 & 1 \\ 1 & 5 \end{vmatrix} - \begin{vmatrix} 1 & 1 \\ 1 & 5 \end{vmatrix} + \begin{vmatrix} 1 & 5 \\ 1 & 1 \end{vmatrix} \\ - \begin{vmatrix} 1 & 1 \\ 1 & 5 \end{vmatrix} + \begin{vmatrix} 5 & 1 \\ 1 & 5 \end{vmatrix} - \begin{vmatrix} 5 & 1 \\ 1 & 1 \end{vmatrix} \\ + \begin{vmatrix} 1 & 1 \\ 5 & 1 \end{vmatrix} - \begin{vmatrix} 5 & 1 \\ 1 & 1 \end{vmatrix} + \begin{vmatrix} 5 & 1 \\ 1 & 5 \end{vmatrix} \end{bmatrix}^T \\ &= \begin{bmatrix} +(25-1)-(5-1)+(1-5) \\ -(5-1)+(25-1)-(5-1) \\ +(1-5)-(5-1)+(25-1) \end{bmatrix}^T \\ &= \begin{bmatrix} 24 & -4 & -4 \\ -4 & 24 & -4 \\ -4 & -4 & 24 \end{bmatrix}^T = \begin{bmatrix} 24 & -4 & -4 \\ -4 & 24 & -4 \\ -4 & -4 & 24 \end{bmatrix} \end{aligned}$$

Taking 4 common from every entry we get,

$$\begin{aligned} \text{adj } A &= 4 \begin{bmatrix} 6 & -1 & -1 \\ -1 & 6 & -1 \\ -1 & -1 & 6 \end{bmatrix} \\ \therefore A^{-1} &= \frac{1}{|A|} \text{adj } A = \frac{1}{112} \cdot 4 \begin{bmatrix} 6 & -1 & -1 \\ -1 & 6 & -1 \\ -1 & -1 & 6 \end{bmatrix} \end{aligned}$$

$$= \frac{1}{28} \begin{bmatrix} 6 & -1 & -1 \\ -1 & 6 & -1 \\ -1 & -1 & 6 \end{bmatrix}$$

$$(iii) \text{ Let } A = \begin{bmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{bmatrix}$$

Expanding along R_1 we get,

$$\begin{aligned} |A| &= 2 \begin{vmatrix} 4 & 1 \\ 7 & 2 \end{vmatrix} - 3 \begin{vmatrix} 3 & 1 \\ 3 & 2 \end{vmatrix} + 1 \begin{vmatrix} 3 & 4 \\ 3 & 7 \end{vmatrix} \\ &= 2(8 - 7) - 3(6 - 3) + 1(21 - 12) \\ &= 2(1) - 3(3) + 1(9) \\ &= 2 - 9 + 9 = 2 \neq 0 \end{aligned}$$

Since A is a non-singular matrix, A^{-1} exists

$$\text{adj } A = \begin{bmatrix} + \begin{vmatrix} 4 & 1 \\ 7 & 2 \end{vmatrix} - \begin{vmatrix} 3 & 1 \\ 3 & 2 \end{vmatrix} + \begin{vmatrix} 3 & 4 \\ 3 & 7 \end{vmatrix} \\ - \begin{vmatrix} 3 & 1 \\ 7 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 3 & 2 \end{vmatrix} - \begin{vmatrix} 2 & 3 \\ 3 & 7 \end{vmatrix} \\ + \begin{vmatrix} 3 & 1 \\ 4 & 1 \end{vmatrix} - \begin{vmatrix} 2 & 1 \\ 3 & 1 \end{vmatrix} + \begin{vmatrix} 2 & 3 \\ 3 & 4 \end{vmatrix} \end{bmatrix}^T$$

$$= \begin{bmatrix} +(8-7)-(6-3)+(21-12) \\ -(6-7)+(4-3)-(14-9) \\ +(3-4)-(2-3)+(8-9) \end{bmatrix}^T = \begin{bmatrix} 1 & -3 & 9 \\ 1 & 1 & -5 \\ -1 & 1 & -1 \end{bmatrix}^T$$

$$\text{adj } A = \begin{bmatrix} 1 & 1 & -1 \\ -3 & 1 & 1 \\ 9 & -5 & -1 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$\Rightarrow A^{-1} = \frac{1}{2} \begin{bmatrix} 1 & 1 & -1 \\ -3 & 1 & 1 \\ 9 & -5 & -1 \end{bmatrix}$$

$$3. \text{ If } F(\alpha) = \begin{bmatrix} \cos \alpha & 0 & \sin \alpha \\ 0 & 1 & 0 \\ -\sin \alpha & 0 & \cos \alpha \end{bmatrix}, \text{ show that}$$

$$[F(\alpha)]^{-1} = F(-\alpha).$$

[Hy - 2019 & 2024; FRT - 2022; Mar - 2023]

$$\text{Sol. Given that } F(\alpha) = \begin{bmatrix} \cos \alpha & 0 & \sin \alpha \\ 0 & 1 & 0 \\ -\sin \alpha & 0 & \cos \alpha \end{bmatrix}.$$

Expanding along R_1 we get,

$$|F(\alpha)| = \cos \alpha \begin{vmatrix} 1 & 0 \\ 0 & \cos \alpha \end{vmatrix} - 0 + \sin \alpha \begin{vmatrix} 0 & 1 \\ -\sin \alpha & 0 \end{vmatrix}$$

$$= \cos \alpha (\cos - 0) + \sin \alpha (0 + \sin \alpha)$$

$$= \cos^2 \alpha + \sin^2 \alpha = 1 \neq 0$$

Since $F(\alpha)$ is a non-singular matrix, $[F(\alpha)]^{-1}$ exists.

Now, $\text{adj } (F(\alpha)) =$

$$\begin{bmatrix} + \begin{vmatrix} 1 & 0 \\ 0 & \cos \alpha \end{vmatrix} - \begin{vmatrix} 0 & 0 \\ \sin \alpha & \cos \alpha \end{vmatrix} + \begin{vmatrix} 0 & 1 \\ -\sin \alpha & 0 \end{vmatrix} \\ - \begin{vmatrix} 0 & \sin \alpha \\ 0 & \cos \alpha \end{vmatrix} + \begin{vmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{vmatrix} - \begin{vmatrix} \cos \alpha & 0 \\ -\sin \alpha & 0 \end{vmatrix} \\ + \begin{vmatrix} 0 & \sin \alpha \\ 1 & 0 \end{vmatrix} - \begin{vmatrix} \cos \alpha & \sin \alpha \\ 0 & 0 \end{vmatrix} + \begin{vmatrix} \cos \alpha & 0 \\ 0 & 1 \end{vmatrix} \end{bmatrix}^T$$

$$= \begin{bmatrix} +(\cos \alpha - 0) & -(0) & +(0 + \sin \alpha) \\ -(0) & +(\cos^2 \alpha + \sin^2 \alpha) & -(0) \\ +(0 - \sin \alpha) & -(0) & +(\cos - 0) \end{bmatrix}^T$$

$$= \begin{bmatrix} \cos \alpha & 0 & +\sin \alpha \\ 0 & 1 & 0 \\ -\sin \alpha & 0 & \cos \alpha \end{bmatrix}^T = \begin{bmatrix} \cos \alpha & 0 & -\sin \alpha \\ 0 & 1 & 0 \\ \sin \alpha & 0 & \cos \alpha \end{bmatrix}$$

$$\therefore F(\alpha)^{-1} = \frac{1}{|F(\alpha)|} \text{adj } (F(\alpha))$$

$$[F(\alpha)]^{-1} = \frac{1}{1} \begin{bmatrix} \cos \alpha & 0 & -\sin \alpha \\ 0 & 1 & 0 \\ \sin \alpha & 0 & \cos \alpha \end{bmatrix}$$

$$= \begin{bmatrix} \cos \alpha & 0 & -\sin \alpha \\ 0 & 1 & 0 \\ \sin \alpha & 0 & \cos \alpha \end{bmatrix} \quad \dots (1)$$

$$\text{Now, } F(-\alpha) = \begin{bmatrix} \cos(-\alpha) & 0 & \sin(-\alpha) \\ 0 & 1 & 0 \\ -\sin(-\alpha) & 0 & \cos(-\alpha) \end{bmatrix}$$

$$= \begin{bmatrix} \cos \alpha & 0 & -\sin \alpha \\ 0 & 1 & 0 \\ \sin \alpha & 0 & \cos \alpha \end{bmatrix} \quad \dots (2)$$

$\because \cos \alpha$ is an even function, $\cos(-\alpha) = \cos \alpha$ and $\sin \alpha$ is an odd function, $\sin(-\alpha) = -\sin \alpha$

From (1) and (2)

$$[F(\alpha)]^{-1} = F(-\alpha)$$

Hence proved.

4. If $A = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$, show that $A^2 - 3A - 7I_2 = 0_2$.

Hence find A^{-1} .

Sol. Given $A = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$

$$A^2 = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} = \begin{bmatrix} 25-3 & 15-6 \\ -5+2 & -3+4 \end{bmatrix}$$

$$= \begin{bmatrix} 22 & 9 \\ -3 & 1 \end{bmatrix} \quad \therefore A^2 - 3A - 7I_2$$

$$= \begin{bmatrix} 22 & 9 \\ -3 & 1 \end{bmatrix} - 3 \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} - 7 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 22-15-7 & 9-9+0 \\ -3+3+0 & 1+6-7 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0_2$$

Hence proved.

$$\therefore A^2 - 3A - 7I_2 = 0$$

Post-multiplying by A^{-1} we get,

$$A^2 \cdot A^{-1} - 3AA^{-1} - 7I_2 A^{-1} = 0 \cdot A^{-1}$$

$$\Rightarrow A(AA^{-1}) - 3(AA^{-1}) - 7(A^{-1}) = 0$$

$$[\because I_2 A^{-1} = A^{-1} \text{ and } (0)A^{-1} = 0]$$

$$\Rightarrow AI - 3I - 7A^{-1} = 0$$

$$[\because AA^{-1} = I]$$

$$\Rightarrow AI - 3I = 7A^{-1}$$

$$\Rightarrow A^{-1} = \frac{1}{7} [A - 3I] \quad [\because AI = A]$$

$$\Rightarrow A^{-1} = \frac{1}{7} \left[\begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} - 3 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right]$$

$$\Rightarrow A^{-1} = \frac{1}{7} \begin{bmatrix} 5-3 & 3-0 \\ -1-0 & -2-3 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 2 & 3 \\ -1 & -5 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{7} \begin{bmatrix} 2 & 3 \\ -1 & -5 \end{bmatrix}$$

5. If $A = \frac{1}{9} \begin{bmatrix} -8 & 1 & 4 \\ 4 & 4 & 7 \\ 1 & -8 & 4 \end{bmatrix}$, prove that $A^{-1} = A^T$.

Sol.

To prove $A^{-1} = A^T$

$$AA^{-1} = AA^T$$

It is enough to prove $AA^T = I$

$$AA^T = \frac{1}{9} \begin{bmatrix} -8 & 1 & 4 \\ 4 & 4 & 7 \\ 1 & -8 & 4 \end{bmatrix} \frac{1}{9} \begin{bmatrix} -8 & 1 & 4 \\ 1 & 4 & -8 \\ 4 & 7 & 4 \end{bmatrix}$$

$$= \frac{1}{81} \begin{bmatrix} 64+1+16 & -32+4+28 & -8-8+16 \\ -32+4+28 & 16+16+49 & 4-32-28 \\ -8-8+16 & 4-32+28 & 1+64+16 \end{bmatrix}$$

$$= \frac{1}{81} \begin{bmatrix} 81 & 0 & 0 \\ 0 & 81 & 0 \\ 0 & 0 & 81 \end{bmatrix} = \frac{81}{81} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

Therefore $A^{-1} = A^T$

6. If $A = \begin{bmatrix} 8 & -4 \\ -5 & 3 \end{bmatrix}$, verify that $A(\text{adj } A) = (\text{adj } A)A = |A| I_2$. [Sep. - 2020; Aug. - 2021]

Sol. Given $A = \begin{bmatrix} 8 & -4 \\ -5 & 3 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 3 & 4 \\ 5 & 8 \end{bmatrix}$$

[Interchange the elements in the leading diagonal and change the sign of the elements in the off diagonal]

$$|A| = 24 - 20 = 4$$

$$\therefore A(\text{adj } A) = \begin{bmatrix} 8 & -4 \\ -5 & 3 \end{bmatrix} \begin{bmatrix} 3 & 4 \\ 5 & 8 \end{bmatrix}$$

$$= \begin{bmatrix} 24-20 & 32-32 \\ -15+15 & -20+24 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \quad \dots(1)$$

$$(\text{adj } A)(A) = \begin{bmatrix} 3 & 4 \\ 5 & 8 \end{bmatrix} \begin{bmatrix} 8 & -4 \\ -5 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 24-20 & -12+12 \\ 40-40 & -20+24 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \quad \dots(2)$$

$$|A| I_2 = 4 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \quad \dots(3)$$

From (1), (2) and (3), it is proved that

$A(\text{adj } A) = (\text{adj } A)A = |A| I_2$ is verified.

7. If $A = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & -3 \\ 5 & 2 \end{bmatrix}$, verify that $(AB)^{-1} = B^{-1}A^{-1}$. [Qy. - 2023]

Sol. Given $A = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & -3 \\ 5 & 2 \end{bmatrix}$

$$AB = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix} \begin{bmatrix} -1 & -3 \\ 5 & 2 \end{bmatrix} = \begin{bmatrix} -3+10 & -9+4 \\ -7+25 & -21+10 \end{bmatrix}$$

$$= \begin{bmatrix} 7 & -5 \\ 18 & -11 \end{bmatrix}$$

$$|AB| = -77 + 90 = 13 \neq 0 \Rightarrow (AB)^{-1} \text{ exists}$$

$$|A| = 15 - 14 = 1 \neq 0 \Rightarrow A^{-1} \text{ exists}$$

$$|B| = -2 + 15 = 13 \neq 0 \Rightarrow B^{-1} \text{ exists}$$

$$(AB)^{-1} = \frac{1}{|AB|} \text{adj } (AB) = \frac{1}{13} \begin{pmatrix} -11 & 5 \\ -18 & 7 \end{pmatrix} \dots(1)$$

$$B^{-1} = \frac{1}{|B|} (\text{adj } B) = \frac{1}{13} \begin{pmatrix} 2 & 3 \\ -5 & -1 \end{pmatrix}$$

$$A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{1} \begin{pmatrix} 5 & -2 \\ -7 & 3 \end{pmatrix}$$

$$= \begin{pmatrix} 5 & -2 \\ -7 & 3 \end{pmatrix}$$

$$\therefore B^{-1}A^{-1} = \frac{1}{13} \begin{pmatrix} 2 & 3 \\ -5 & -1 \end{pmatrix} \begin{pmatrix} 5 & -2 \\ -7 & 3 \end{pmatrix}$$

$$= \frac{1}{13} \begin{pmatrix} 10-21 & -4+9 \\ -25+7 & 10-3 \end{pmatrix}$$

$$= \frac{1}{13} \begin{pmatrix} -11 & 5 \\ -18 & 7 \end{pmatrix} \quad \dots(2)$$

From (1) and (2) it is prove that

$$(AB)^{-1} = B^{-1}A^{-1}.$$

8. If $\text{adj } (A) = \begin{bmatrix} 2 & -4 & 2 \\ -3 & 12 & -7 \\ -2 & 0 & 2 \end{bmatrix}$, find A .

Sol. Given $\text{adj } A = \begin{bmatrix} 2 & -4 & 2 \\ -3 & 12 & -7 \\ -2 & 0 & 2 \end{bmatrix}$

$$\text{We know that } A = \pm \frac{1}{\sqrt{|\text{adj } A|}} \text{adj } (\text{adj } A) \quad \dots(1)$$

$$|\text{adj } A| = 2 \begin{vmatrix} 12 & -7 \\ 0 & 2 \end{vmatrix} + 4 \begin{vmatrix} -3 & -7 \\ -2 & 2 \end{vmatrix} + 2 \begin{vmatrix} -3 & 12 \\ -2 & 0 \end{vmatrix}$$

$$= 2(24-0) + 4(-6-14) + 2(0+24)$$

$$= 2(24) + 4(-20) + 2(24) = 48 - 80 + 48$$

$$= 96 - 80 = 16 \quad \dots(2)$$

Now, $\text{adj } (\text{adj } A)$

$$= \begin{bmatrix} + \begin{vmatrix} 12 & -7 \\ 0 & 2 \end{vmatrix} - \begin{vmatrix} -3 & -7 \\ -2 & 2 \end{vmatrix} + \begin{vmatrix} -3 & 12 \\ -2 & 0 \end{vmatrix} \\ - \begin{vmatrix} -4 & 2 \\ 0 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 2 \\ -2 & 2 \end{vmatrix} - \begin{vmatrix} 2 & -4 \\ -2 & 0 \end{vmatrix} \\ + \begin{vmatrix} -4 & 2 \\ 12 & -7 \end{vmatrix} - \begin{vmatrix} 2 & 2 \\ -3 & -7 \end{vmatrix} + \begin{vmatrix} 2 & -4 \\ -3 & 12 \end{vmatrix} \end{bmatrix}^T$$

$$\begin{aligned}
 &= \begin{bmatrix} +(24-0) - (-6-14) + (0+24) \\ -(-8-0) + (4+4) - (0-8) \\ +(28-24) - (-14+6) + (24-12) \end{bmatrix}^T \\
 &= \begin{bmatrix} 24 & 20 & 24 \\ 8 & 8 & 8 \\ 4 & 8 & 12 \end{bmatrix}^T = \begin{bmatrix} 24 & 8 & 4 \\ 20 & 8 & 8 \\ 24 & 8 & 12 \end{bmatrix} \\
 &= 4 \begin{bmatrix} 6 & 2 & 1 \\ 5 & 2 & 2 \\ 6 & 2 & 3 \end{bmatrix}
 \end{aligned}$$

Substituting (2) and (3) in (1) we get,

$$A = \pm \frac{1}{\sqrt{16}} \cdot 4 \begin{bmatrix} 6 & 2 & 1 \\ 5 & 2 & 2 \\ 6 & 2 & 3 \end{bmatrix}$$

$$A = \pm \frac{4}{4} \begin{bmatrix} 6 & 2 & 1 \\ 5 & 2 & 2 \\ 6 & 2 & 3 \end{bmatrix} = \pm \begin{bmatrix} 6 & 2 & 1 \\ 5 & 2 & 2 \\ 6 & 2 & 3 \end{bmatrix}$$

9. If $\text{adj}(A) = \begin{bmatrix} 0 & -2 & 0 \\ 6 & 2 & -6 \\ -3 & 0 & 6 \end{bmatrix}$, find A^{-1} .

[PTA - 6; FRT - 2022; June & Qy. - 2023; July - 2024]

Sol. Given $\text{adj}(A) = \begin{bmatrix} 0 & -2 & 0 \\ 6 & 2 & -6 \\ -3 & 0 & 6 \end{bmatrix}$

We know that $A^{-1} = \pm \frac{1}{\sqrt{|\text{adj} A|}} (\text{adj} A) \dots (1)$

$$|\text{adj} A| = 0 + 2 \begin{vmatrix} 6 & -6 \\ -3 & 6 \end{vmatrix} + 0$$

[Expanded along R_1]

$$= 2(36 - 18) = 2(18) = 36$$

$$\therefore A^{-1} = \pm \frac{1}{\sqrt{36}} \begin{bmatrix} 0 & -2 & 0 \\ 6 & 2 & -6 \\ -3 & 0 & 6 \end{bmatrix}$$

$$= \pm \frac{1}{6} \begin{bmatrix} 0 & -2 & 0 \\ 6 & 2 & -6 \\ -3 & 0 & 6 \end{bmatrix}$$

10. Find $\text{adj}(\text{adj}(A))$ if $\text{adj} A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix}$.

Sol. Given $\text{adj} A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix}$

Now $\text{adj}(\text{adj} A) = \begin{bmatrix} + \begin{vmatrix} 2 & 0 \\ 0 & 1 \end{vmatrix} - \begin{vmatrix} 0 & 0 \\ -1 & 1 \end{vmatrix} + \begin{vmatrix} 0 & 2 \\ -1 & 0 \end{vmatrix} \\ - \begin{vmatrix} 0 & 1 \\ 0 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 1 \\ -1 & 1 \end{vmatrix} - \begin{vmatrix} 1 & 0 \\ -1 & 0 \end{vmatrix} \\ + \begin{vmatrix} 0 & 1 \\ 2 & 0 \end{vmatrix} - \begin{vmatrix} 1 & 1 \\ 0 & 0 \end{vmatrix} + \begin{vmatrix} 1 & 0 \\ 0 & 2 \end{vmatrix} \end{bmatrix}^T$

$$= \begin{bmatrix} +(2-0) & -(0) & +(0+2) \\ -(0) & +(1+1) & -(0) \\ +(0-2) & -(0) & +(2-0) \end{bmatrix}^T = \begin{bmatrix} 2 & 0 & 2 \\ 0 & 2 & 0 \\ -2 & 0 & 2 \end{bmatrix}^T$$

$$\text{adj}(\text{adj} A) = \begin{bmatrix} 2 & 0 & -2 \\ 0 & 2 & 0 \\ 2 & 0 & 2 \end{bmatrix}$$

11. $A = \begin{bmatrix} 1 & \tan x \\ -\tan x & 1 \end{bmatrix}$, show that

$$A^T A^{-1} = \begin{bmatrix} \cos 2x & -\sin 2x \\ \sin 2x & \cos 2x \end{bmatrix}.$$

Sol. $A = \begin{bmatrix} 1 & \tan x \\ -\tan x & 1 \end{bmatrix}$

$$|A| = \begin{vmatrix} 1 & \tan x \\ -\tan x & 1 \end{vmatrix} = 1 + \tan^2 x = \sec^2 x$$

$$\text{adj} A = \begin{bmatrix} 1 & -\tan x \\ \tan x & 1 \end{bmatrix}$$

We know, $A^{-1} = \frac{1}{|A|} \text{Adj} A$

$$= \frac{1}{\sec^2 x} \begin{bmatrix} 1 & -\tan x \\ \tan x & 1 \end{bmatrix} = \cos^2 x \begin{bmatrix} 1 & -\tan x \\ \tan x & 1 \end{bmatrix}$$

$$= \cos^2 x \begin{bmatrix} 1 & -\frac{\sin x}{\cos x} \\ \frac{\sin x}{\cos x} & 1 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} \cos^2 x & -\cos x \sin x \\ \cos x \sin x & \cos^2 x \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & \tan x \\ -\tan x & 1 \end{bmatrix}$$

$$A^T = \begin{bmatrix} 1 & -\frac{\sin x}{\cos x} \\ \frac{\sin x}{\cos x} & 1 \end{bmatrix}$$

$$A^T A^{-1} = \begin{bmatrix} 1 & -\frac{\sin x}{\cos x} \\ \frac{\sin x}{\cos x} & 1 \end{bmatrix} \begin{bmatrix} \cos^2 x & -\cos x \sin x \\ \cos x \sin x & \cos^2 x \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2 x - \sin^2 x & -2 \sin x \cos x \\ 2 \sin x \cos x & \cos^2 x - \sin^2 x \end{bmatrix} = \begin{bmatrix} \cos 2x & -\sin 2x \\ \sin 2x & \cos 2x \end{bmatrix}$$

12. Find the matrix A for which

$$A \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} = \begin{bmatrix} 14 & 7 \\ 7 & 7 \end{bmatrix}$$

Sol. Given $A \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} = \begin{bmatrix} 14 & 7 \\ 7 & 7 \end{bmatrix}$

Let $B = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$ and $C = \begin{bmatrix} 14 & 7 \\ 7 & 7 \end{bmatrix}$

$$\therefore AB = C$$

Post multiply by B^{-1} we get

$$A(BB^{-1}) = CB^{-1}$$

$$\Rightarrow A = CB^{-1} \quad [\because BB^{-1} = I]$$

$$|B| = \begin{vmatrix} 5 & 3 \\ -1 & -2 \end{vmatrix} = -10 + 3 = -7 \neq 0$$

$\therefore B^{-1}$ exists

$$B^{-1} = \frac{1}{|B|} \text{adj } B = \frac{-1}{7} \begin{bmatrix} -2 & -3 \\ 1 & 5 \end{bmatrix}$$

$$A = CB^{-1}$$

$$= \begin{bmatrix} 14 & 7 \\ 7 & 7 \end{bmatrix} \cdot \left(\frac{-1}{7}\right) \begin{bmatrix} -2 & -3 \\ 1 & 5 \end{bmatrix}$$

$$= 7 \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \left(\frac{-1}{7}\right) \begin{bmatrix} -2 & -3 \\ 1 & 5 \end{bmatrix}$$

$$= - \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} -2 & -3 \\ 1 & 5 \end{bmatrix} = - \begin{bmatrix} -4+1 & -6+5 \\ -2+1 & -3+5 \end{bmatrix}$$

$$= - \begin{bmatrix} -3 & -1 \\ -1 & 2 \end{bmatrix} \Rightarrow A = \begin{bmatrix} 3 & 1 \\ 1 & -2 \end{bmatrix}$$

13. Given $A = \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 3 & -2 \\ 1 & 1 \end{bmatrix}$ and

$C = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$, find a matrix X such that $AXB = C$.

Sol. Given $A = \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 3 & -2 \\ 1 & 1 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$

$$\text{Also, } A X B = C$$

Pre-multiply by A^{-1} we get,

$$(A^{-1} A) X B = A^{-1} C$$

$$\Rightarrow X B = A^{-1} \cdot C \quad [\because A^{-1} A = I]$$

Post Multiply by B^{-1} we get

$$(X B) B^{-1} = (A^{-1} C) B^{-1} \Rightarrow X = (A^{-1} C) B^{-1}$$

$$|A| = \begin{vmatrix} 1 & -1 \\ 2 & 0 \end{vmatrix} = 0 + 2 = 2 \neq 0$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{2} \begin{bmatrix} 0 & 1 \\ -2 & 1 \end{bmatrix}$$

$$|B| = \begin{vmatrix} 3 & -2 \\ 1 & 1 \end{vmatrix} = 3 + 2 = 5 \neq 0$$

$$\therefore B^{-1} = \frac{1}{|B|} \text{adj } B = \frac{1}{5} \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix}$$

$$A^{-1} C = \frac{1}{2} \begin{bmatrix} 0 & 1 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} 0+2 & 0+2 \\ -2+2 & -2+2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 2 & 2 \\ 0 & 0 \end{bmatrix}$$

$$= \frac{1}{2} (2) \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\therefore X = (A^{-1} C) \cdot B^{-1}$$

$$= \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \frac{1}{5} \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix}$$

$$= \frac{1}{5} \begin{bmatrix} 1-1 & 2+3 \\ 0+0 & 0+0 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 0 & 5 \\ 0 & 0 \end{bmatrix}$$

$$= \frac{1}{5} (5) \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\therefore X = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

14. If $A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$, show that $A^{-1} = \frac{1}{2}(A^2 - 3I)$.

Sol. Given $A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$

$$|A| = 0 - 1 \begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} + 1 \begin{vmatrix} 1 & 0 \\ 1 & 1 \end{vmatrix}$$

$$= -1(0-1) + 1(1-0) = 1 + 1 = 2$$

$$|A| = 2 \neq 0,$$

Hence A^{-1} exists

$$\begin{aligned} \text{adj } A &= \begin{bmatrix} + \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} & - \begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} & + \begin{vmatrix} 1 & 0 \\ 1 & 1 \end{vmatrix} \\ - \begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} & + \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} & - \begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix} \\ + \begin{vmatrix} 1 & 1 \\ 0 & 1 \end{vmatrix} & - \begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix} & + \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} \end{bmatrix}^T \\ &= \begin{bmatrix} (0-1) & -(0-1) & +(1-0) \\ -(0-1) & +(0-1) & -(0-1) \\ +(1-0) & -(0-1) & +(0-1) \end{bmatrix}^T \\ &= \begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix}^T = \begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix} \\ \therefore A^{-1} &= \frac{1}{2} \begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix} \quad \dots(1) \end{aligned}$$

$$\begin{aligned} \text{Now } A^2 &= \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0+1+1 & 0+0+1 & 0+1+0 \\ 0+0+1 & 1+0+1 & 1+0+0 \\ 0+1+0 & 1+0+0 & 1+1+0 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix} \\ A^2 - 3I &= \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix} - 3 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 2-3 & 1-0 & 1-0 \\ 1-0 & 2-3 & 1-0 \\ 1-0 & 1-0 & 2-3 \end{bmatrix} \\ &= \frac{1}{2} (A^2 - 3I) = \frac{1}{2} \begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix} \quad \dots(2) \end{aligned}$$

From (1) and (2), it is proved that $A^{-1} = \frac{1}{2} [A^2 - 3I]$

15. Decrypt the received encoded message $[2 \ -3] \ [20 \ 4]$ with the encryption matrix $\begin{bmatrix} -1 & -1 \\ 2 & 1 \end{bmatrix}$ and the decryption matrix as its inverse, where the system of codes are described by the numbers 1 – 26 to the letters A – Z respectively, and the number 0 to a blank space.

Sol. Let the encryption matrix be $A = \begin{bmatrix} -1 & -1 \\ 2 & 1 \end{bmatrix}$

$$|A| = -1 + 2 = 1 \neq 0$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{1} \begin{bmatrix} 1 & 1 \\ -2 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ -2 & -1 \end{bmatrix}$$

Hence the decryption matrix is $\begin{bmatrix} 1 & 1 \\ -2 & -1 \end{bmatrix}$

Coded row matrix	Decoding matrix	Decoded row matrix
$[2 \ -3]$	$\begin{bmatrix} 1 & 1 \\ -2 & -1 \end{bmatrix}$	$= [2+6 \ 2+3] = [8 \ 5]$
$[20 \ 4]$	$\begin{bmatrix} 1 & 1 \\ -2 & -1 \end{bmatrix}$	$= [20-8 \ 20-4] = [12 \ 16]$

So, the sequence of decoded row matrices is $[8 \ 5], [12 \ 16]$

Now the 8th English alphabet is H.

5th English alphabet is E.

12th English alphabet is L.

and the 16th English alphabet is P.

Thus the receiver reads the message as “HELP”.

EXERCISE 1.2

1. Find the rank of the following matrices by minor method:

(i) $\begin{bmatrix} 2 & -4 \\ -1 & 2 \end{bmatrix}$ (ii) $\begin{bmatrix} -1 & 3 \\ 4 & -7 \\ 3 & -4 \end{bmatrix}$

[Mar. - 2024]

(iii) $\begin{bmatrix} 1 & -2 & -1 & 0 \\ 3 & -6 & -3 & 1 \end{bmatrix}$ [PTA - 5]

(iv) $\begin{bmatrix} 1 & -2 & 3 \\ 2 & 4 & -6 \\ 5 & 1 & -1 \end{bmatrix}$ [July - 2022]

(v) $\begin{bmatrix} 0 & 1 & 2 & 1 \\ 0 & 2 & 4 & 3 \\ 8 & 1 & 0 & 2 \end{bmatrix}$ [Aug. - 2021]

Sol. (i) Let $A = \begin{bmatrix} 2 & -4 \\ -1 & 2 \end{bmatrix}$

A is a matrix of order 2×2

$$\therefore \rho(A) \leq \min(2, 2) = 2$$

The highest order of minor of A is 2

$$\text{It is } \begin{vmatrix} 2 & -4 \\ -1 & 2 \end{vmatrix} = 4 - 4 = 0$$

$$\text{So, } \rho(A) < 2$$

Next consider the minor of order 1 $|2| = 2 \neq 0$

$$\therefore \rho(A) = 1$$

(ii) Let $A = \begin{bmatrix} -1 & 3 \\ 4 & -7 \\ 3 & -4 \end{bmatrix}$

A is a matrix of order 3×2

$$\therefore \rho(A) \leq 2$$

We find that there is a second order minor,

$$\begin{vmatrix} -1 & 3 \\ 4 & -7 \end{vmatrix} = 7 - 12 = -5 \neq 0$$

$$\therefore \rho(A) = 2.$$

(iii) Let $A = \begin{bmatrix} 1 & -2 & -1 & 0 \\ 3 & -6 & -3 & 1 \end{bmatrix}$

A is a matrix of order (2×4)

$$\therefore \rho(A) \leq \min(2, 4) = 2$$

The highest order of minor of A is 2

$$\text{It is } \begin{vmatrix} 1 & -2 \\ 3 & -6 \end{vmatrix} = -6 + 6 = 0$$

$$\text{Also, } \begin{vmatrix} -1 & 0 \\ -3 & 1 \end{vmatrix} = -1 + 0 = -1 \neq 0.$$

$$\therefore \rho(A) = 2.$$

(iv) Let $A = \begin{bmatrix} 1 & -2 & 3 \\ 2 & 4 & -6 \\ 5 & 1 & -1 \end{bmatrix}$

A is a matrix of order 3×3

$$\therefore \rho(A) \leq \min(3, 3) = 3$$

The highest order of minor of A is 3.

$$(I.E) \begin{vmatrix} 1 & -2 & 3 \\ 2 & 4 & -6 \\ 5 & 1 & -1 \end{vmatrix} = 1 \begin{vmatrix} 4 & -6 \\ 1 & -1 \end{vmatrix} + 2 \begin{vmatrix} 2 & -6 \\ 5 & -1 \end{vmatrix} + 3 \begin{vmatrix} 2 & 4 \\ 5 & 1 \end{vmatrix}$$

[Expanded along R_1]

$$= 1(-4 + 6) + 2(-2 + 30) + 3(2 - 20)$$

$$= 1(2) + 2(28) + 3(-18)$$

$$= 2 + 56 - 54 = 58 - 54 = 4 \neq 0$$

$$\therefore \rho(A) = 3.$$

(v) Let $A = \begin{bmatrix} 0 & 1 & 2 & 1 \\ 0 & 2 & 4 & 3 \\ 8 & 1 & 0 & 2 \end{bmatrix}$

A is a matrix of order 3×4

$$\therefore \rho(A) \leq \min(3, 4) = 3$$

The highest order of minor of A is 3

$$(I.E) \begin{vmatrix} 0 & 1 & 2 \\ 0 & 2 & 4 \\ 8 & 1 & 0 \end{vmatrix} = 0 + 0 + 8(4 - 4) = 0$$

[Expanded along C_1]

$$\text{Also, } \begin{vmatrix} 0 & 2 & 1 \\ 0 & 4 & 3 \\ 8 & 0 & 2 \end{vmatrix} = 0 + 0 - 8 \begin{vmatrix} 2 & 1 \\ 4 & 3 \end{vmatrix}$$

[Expanded along C_1]

$$= -8(6 - 4) = -8(2) = -16 \neq 0$$

$$\therefore \rho(A) = 3$$

2. Find the rank of the following matrices by row reduction method :

(i) $\begin{bmatrix} 1 & 1 & 1 & 3 \\ 2 & -1 & 3 & 4 \\ 5 & -1 & 7 & 11 \end{bmatrix}$ [PTA-1; FRT -2022]

(ii) $\begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 1 & -2 & 3 \\ 1 & -1 & 1 \end{bmatrix}$ [May - 2022; Qy. - 2024]

(iii) $\begin{bmatrix} 3 & -8 & 5 & 2 \\ 2 & -5 & 1 & 4 \\ -1 & 2 & 3 & -2 \end{bmatrix}$ [July - 2024]

Sol. (i) Let $A = \begin{bmatrix} 1 & 1 & 1 & 3 \\ 2 & -1 & 3 & 4 \\ 5 & -1 & 7 & 11 \end{bmatrix}$

$$A = \begin{bmatrix} 1 & 1 & 1 & 3 \\ 2 & -1 & 3 & 4 \\ 5 & -1 & 7 & 11 \end{bmatrix} \xrightarrow{R_2 \rightarrow R_2 - 2R_1} \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & -3 & 1 & -2 \\ 5 & -1 & 7 & 11 \end{bmatrix}$$

$$\xrightarrow{R_3 \rightarrow R_3 - 5R_1} \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & -3 & 1 & -2 \\ 0 & -6 & 2 & -4 \end{bmatrix}$$

$$\xrightarrow{R_3 \rightarrow R_3 - 2R_2} \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & -3 & 1 & -2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

The last equivalent matrix is in row echelon form
it has two non-zero rows

$$\therefore \rho(A) = 2.$$

(ii) Let $A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 1 & -2 & 3 \\ 1 & -1 & 1 \end{bmatrix}$

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 1 & -2 & 3 \\ 1 & -1 & 1 \end{bmatrix} \xrightarrow{\begin{matrix} R_2 \rightarrow R_2 - 3R_1 \\ R_3 \rightarrow R_3 - R_1 \\ R_4 \rightarrow R_4 - R_1 \end{matrix}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & -4 & 4 \\ 0 & -3 & 2 \end{bmatrix}$$

$$\begin{array}{l}
 R_3 \rightarrow R_3 \div 4 \rightarrow \begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & -1 & 1 \\ 0 & -3 & 2 \end{bmatrix} \xrightarrow{R_4 \rightarrow R_4 - 3R_3} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & -1 & 1 \\ 0 & 0 & -1 \end{bmatrix} \\
 R_3 \rightarrow 7R_3 - R_2 \rightarrow \begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & 0 & 2 \\ 0 & 0 & -1 \end{bmatrix} \xrightarrow{R_4 \rightarrow 2R_4 + R_3} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{bmatrix}
 \end{array}$$

The last equivalent matrix is in row echelon form
it has two non-zero rows

$$\therefore \rho(A) = 3$$

(iii) Let $A = \begin{bmatrix} 3 & -8 & 5 & 2 \\ 2 & -5 & 1 & 4 \\ -1 & 2 & 3 & -2 \end{bmatrix}$

$$\begin{array}{l}
 A \xrightarrow{R_3 \leftrightarrow R_1} \begin{bmatrix} -1 & 2 & 3 & -2 \\ 2 & -5 & 1 & 4 \\ 3 & -8 & 5 & 2 \end{bmatrix} \\
 \xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 + 2R_1 \\ R_3 \rightarrow R_3 + 3R_1 \end{array}} \begin{bmatrix} -1 & 2 & 3 & -2 \\ 0 & -1 & 7 & 0 \\ 0 & -2 & 14 & -4 \end{bmatrix} \\
 \xrightarrow{R_3 \rightarrow R_3 \div 2} \begin{bmatrix} -1 & 2 & 3 & -2 \\ 0 & -1 & 7 & 0 \\ 0 & -1 & 7 & -2 \end{bmatrix} \\
 \xrightarrow{R_3 \rightarrow R_3 - R_2} \begin{bmatrix} -1 & 2 & 3 & -2 \\ 0 & -1 & 7 & 0 \\ 0 & 0 & 0 & -2 \end{bmatrix}
 \end{array}$$

The last equivalent matrix is in row-echelon form. It has three non-zero rows.

$$\therefore \rho(A) = 3$$

3. Find the inverse of each of the following by Gauss – Jordan method :

(i) $\begin{bmatrix} 2 & -1 \\ 5 & -2 \end{bmatrix}$ [Govt. MQP-2019; Hy. - 2024]

(ii) $\begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \\ 6 & -2 & -3 \end{bmatrix}$ (iii) $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 3 \\ 1 & 0 & 8 \end{bmatrix}$

Sol. (i) Let $A = \begin{bmatrix} 2 & -1 \\ 5 & -2 \end{bmatrix}$

Applying Gauss – Jordan method, we get

$$\begin{array}{l}
 [A|I_2] = \left[\begin{array}{cc|cc} 2 & -1 & 1 & 0 \\ 5 & -2 & 0 & 1 \end{array} \right] \\
 \xrightarrow{R_1 \rightarrow R_1 \div 2} \left[\begin{array}{cc|cc} 1 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 5 & -2 & 0 & 1 \end{array} \right]
 \end{array}$$

$$\xrightarrow{R_2 \rightarrow R_2 - 5R_1} \left[\begin{array}{cc|cc} 1 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & \frac{1}{2} & -\frac{5}{2} & 1 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 \times 2} \left[\begin{array}{cc|cc} 1 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 1 & -5 & 2 \end{array} \right]$$

$$\xrightarrow{R_1 \rightarrow R_1 + \frac{1}{2}R_2} \left[\begin{array}{cc|cc} 1 & 0 & -2 & 1 \\ 0 & 1 & -5 & 2 \end{array} \right]$$

$$\therefore \text{We get } A^{-1} = \begin{bmatrix} -2 & 1 \\ -5 & 2 \end{bmatrix}$$

(ii) Let $A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \\ 6 & -2 & -3 \end{bmatrix}$

Applying Gauss – Jordan method, we get

$$[A|I_3] = \left[\begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 1 & 0 & -1 & 0 & 1 & 0 \\ 6 & -2 & -3 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 - R_1} \left[\begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 6 & -2 & -3 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - 6R_1} \left[\begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 4 & -3 & -6 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - 4R_2} \left[\begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -2 & -4 & 1 \end{array} \right]$$

$$\xrightarrow{R_1 \rightarrow R_1 + R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & -1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -2 & -4 & 1 \end{array} \right]$$

$$\xrightarrow{R_1 \rightarrow R_1 + R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & -3 & 1 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -2 & -4 & 1 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 + R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & -3 & 1 \\ 0 & 1 & 0 & -3 & -3 & 1 \\ 0 & 0 & 1 & -2 & -4 & 1 \end{array} \right]$$

$$\text{So, we get } A^{-1} = \begin{bmatrix} -2 & -3 & 1 \\ -3 & -3 & 1 \\ -2 & -4 & 1 \end{bmatrix}$$

$$(iii) \text{ Let } A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 3 \\ 1 & 0 & 8 \end{bmatrix}$$

Applying Gauss Jordan method, we get

$$[A|I_3] = \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 5 & 3 & 0 & 1 & 0 \\ 1 & 0 & 8 & 0 & 0 & 1 \end{array} \right]$$

$$\begin{array}{l} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - R_1 \end{array} \rightarrow \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & -3 & -2 & 1 & 0 \\ 0 & -2 & 5 & -1 & 0 & 1 \end{array} \right]$$

$$R_1 \rightarrow R_1 + R_3 \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 8 & 0 & 0 & 1 \\ 0 & 1 & -3 & -2 & 1 & 0 \\ 0 & -2 & 5 & -1 & 0 & 1 \end{array} \right]$$

$$R_3 \rightarrow R_3 + 2R_2 \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 8 & 0 & 0 & 1 \\ 0 & 1 & -3 & -2 & 1 & 0 \\ 0 & 0 & -1 & -5 & 2 & 1 \end{array} \right]$$

$$R_2 \rightarrow R_2 - 3R_3 \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 8 & 0 & 0 & 1 \\ 0 & 1 & 0 & 13 & -5 & -3 \\ 0 & 0 & -1 & -5 & 2 & 1 \end{array} \right]$$

$$R_1 \rightarrow R_1 + 8R_3 \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -40 & 16 & 9 \\ 0 & 1 & 0 & 13 & -5 & -3 \\ 0 & 0 & -1 & -5 & 2 & 1 \end{array} \right]$$

$$R_3 \rightarrow R_3 \times (-1) \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -40 & 16 & 9 \\ 0 & 1 & 0 & 13 & -5 & -3 \\ 0 & 0 & 1 & 5 & -2 & -1 \end{array} \right]$$

$$\text{So, we get } A^{-1} = \begin{bmatrix} -40 & 16 & 9 \\ 13 & -5 & -3 \\ 5 & -2 & -1 \end{bmatrix}$$

EXERCISE 1.3

1. Solve the following system of linear equations by matrix inversion method:

(i) $2x + 5y = -2, x + 2y = -3$

[June - 2023; Mar. - 2025]

(ii) $2x - y = 8, 3x + 2y = -2$ [PTA -3; June - 2025]

(iii) $2x + 3y - z = 9, x + y + z = 9, 3x - y - z = -1$

[July - 2024]

(iv) $x + y + z - 2 = 0, 6x - 4y + 5z - 31 = 0,$
 $5x + 2y + 2z = 13$

• Sol. (i) $2x + 5y = -2, x + 2y = -3$

The matrix form of the system is

$$= \begin{pmatrix} 2 & 5 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -2 \\ -3 \end{pmatrix}$$

$$\Rightarrow AX = B \text{ where}$$

$$A = \begin{pmatrix} 2 & 5 \\ 1 & 2 \end{pmatrix}, B = \begin{pmatrix} -2 \\ -3 \end{pmatrix},$$

$$X = \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\Rightarrow X = A^{-1} B$$

$$|A| = \begin{vmatrix} 2 & 5 \\ 1 & 2 \end{vmatrix} = 4 - 5 = -1 \neq 0.$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$= \frac{1}{-1} \begin{bmatrix} 2 & -5 \\ -1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} -2 & 5 \\ 1 & -2 \end{bmatrix}$$

$$\therefore X = A^{-1} B = \begin{bmatrix} -2 & 5 \\ 1 & -2 \end{bmatrix} \begin{bmatrix} -2 \\ -3 \end{bmatrix}$$

$$= \begin{bmatrix} 4 - 15 \\ -2 + 6 \end{bmatrix} = \begin{bmatrix} -11 \\ 4 \end{bmatrix}$$

$$\therefore x = -11, y = 4.$$

(ii) $2x - y = 8, 3x + 2y = -2$

The matrix form of the system is

$$\begin{bmatrix} 2 & -1 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ -2 \end{bmatrix}$$

$$\Rightarrow AX = B \text{ where } A = \begin{bmatrix} 2 & -1 \\ 3 & 2 \end{bmatrix},$$

$$B = \begin{bmatrix} 8 \\ -2 \end{bmatrix}$$

$$\Rightarrow X = A^{-1} B.$$

$$\text{Now, } |A| = \begin{vmatrix} 2 & -1 \\ 3 & 2 \end{vmatrix} = 4 + 3 = 7$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$= \frac{1}{7} \begin{bmatrix} 2 & 1 \\ -3 & 2 \end{bmatrix}$$

$$\therefore X = A^{-1} B = \frac{1}{7} \begin{bmatrix} 2 & 1 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 8 \\ -2 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 16 - 2 \\ -24 - 4 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 14 \\ -28 \end{bmatrix} = \begin{bmatrix} \frac{14}{7} \\ \frac{-28}{7} \end{bmatrix} = \begin{bmatrix} 2 \\ -4 \end{bmatrix}$$

$$\therefore x = 2, y = -4$$

(iii) $2x + 3y - z = 9, x + y + z = 9, 3x - y - z = -1.$

The matrix form of the system is

$$\begin{bmatrix} 2 & 3 & -1 \\ 1 & 1 & 1 \\ 3 & -1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 9 \\ 9 \\ -1 \end{bmatrix}$$

$$\Rightarrow AX = B \text{ where } A = \begin{bmatrix} 2 & 3 & -1 \\ 1 & 1 & 1 \\ 3 & -1 & -1 \end{bmatrix},$$

$$X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \text{ and } B = \begin{bmatrix} 9 \\ 9 \\ -1 \end{bmatrix}$$

$$\Rightarrow X = A^{-1} B$$

$$|A| = \begin{vmatrix} 2 & 3 & -1 \\ 1 & 1 & 1 \\ 3 & -1 & -1 \end{vmatrix} = 2 \begin{vmatrix} 1 & 1 \\ -1 & -1 \end{vmatrix} - 3 \begin{vmatrix} 1 & 1 \\ 3 & -1 \end{vmatrix} - 1 \begin{vmatrix} 1 & 1 \\ 3 & -1 \end{vmatrix}$$

[Expanded along R_1]

$$= 2(-1+1) - 3(-1-3) - 1(-1-3)$$

$$= 0 - 3(-4) - 1(-4) = 12 + 4 = 16.$$

$$\text{adj } A = \begin{bmatrix} + \begin{vmatrix} 1 & 1 \\ -1 & -1 \end{vmatrix} & - \begin{vmatrix} 1 & 1 \\ 3 & -1 \end{vmatrix} & + \begin{vmatrix} 1 & 1 \\ 3 & -1 \end{vmatrix} \\ - \begin{vmatrix} 3 & -1 \\ -1 & -1 \end{vmatrix} & + \begin{vmatrix} 2 & -1 \\ 3 & -1 \end{vmatrix} & - \begin{vmatrix} 2 & 3 \\ 3 & -1 \end{vmatrix} \\ + \begin{vmatrix} 3 & -1 \\ 1 & 1 \end{vmatrix} & - \begin{vmatrix} 2 & -1 \\ 1 & 1 \end{vmatrix} & + \begin{vmatrix} 2 & 3 \\ 1 & 1 \end{vmatrix} \end{bmatrix}^T$$

$$= \begin{bmatrix} +(-1+1) & -(-1-3) & +(-1-3) \\ -(-3-1) & +(-2+3) & -(-2-9) \\ +(3+1) & -(2+1) & +(2-3) \end{bmatrix}^T$$

$$= \begin{bmatrix} 0 & 4 & -4 \\ 4 & 1 & 11 \\ 4 & -3 & -1 \end{bmatrix}^T = \begin{bmatrix} 0 & 4 & 4 \\ 4 & 1 & -3 \\ -4 & 11 & -1 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{16} \begin{bmatrix} 0 & 4 & 4 \\ 4 & 1 & -3 \\ -4 & 11 & -1 \end{bmatrix}$$

$$\therefore X = A^{-1} B = \frac{1}{16} \begin{bmatrix} 0 & 4 & 4 \\ 4 & 1 & -3 \\ -4 & 11 & -1 \end{bmatrix} \begin{bmatrix} 9 \\ 9 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{16} \begin{bmatrix} 0+36-4 \\ 36+9+3 \\ -36+99+1 \end{bmatrix} = \frac{1}{16} \begin{bmatrix} 32 \\ 48 \\ 64 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}$$

$$\therefore x = 2, y = 3, z = 4$$

(iv) $x + y + z - 2 = 0, 6x - 4y + 5z - 31 = 0, 5x + 2y + 2z = 13$

The matrix form of the system is

$$\begin{bmatrix} 1 & 1 & 1 \\ 6 & -4 & 5 \\ 5 & 2 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 31 \\ 13 \end{bmatrix}$$

$$AX = B \text{ where } A = \begin{bmatrix} 1 & 1 & 1 \\ 6 & -4 & 5 \\ 5 & 2 & 2 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 2 \\ 31 \\ 13 \end{bmatrix}$$

$$\Rightarrow X = A^{-1} B$$

$$|A| = \begin{vmatrix} 1 & 1 & 1 \\ 6 & -4 & 5 \\ 5 & 2 & 2 \end{vmatrix} = 1 \begin{vmatrix} -4 & 5 \\ 2 & 2 \end{vmatrix} - 1 \begin{vmatrix} 6 & 5 \\ 5 & 2 \end{vmatrix} + 1 \begin{vmatrix} 6 & -4 \\ 5 & 2 \end{vmatrix}$$

$$= 1(-8-10) - 1(12-25) + 1(12+20)$$

$$= 1(-18) - 1(-13) + 1(32) = -18 + 13 + 32 = 27$$

$$\text{adj } A = \begin{bmatrix} + \begin{vmatrix} -4 & 5 \\ 2 & 2 \end{vmatrix} & - \begin{vmatrix} 6 & 5 \\ 5 & 2 \end{vmatrix} & + \begin{vmatrix} 6 & -4 \\ 5 & 2 \end{vmatrix} \\ - \begin{vmatrix} 1 & 1 \\ 2 & 2 \end{vmatrix} & + \begin{vmatrix} 1 & 1 \\ 5 & 2 \end{vmatrix} & - \begin{vmatrix} 1 & 1 \\ 5 & 2 \end{vmatrix} \\ + \begin{vmatrix} 1 & 1 \\ -4 & 5 \end{vmatrix} & - \begin{vmatrix} 1 & 1 \\ 6 & 5 \end{vmatrix} & + \begin{vmatrix} 1 & 1 \\ 6 & -4 \end{vmatrix} \end{bmatrix}^T$$

$$= \begin{bmatrix} +(-8-10) & -(12-25) & +(12+20) \\ -(2-2) & +(2-5) & -(2-5) \\ +(5+4) & -(5-6) & +(-4-6) \end{bmatrix}^T$$

$$= \begin{bmatrix} -18 & 13 & 32 \\ 0 & -3 & 3 \\ 9 & 1 & -10 \end{bmatrix}^T = \begin{bmatrix} -18 & 0 & 9 \\ 13 & -3 & 1 \\ 32 & 3 & -10 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{27} \begin{bmatrix} -18 & 0 & 9 \\ 13 & -3 & 1 \\ 32 & 3 & -10 \end{bmatrix}$$

$$\therefore X = A^{-1} B = \frac{1}{27} \begin{bmatrix} -18 & 0 & 9 \\ 13 & -3 & 1 \\ 32 & 3 & -10 \end{bmatrix} \begin{bmatrix} 2 \\ 31 \\ 13 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{27} \begin{bmatrix} -36+0+117 \\ 26-93+13 \\ 64+93-130 \end{bmatrix} = \frac{1}{27} \begin{bmatrix} 81 \\ -54 \\ 27 \end{bmatrix} = \begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix}$$

$$\therefore x=3, y=-2, z=1$$

2. If $A = \begin{bmatrix} -5 & 1 & 3 \\ 7 & 1 & -5 \\ 1 & -1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$, find

the products AB and BA and hence solve the system of equations $x+y+2z=1$, $3x+2y+z=7$, $2x+y+3z=2$. [Qy. - 2023]

Sol. Given $A = \begin{bmatrix} -5 & 1 & 3 \\ 7 & 1 & -5 \\ 1 & -1 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$

$$AB = \begin{bmatrix} -5 & 1 & 3 \\ 7 & 1 & -5 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} -5+3+6 & -5+2+3 & -10+1+9 \\ 7+3-10 & 7+2-5 & 14+1-15 \\ 1-3+2 & 1-2+1 & 2-1+3 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix} = 4 \cdot I_3$$

$$BA = \begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix} \begin{bmatrix} -5 & 1 & 3 \\ 7 & 1 & -5 \\ 1 & -1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} -5+7+2 & 1+1-2 & 3-5+2 \\ -15+14+1 & 3+2-1 & 9-10+1 \\ -10+7+3 & 2+1-3 & 6-5+3 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix} = 4 \cdot I_3$$

So, we get $AB = BA = 4 \cdot I_3$

$$\Rightarrow \left(\frac{1}{4}A\right) B = B \left(\frac{1}{4}A\right) = I$$

$$\Rightarrow B^{-1} = \frac{1}{4} A$$

Writing the given set of equations in matrix form we get,

$$\begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 7 \\ 2 \end{bmatrix}$$

$$\Rightarrow B \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 7 \\ 2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = B^{-1} \begin{bmatrix} 1 \\ 7 \\ 2 \end{bmatrix} = \left[\frac{1}{4}A\right] \begin{bmatrix} 1 \\ 7 \\ 2 \end{bmatrix}$$

$$= \frac{1}{4} \begin{bmatrix} -5 & 1 & 3 \\ 7 & 1 & -5 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 7 \\ 2 \end{bmatrix} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{4} \begin{bmatrix} -5+7+6 \\ 7+7-10 \\ 1-7+2 \end{bmatrix}$$

$$= \frac{1}{4} \begin{bmatrix} 8 \\ 4 \\ -4 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix}$$

$$\therefore x=2, y=1, z=-1$$

3. A man is appointed in a job with a monthly salary of certain amount and a fixed amount of annual increment. If his salary was ₹ 19,800 per month at the end of the first month after 3 years of service and ₹ 23,400 per month at the end of the first month after 9 years of service, find his starting salary and his annual increment. (Use matrix inversion method to solve the problem.)

Sol. Let the man's starting salary be ₹ x and his annual increment be ₹ y .

By the given data $x+3y=19800$ and $x+9y=23,400$.

The matrix form of the given system of equations is

$$\begin{bmatrix} 1 & 3 \\ 1 & 9 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 19800 \\ 23400 \end{bmatrix}$$

$$\Rightarrow AX = B \text{ where } A = \begin{bmatrix} 1 & 3 \\ 1 & 9 \end{bmatrix} \text{ and } B = \begin{bmatrix} 19800 \\ 23400 \end{bmatrix}$$

$$\Rightarrow X = A^{-1} B$$

$$|A| = \begin{vmatrix} 1 & 3 \\ 1 & 9 \end{vmatrix}$$

$$= 9-3=6 \neq 0$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj}A = \frac{1}{6} \begin{bmatrix} 9 & -3 \\ -1 & 1 \end{bmatrix}$$

$$\therefore X = A^{-1} B$$

$$= \frac{1}{6} \begin{bmatrix} 9 & -3 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 19800 \\ 23400 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{6} \begin{bmatrix} 178200 - 70200 \\ -19800 + 23400 \end{bmatrix} = \frac{1}{6} \begin{bmatrix} 108000 \\ 3600 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 18000 \\ 600 \end{bmatrix}$$

∴ $x = 18000, y = 600$.

Hence the man's starting salary is ₹ 18000 and his annual increment is ₹ 600.

4. Four men and 4 women can finish a piece of work jointly in 3 days while 2 men and 5 women can finish the same work jointly in 4 days. Find the time taken by one man alone and that of one woman alone to finish the same work by using matrix inversion method.

Sol. Let the time by one man alone be x days and one woman alone be y days

∴ By the given data,

$$\frac{4}{x} + \frac{4}{y} = \frac{1}{3} \text{ and } \frac{2}{x} + \frac{5}{y} = \frac{1}{4}$$

$$\text{put } \frac{1}{x} = s \text{ and } \frac{1}{y} = t$$

$$\therefore 4s + 4t = \frac{1}{3} \text{ and } 2s + 5t = \frac{1}{4}$$

The matrix form of the system of equation is

$$\begin{bmatrix} 4 & 4 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} s \\ t \end{bmatrix} = \begin{bmatrix} \frac{1}{3} \\ \frac{1}{4} \end{bmatrix} \Rightarrow AX = B \text{ where}$$

$$A = \begin{bmatrix} 4 & 4 \\ 2 & 5 \end{bmatrix} \text{ and } B = \begin{bmatrix} \frac{1}{3} \\ \frac{1}{4} \end{bmatrix}$$

$$X = A^{-1} B$$

$$\text{Now } |A| = \begin{vmatrix} 4 & 4 \\ 2 & 5 \end{vmatrix} = 20 - 8 = 12 \neq 0$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{12} \begin{bmatrix} 5 & -4 \\ -2 & 4 \end{bmatrix}$$

$$\therefore X = A^{-1} B = \frac{1}{12} \begin{bmatrix} 5 & -4 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} \frac{1}{3} \\ \frac{1}{4} \end{bmatrix} = \frac{1}{12} \begin{bmatrix} \frac{5}{3} - 1 \\ -\frac{2}{3} + 1 \end{bmatrix}$$

$$= \frac{1}{12} \begin{bmatrix} \frac{2}{3} \\ \frac{1}{3} \end{bmatrix} = \begin{bmatrix} \frac{2}{3} \times \frac{1}{12} \\ \frac{1}{3} \times \frac{1}{12} \end{bmatrix} = \begin{bmatrix} \frac{1}{18} \\ \frac{1}{36} \end{bmatrix}$$

$$\therefore s = \frac{1}{18} \Rightarrow \frac{1}{x} = \frac{1}{18}$$

$$\Rightarrow x = 18$$

$$t = \frac{1}{36} \Rightarrow \frac{1}{y} = \frac{1}{36}$$

$$\Rightarrow y = 36.$$

Hence, the time taken by 1 man alone is 18 days and the time taken by 1 woman alone is 36 days.

5. The prices of three commodities A, B and C are ₹ x, y and ₹ z per units respectively. A person P purchases 4 units of B and sells two units of A and 5 units of C. Person Q purchases 2 units of C and sells 3 units of A and one unit of B. Person R purchases one unit of A and sells 3 unit of B and one unit of C. In the process, P, Q and R earn ₹ 15,000, ₹ 1,000 and ₹ 4,000 respectively. Find the prices per unit of A, B and C. (Use matrix inversion method to solve the problem.) [Qy. - 2024]

Sol. Let the prices per unit for the commodities A, B and C be ₹ x, y and ₹ z .

By the given data,

$$2x - 4y + 5z = 15000$$

$$3x + y - 2z = 1000$$

$$-x + 3y + z = 4000$$

The matrix form of the system of equations is

$$\begin{bmatrix} 2 & -4 & 5 \\ 3 & 1 & -2 \\ -1 & 3 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 15000 \\ 1000 \\ 4000 \end{bmatrix}$$

$$AX = B \text{ where } A = \begin{bmatrix} 2 & -4 & 5 \\ 3 & 1 & -2 \\ -1 & 3 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix},$$

$$\text{and } B = \begin{bmatrix} 15000 \\ 1000 \\ 4000 \end{bmatrix} \Rightarrow X = A^{-1} B$$

$$|A| = \begin{vmatrix} 2 & -4 & 5 \\ 3 & 1 & -2 \\ -1 & 3 & 1 \end{vmatrix}$$

$$= 2 \begin{vmatrix} 1 & -2 \\ 3 & 1 \end{vmatrix} + 4 \begin{vmatrix} 3 & -2 \\ -1 & 1 \end{vmatrix} + 5 \begin{vmatrix} 3 & 1 \\ -1 & 3 \end{vmatrix}$$

$$= 2(1+6) + 4(3-2) + 5(9+1)$$

$$= 2(7) + 4(1) + 5(10) = 14 + 4 + 50 = 68.$$

$$\text{adj } A = \begin{bmatrix} + \begin{vmatrix} 1 & -2 \\ 3 & 1 \end{vmatrix} & - \begin{vmatrix} 3 & -2 \\ -1 & 1 \end{vmatrix} & + \begin{vmatrix} 3 & 1 \\ -1 & 3 \end{vmatrix} \\ - \begin{vmatrix} -4 & 5 \\ 3 & 1 \end{vmatrix} & + \begin{vmatrix} 2 & 5 \\ -1 & 1 \end{vmatrix} & - \begin{vmatrix} 2 & -4 \\ -1 & 3 \end{vmatrix} \\ + \begin{vmatrix} -4 & 5 \\ 1 & -2 \end{vmatrix} & - \begin{vmatrix} 2 & 5 \\ 3 & -2 \end{vmatrix} & + \begin{vmatrix} 2 & -4 \\ 3 & 1 \end{vmatrix} \end{bmatrix}^T$$

$$= \begin{bmatrix} +(1+6) & -(3-2) & +(9+1) \\ -(-4-15) & +(2+5) & -(6-4) \\ +(8-5) & -(-4-15) & +(2+12) \end{bmatrix}^T$$

$$= \begin{bmatrix} 7 & -1 & 10 \\ 19 & 7 & -2 \\ 3 & 19 & 14 \end{bmatrix}^T = \begin{bmatrix} 7 & 19 & 3 \\ -1 & 7 & 19 \\ 10 & -2 & 14 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{68} \begin{bmatrix} 7 & 19 & 3 \\ -1 & 7 & 19 \\ 10 & -2 & 14 \end{bmatrix}$$

$$\therefore X = A^{-1} B = \frac{1}{68} \begin{bmatrix} 7 & 19 & 3 \\ -1 & 7 & 19 \\ 10 & -2 & 14 \end{bmatrix} \begin{bmatrix} 15000 \\ 1000 \\ 4000 \end{bmatrix}$$

$$= \frac{1}{68} \begin{bmatrix} 105000 + 19000 + 12000 \\ -15000 + 7000 + 76000 \\ 150000 - 2000 + 56000 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{68} \begin{bmatrix} 136000 \\ 68000 \\ 204000 \end{bmatrix} = \begin{bmatrix} 2000 \\ 1000 \\ 3000 \end{bmatrix}$$

$$\therefore x = 2000, y = 1000, z = 3000.$$

Hence the prices per unit of the commodities

A, B and C are ₹ 2000, ₹ 1000 and ₹ 3000

respectively.

EXERCISE 1.4

1. Solve the following systems of linear equations by Cramer's rule :

(i) $5x - 2y + 16 = 0, x + 3y - 7 = 0$

(ii) $\frac{3}{x} + 2y = 12, \frac{2}{x} + 3y = 13$ [Qy. - 2023 & 2024]

(iii) $3x + 3y - z = 11, 2x - y + 2z = 9,$
 $4x + 3y + 2z = 25$ [Hy - 2019; July - 2022]

(iv) $\frac{3}{x} - \frac{4}{y} - \frac{2}{z} - 1 = 0, \frac{1}{x} + \frac{2}{y} + \frac{1}{z} - 2 = 0,$
 $\frac{2}{x} - \frac{5}{y} - \frac{4}{z} + 1 = 0$ [FRT -2022; Mar. - 2024]

Sol. (i) Given $\Delta = \begin{vmatrix} 5 & -2 \\ 1 & 3 \end{vmatrix} = 15 + 2 = 17$

$$\Delta_1 = \begin{vmatrix} -16 & -2 \\ 7 & 3 \end{vmatrix} = -48 + 14 = -34$$

$$\Delta_2 = \begin{vmatrix} 5 & -16 \\ 1 & 7 \end{vmatrix} = 35 + 16 = 51$$

$$\therefore x = \frac{\Delta_1}{\Delta} = \frac{-34}{17} = -2$$

$$y = \frac{\Delta_2}{\Delta} = \frac{51}{17} = 3$$

$$x = -2, y = 3.$$

(ii) Let $\frac{1}{x} = z$
 $\therefore 3z + 2y = 12, 2z + 3y = 13$

$$\therefore \Delta = \begin{vmatrix} 3 & 2 \\ 2 & 3 \end{vmatrix} = 9 - 4 = 5$$

$$\Delta_1 = \begin{vmatrix} 12 & 2 \\ 13 & 3 \end{vmatrix} = 36 - 26 = 10$$

$$\Delta_2 = \begin{vmatrix} 3 & 12 \\ 2 & 13 \end{vmatrix} = 39 - 24 = 15$$

$$\therefore z = \frac{\Delta_1}{\Delta} = \frac{10}{5} = 2 \Rightarrow \frac{1}{x} = 2 \Rightarrow x = \frac{1}{2}$$

$$y = \frac{\Delta_2}{\Delta} = \frac{15}{5} = 3$$

$$\therefore x = \frac{1}{2}, y = 3$$

$$\begin{aligned}
 \text{(iii)} \quad \Delta &= \begin{vmatrix} 3 & 3 & -1 \\ 2 & -1 & 2 \\ 4 & 3 & 2 \end{vmatrix} \\
 &= 3 \begin{vmatrix} -1 & 2 \\ 3 & 2 \end{vmatrix} - 3 \begin{vmatrix} 2 & 2 \\ 4 & 2 \end{vmatrix} - 1 \begin{vmatrix} 2 & -1 \\ 4 & 3 \end{vmatrix} \\
 &= 3(-2-6) - 3(4-8) - 1(6+4) \\
 &= 3(-8) - 3(-4) - 1(10) \\
 &= -24 + 12 - 10 = -22 \\
 \Delta_x &= \begin{vmatrix} 11 & 3 & -1 \\ 9 & -1 & 2 \\ 25 & 3 & 2 \end{vmatrix} \\
 &= 11 \begin{vmatrix} -1 & 2 \\ 3 & 2 \end{vmatrix} - 3 \begin{vmatrix} 9 & 2 \\ 25 & 2 \end{vmatrix} - 1 \begin{vmatrix} 9 & -1 \\ 25 & 3 \end{vmatrix} \\
 &= 11(-2-6) - 3(18-50) - 1(27+25) \\
 &= 11(-8) - 3(-32) - 1(52) \\
 &= -88 + 96 - 52 = -44 \\
 \Delta_y &= \begin{vmatrix} 3 & 11 & -1 \\ 2 & 9 & 2 \\ 4 & 25 & 2 \end{vmatrix} \\
 &= 3 \begin{vmatrix} 9 & 2 \\ 25 & 2 \end{vmatrix} - 11 \begin{vmatrix} 2 & 2 \\ 4 & 2 \end{vmatrix} - 1 \begin{vmatrix} 2 & 9 \\ 4 & 25 \end{vmatrix} \\
 &= 3(18-50) - 11(4-8) - 1(50-36) \\
 &= 3(-32) - 11(-4) - 1(14) \\
 &= -96 + 44 - 14 = -66 \\
 \Delta_z &= \begin{vmatrix} 3 & 3 & 11 \\ 2 & -1 & 9 \\ 4 & 3 & 25 \end{vmatrix} \\
 &= 3 \begin{vmatrix} -1 & 9 \\ 3 & 25 \end{vmatrix} - 3 \begin{vmatrix} 2 & 9 \\ 4 & 25 \end{vmatrix} + 11 \begin{vmatrix} 2 & -1 \\ 4 & 3 \end{vmatrix} \\
 &= 3(-25-27) - 3(50-36) + 11(6+4) \\
 &= 3(-52) - 3(14) + 11(10) \\
 &= -156 - 42 + 110 = -88 \\
 \therefore x &= \frac{\Delta_1}{\Delta} = \frac{-44}{-22} = 2 \\
 y &= \frac{\Delta_2}{\Delta} = \frac{-66}{-22} = 3 \\
 z &= \frac{\Delta_3}{\Delta} = \frac{-88}{-22} = 4
 \end{aligned}$$

$$\therefore x = 2, y = 3, z = 4.$$

$$\text{(iv)} \quad \text{Put } \frac{1}{x} = X, \frac{1}{y} = Y, \frac{1}{z} = Z$$

$$\text{We get } 3X - 4Y - 2Z = 1, X + 2Y + Z = 2, \\ 2X - 5Y - 4Z = -1$$

$$\begin{aligned}
 \therefore \Delta &= \begin{vmatrix} 3 & -4 & -2 \\ 1 & 2 & 1 \\ 2 & -5 & -4 \end{vmatrix} = 3 \begin{vmatrix} 2 & 1 \\ -5 & -4 \end{vmatrix} + 4 \begin{vmatrix} 1 & 1 \\ 2 & -4 \end{vmatrix} - 2 \begin{vmatrix} 1 & 2 \\ 2 & -5 \end{vmatrix} \\
 &= 3(-8+5) + 4(-4-2) - 2(-5-4) \\
 &= 3(-3) + 4(-6) - 2(-9) \\
 &= -9 - 24 + 18 = -15
 \end{aligned}$$

$$\begin{aligned}
 \Delta_x &= \begin{vmatrix} 1 & -4 & -2 \\ 2 & 2 & 1 \\ -1 & -5 & -4 \end{vmatrix} \\
 &= 1 \begin{vmatrix} 2 & 1 \\ -5 & -4 \end{vmatrix} + 4 \begin{vmatrix} 2 & 1 \\ -1 & -4 \end{vmatrix} - 2 \begin{vmatrix} 2 & 2 \\ -1 & -5 \end{vmatrix} \\
 &= 1(-8+5) + 4(-8+1) - 2(-10+2) \\
 &= 1(-3) + 4(-7) - 2(-8) \\
 &= -3 - 28 + 16 = -15
 \end{aligned}$$

$$\begin{aligned}
 \Delta_y &= \begin{vmatrix} 3 & 1 & -2 \\ 1 & 2 & 1 \\ 2 & -1 & -4 \end{vmatrix} \\
 &= 3 \begin{vmatrix} 2 & 1 \\ -1 & -4 \end{vmatrix} - 1 \begin{vmatrix} 1 & 1 \\ 2 & -4 \end{vmatrix} - 2 \begin{vmatrix} 1 & 2 \\ 2 & -1 \end{vmatrix} \\
 &= 3(-8+1) - 1(-4-2) - 2(-1-4) \\
 &= 3(-7) - 1(-6) - 2(-5) = -21 + 6 + 10 = -5 \\
 \Delta_z &= \begin{vmatrix} 3 & -4 & 1 \\ 1 & 2 & 2 \\ 2 & -5 & -1 \end{vmatrix} = 3 \begin{vmatrix} 2 & 2 \\ -5 & -1 \end{vmatrix} + 4 \begin{vmatrix} 1 & 2 \\ 2 & -1 \end{vmatrix} + 1 \begin{vmatrix} 1 & 2 \\ 2 & -5 \end{vmatrix} \\
 &= 3(-2+10) + 4(-1-4) + 1(-5-4) \\
 &= 3(8) + 4(-5) + 1(-9) \\
 &= 24 - 20 - 9 = -5
 \end{aligned}$$

$$\therefore X = \frac{\Delta_x}{\Delta} = \frac{-15}{-15} = 1 \Rightarrow \frac{1}{x} = 1$$

$$\Rightarrow x = 1$$

$$Y = \frac{\Delta_y}{\Delta} = \frac{-5}{-15} = \frac{1}{3}$$

$$\Rightarrow \frac{1}{y} = \frac{1}{3} \Rightarrow y = 3$$

$$Z = \frac{\Delta_z}{\Delta} = \frac{-5}{-15} = \frac{1}{3} \Rightarrow \frac{1}{z} = \frac{1}{3} \Rightarrow z = 3$$

$$\therefore x = 1, y = 3, z = 3.$$

2. In a competitive examination, one mark is awarded for every correct answer while $\frac{1}{4}$ mark is deducted for every wrong answer. A student answered 100 questions and got 80 marks. How many questions did he answer correctly? (Use Cramer's rule to solve the problem). [Qy - 2019]

Sol. Let x represent the number of question with correct answer and y represent the number of questions with wrong answers.

By the given data, $x + y = 100$ and ... (1)

$$1 \cdot x - \frac{1}{4} y = 80$$

Multiplying by 4 we get,

$$4x - y = 320 \quad \dots (2)$$

From (1) and (2)

$$\Delta = \begin{vmatrix} 1 & 1 \\ 4 & -1 \end{vmatrix} = -1 - 4 = -5$$

$$\Delta_1 = \begin{vmatrix} 100 & 1 \\ 320 & -1 \end{vmatrix} = -100 - 320 = -420$$

$$\Delta_2 = \begin{vmatrix} 1 & 100 \\ 4 & 320 \end{vmatrix} = 320 - 400 = -80$$

$$\therefore x = \frac{\Delta_1}{\Delta} = \frac{-420}{-5} = +84$$

$$\text{and } y = \frac{\Delta_2}{\Delta} = \frac{-80}{-5} = 16$$

Hence, the number of questions with correct answer is 84.

3. A chemist has one solution which is 50% acid and another solution which is 25% acid. How much each should be mixed to make 10 litres of a 40% acid solution ? (Use Cramer's rule to solve the problem). [Hy. - 2023]

Sol. Let the amount of 50% acid be x litres and the amount of 25% acid be y litres

By the given data, $x + y = 10$... (1)

$$\text{and } x \left(\frac{50}{100} \right) + y \left(\frac{25}{100} \right) = 10 \left(\frac{40}{100} \right)$$

$$\Rightarrow 50x + 25y = 400 \Rightarrow 2x + y = 16 \quad \dots (2)$$

$$\Delta = \begin{vmatrix} 1 & 1 \\ 2 & 1 \end{vmatrix} = -1$$

$$\Delta_x = \begin{vmatrix} 10 & 1 \\ 16 & 1 \end{vmatrix} = 10 - 16 = -6$$

$$\Rightarrow \Delta_y = \begin{vmatrix} 1 & 10 \\ 2 & 16 \end{vmatrix} = 16 - 20 = -4$$

$$x = \frac{\Delta_x}{\Delta} = \frac{-6}{-1} = 6$$

$$y = \frac{\Delta_y}{\Delta} = \frac{-4}{-1} = 4$$

i.e., 6 litres of 50% acid and 4 litres of 25% acid solution to be mixed to get 10 litres of 40% of acid solution.

4. A fish tank can be filled in 10 minutes using both pumps A and B simultaneously. However, pump B can pump water in or out at the same rate. If pump B is inadvertently run in reverse, then the tank will be filled in 30 minutes. How long would it take each pump to fill the tank by itself ? (Use Cramer's rule to solve the problem).

Sol. Let the pump A can fill the tank in x minutes, and the pump B can fill the tank in y minutes

In 1 minute A can fill $\frac{1}{x}$ units and in 1 minute B can fill $\frac{1}{y}$ units

$$\therefore \frac{1}{x} + \frac{1}{y} = \frac{1}{10} \text{ and } \frac{1}{x} - \frac{1}{y} = \frac{1}{30}$$

$$\text{Put } \frac{1}{x} = a \text{ and } \frac{1}{y} = b$$

$$\Rightarrow a + b = \frac{1}{10} \quad \dots (1)$$

$$\text{and } a - b = \frac{1}{30} \quad \dots (2)$$

$$\Delta = \begin{vmatrix} 1 & 1 \\ 1 & -1 \end{vmatrix} = -1 - 1 = -2$$

$$\Delta_1 = \begin{vmatrix} \frac{1}{10} & 1 \\ \frac{1}{30} & -1 \end{vmatrix} = \frac{-1}{10} - \frac{1}{30} = \frac{-3-1}{30} = \frac{-4}{30} = \frac{-2}{15}$$

$$\Delta_2 = \begin{vmatrix} 1 & \frac{1}{10} \\ 1 & \frac{1}{30} \end{vmatrix} = \frac{1}{30} - \frac{1}{10} = \frac{1-3}{30} = \frac{-2}{30} = \frac{-1}{15}$$

$$\therefore a = \frac{\Delta_1}{\Delta} = \frac{-2}{15} \div \frac{-2}{15} = \frac{1}{15} \Rightarrow \frac{1}{x} = \frac{1}{15} \Rightarrow x = 15$$

$$b = \frac{\Delta_2}{\Delta} = \frac{-1}{15} \div \frac{-2}{15} = \frac{1}{30} \Rightarrow \frac{1}{y} = \frac{1}{30} \Rightarrow y = 30$$

Hence the pump A can fill the tank in 15 minutes and the pump B can fill the tank in 30 minutes.

5. A family of 3 people went out for dinner in a restaurant. The cost of two dosai, three idlies and two vadais is ₹ 150. The cost of the two dosai, two idlies and four vadais is ₹ 200. The cost of five dosai, four idlies and two vadais is ₹ 250. The family has ₹ 350 in hand and they ate 3 dosai and six idlies and six vadais. Will

they be able to manage to pay the bill within the amount they had ?

Sol. Let the cost of one dosa be ₹ x

The cost of one idli be ₹ y

and the cost of one vadai be ₹ z

By the given data,

$$2x + 3y + 2z = 150$$

$$2x + 2y + 4z = 200$$

$$5x + 4y + 2z = 250$$

$$\therefore \Delta = \begin{vmatrix} 2 & 3 & 2 \\ 2 & 2 & 4 \\ 5 & 4 & 2 \end{vmatrix}$$

$$= 2 \begin{vmatrix} 2 & 4 \\ 4 & 2 \end{vmatrix} - 3 \begin{vmatrix} 2 & 4 \\ 5 & 2 \end{vmatrix} + 2 \begin{vmatrix} 2 & 2 \\ 5 & 4 \end{vmatrix}$$

$$= 2(4 - 16) - 3(4 - 20) + 2(8 - 10)$$

$$= 2(-12) - 3(-16) + 2(-2)$$

$$= -24 + 48 - 4 = 20$$

$$\Delta_1 = \begin{vmatrix} 150 & 3 & 2 \\ 200 & 2 & 4 \\ 250 & 4 & 2 \end{vmatrix}$$

Taking 50 common from C_1 and 2 common from C_3 we get,

$$= 100 \begin{vmatrix} 3 & 3 & 1 \\ 4 & 2 & 2 \\ 5 & 4 & 1 \end{vmatrix} = 100 \left[3 \begin{vmatrix} 2 & 2 \\ 4 & 1 \end{vmatrix} - 3 \begin{vmatrix} 4 & 2 \\ 5 & 1 \end{vmatrix} + 1 \begin{vmatrix} 4 & 2 \\ 5 & 4 \end{vmatrix} \right]$$

$$= 100[3(2 - 8) - 3(4 - 10) + 1(16 - 10)]$$

$$= 100[3(-6) - 3(-6) + 6]$$

$$= 100[-18 + 18 + 6] = 600.$$

$$\Delta_2 = \begin{vmatrix} 2 & 150 & 2 \\ 2 & 200 & 4 \\ 5 & 250 & 2 \end{vmatrix} = 100 \begin{vmatrix} 2 & 3 & 1 \\ 2 & 4 & 2 \\ 5 & 5 & 1 \end{vmatrix}$$

$$= 100 \left[2 \begin{vmatrix} 4 & 2 \\ 5 & 1 \end{vmatrix} - 3 \begin{vmatrix} 2 & 2 \\ 5 & 1 \end{vmatrix} + 1 \begin{vmatrix} 2 & 4 \\ 5 & 5 \end{vmatrix} \right]$$

$$= 100[2(4 - 10) - 3(2 - 10) + 1(10 - 20)]$$

$$= 100[2(-6) - 3(-8) + 1(-10)]$$

$$= 100[-12 + 24 - 10] = 100[2] = 200.$$

$$\Delta_3 = \begin{vmatrix} 2 & 3 & 150 \\ 2 & 2 & 200 \\ 5 & 4 & 250 \end{vmatrix} = 50 \begin{vmatrix} 2 & 3 & 3 \\ 2 & 2 & 4 \\ 5 & 4 & 5 \end{vmatrix}$$

$$= 50 \left[2 \begin{vmatrix} 2 & 4 \\ 4 & 5 \end{vmatrix} - 3 \begin{vmatrix} 2 & 4 \\ 5 & 5 \end{vmatrix} + 3 \begin{vmatrix} 2 & 2 \\ 5 & 4 \end{vmatrix} \right]$$

$$= 50[2(10 - 16) - 3(10 - 20) + 3(8 - 10)]$$

$$= 50[2(-6) - 3(-10) + 3(-2)]$$

$$= 50[-12 + 30 - 6] = 50[12] = 600.$$

$$\therefore x = \frac{\Delta_1}{\Delta} = \frac{600}{20} = 30$$

$$y = \frac{\Delta_2}{\Delta} = \frac{200}{20} = 10$$

$$z = \frac{\Delta_3}{\Delta} = \frac{600}{20} = 30.$$

Hence, the price of one dosa be ₹ 30, one idli be ₹ 10 and the price of 1 vadai be ₹ 30.

Also the cost of 3 dosa, six idlies and six vadai is

$$= 3x + 6y + 6z = 3(30) + 6(10) + 6(30)$$

$$= 90 + 60 + 180 = ₹ 330$$

Since the family had ₹ 350 in hand, they will be able to manage to pay the bill.

EXERCISE 1.5

1. Solve the following systems of linear equations by Gaussian elimination method :

(i) $2x - 2y + 3z = 2, x + 2y - z = 3, 3x - y + 2z = 1.$

(ii) $2x + 4y + 6z = 22, 3x + 8y + 5z = 27, -x + y + 2z = 2$

Sol. (i) Transforming the augmented matrix to echelon form, we get

$$\left[\begin{array}{ccc|c} 2 & -2 & 3 & 2 \\ 1 & 2 & -1 & 3 \\ 3 & -1 & 2 & 1 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2}$$

$$\left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 2 & -2 & 3 & 2 \\ 3 & -1 & 2 & 1 \end{array} \right] \xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 3R_1 \end{array}} \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -6 & 5 & -4 \\ 0 & -7 & 5 & -8 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -6 & 5 & -4 \\ 0 & -1 & 0 & -4 \end{array} \right] \xrightarrow{R_3 \rightarrow 6R_3 - 2R_2}$$

$$\left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -6 & 5 & -4 \\ 0 & 0 & -5 & -20 \end{array} \right]$$

Writing the equivalent equations from the row-echelon matrix, we get,

$$x + 2y - z = 3 \quad \dots (1)$$

$$\begin{aligned} -6y + 5z &= -4 \\ -5z &= -20 \Rightarrow z = \frac{-20}{-5} = 4. \end{aligned} \quad \dots (2)$$

Substituting $z = 4$ in (2) we get,

$$\begin{aligned} -6y + 5(4) &= -4 \\ \Rightarrow -6y + 20 &= -4 \Rightarrow -4 - 20 = -24 \\ \Rightarrow y &= \frac{-24}{-6} = 4. \end{aligned}$$

Substituting $y = z = 4$ in (1) we get

$$\begin{aligned} x + 2(4) - 4 &= 3 \Rightarrow x + 8 - 4 = 3 \\ \Rightarrow x + 4 &= 3 \\ \Rightarrow x &= 3 - 4 = -1. \\ \therefore x &= -1, y = 4, z = 4. \end{aligned}$$

(ii) $2x + 4y + 6z = 22, 3x + 8y + 5z = 27,$
 $-x + y + 2z = 2$

Reducing the augmented matrix to an equivalent row echelon form by using elementary row operations, we get

$$\left[\begin{array}{ccc|c} 2 & 4 & 6 & 22 \\ 3 & 8 & 5 & 27 \\ -1 & 1 & 2 & 2 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_3} \left[\begin{array}{ccc|c} -1 & 1 & 2 & 2 \\ 3 & 8 & 5 & 27 \\ 2 & 4 & 6 & 22 \end{array} \right]$$

$$\begin{aligned} R_2 &\rightarrow R_2 + 3R_1 \\ R_3 &\rightarrow R_3 + 2R_1 \end{aligned} \rightarrow \left[\begin{array}{ccc|c} -1 & 1 & 2 & 2 \\ 0 & 11 & 11 & 33 \\ 0 & 6 & 10 & 26 \end{array} \right]$$

$$\begin{aligned} R_2 &\rightarrow R_2 \div 11 \\ R_3 &\rightarrow R_3 \div 2 \end{aligned} \rightarrow \left[\begin{array}{ccc|c} -1 & 1 & 2 & 2 \\ 0 & 1 & 1 & 3 \\ 0 & 3 & 5 & 13 \end{array} \right]$$

$$R_3 \rightarrow R_3 - 3R_2 \rightarrow \left[\begin{array}{ccc|c} -1 & 1 & 2 & 2 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 2 & 4 \end{array} \right]$$

$$R_3 \rightarrow R_3 \div 2 \rightarrow \left[\begin{array}{ccc|c} -1 & 1 & 2 & 2 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 1 & 2 \end{array} \right]$$

Writing the equivalent equations from the row echelon matrix we get,

$$\begin{aligned} -x + y + 2z &= 2 & \dots (1) \\ y + z &= 3 & \dots (2) \\ z &= 2 & \dots (3) \end{aligned}$$

Substituting (3) in (2) we get, $y + 2 = 3$
 $\Rightarrow y = 3 - 2 = 1$

Substituting $y = 1$ and $z = 2$ in (1) we get,

$$\begin{aligned} -x + 1 + 2(2) &= 2 & \Rightarrow -x + 1 + 4 = 2 \\ \Rightarrow -x + 5 &= 2 & \Rightarrow -x = 2 - 5 \\ \Rightarrow -x &= -3 & \Rightarrow x = 3 \end{aligned}$$

$$\therefore x = 3, y = 1, z = 2.$$

2. If $ax^2 + bx + c$ is divided by $x + 3$, $x - 5$, and $x - 1$, the remainders are 21, 61 and 9 respectively. Find a , b and c . (Use Gaussian elimination method.) [PTA -3]

Sol.

$$\text{Let } P(x) = ax^2 + bx + c$$

$$\text{Given } P(-3) = 21$$

$$[\because P(x) \div x + 3, \text{ the remainder is } 21]$$

$$\begin{aligned} \Rightarrow a(-3)^2 + b(-3) + c &= 21 \\ \Rightarrow 9a - 3b + c &= 21 \end{aligned} \quad \dots (1)$$

$$\text{Also, } P(5) = 61$$

$$\begin{aligned} \Rightarrow a(5)^2 + b(5) + c &= 61 \\ &[\text{using remainder theorem}] \\ \Rightarrow 25a + 5b + c &= 61 \end{aligned} \quad \dots (2)$$

$$\text{and } P(1) = 9$$

$$\begin{aligned} \Rightarrow a(1)^2 + b(1) + c &= 9 \\ \Rightarrow a + b + c &= 9 \end{aligned} \quad \dots (3)$$

Reducing the augment matrix to an equivalent row-echelon form using elementary row operations, we get

$$\left[\begin{array}{ccc|c} 9 & -3 & 1 & 21 \\ 25 & 5 & 1 & 61 \\ 1 & 1 & 1 & 9 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_3} \left[\begin{array}{ccc|c} 1 & 1 & 1 & 9 \\ 25 & 5 & 1 & 61 \\ 9 & -3 & 1 & 21 \end{array} \right]$$

$$\begin{aligned} R_3 &\rightarrow R_3 - 9R_1 \\ R_2 &\rightarrow R_2 - 25R_1 \end{aligned} \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 1 & 9 \\ 0 & -20 & -24 & -164 \\ 0 & -12 & -8 & -60 \end{array} \right]$$

$$\begin{aligned} R_2 &\rightarrow R_2 \div 4 \\ R_3 &\rightarrow R_3 \div 4 \end{aligned} \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 1 & 9 \\ 0 & -5 & -6 & -41 \\ 0 & -3 & -2 & -15 \end{array} \right]$$

$$R_3 \rightarrow R_3 - \frac{3}{5}R_2 \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 1 & 9 \\ 0 & -5 & -6 & -41 \\ 0 & 0 & \frac{8}{5} & \frac{48}{5} \end{array} \right]$$

$$R_3 \rightarrow 5R_3 \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 1 & 9 \\ 0 & -5 & -6 & -41 \\ 0 & 0 & 8 & 48 \end{array} \right]$$

Writing the equivalent equations from the row-echelon matrix we get,

$$a + b + c = 9 \quad \dots (1)$$

$$-5b - 6c = -41 \quad \dots (2)$$

$$8c = 48$$

$$\Rightarrow c = \frac{48}{8} = 6$$

Substituting $c = 6$ in (2) we get,

$$\Rightarrow -5b - 6(6) = -41 \Rightarrow -5b = 36 - 41$$

$$\Rightarrow -5b = -41 + 36 = -5$$

$$\Rightarrow b = \frac{-5}{-5} = 1$$

Substituting $b = 1, c = 6$ in (1) we get,

$$a + 1 + 6 = 9 \Rightarrow a + 7 = 9$$

$$\Rightarrow a = 9 - 7$$

$$\Rightarrow a = 2$$

$$\therefore a = 2, b = 1 \text{ and } c = 6$$

- 3. An amount of ₹ 65,000 is invested in three bonds at the rates of 6%, 8% and 9% per annum respectively. The total annual income is ₹ 4,800. The income from the third bond is ₹ 600 more than that from the second bond. Determine the price of each bond. (Use Gaussian elimination method.)**

Sol. Let the price of bond invested in 6%, 8% and 9% rates be let ₹ x , ₹ y and ₹ z respectively

$$\therefore \text{By the given data, } x + y + z = 65,000 \quad \dots(1)$$

$$\frac{6 \times x \times 1}{100} + \frac{8 \times y \times 1}{100} + \frac{9 \times z \times 1}{100} = 4,800$$

$$[\because \text{Interest} = \frac{\text{PNR}}{100}]$$

$$\Rightarrow \frac{6x}{100} + \frac{8y}{100} + \frac{9z}{100} = 4,800$$

$$\Rightarrow 6x + 8y + 9z = 4,80,000 \quad \dots(2)$$

$$\text{Also, } \frac{9z}{100} = 600 + \frac{8y}{100}$$

$$\Rightarrow \frac{-8y}{100} + \frac{9z}{100} = 600$$

$$\Rightarrow -8y + 9z = 60,000 \quad \dots(3)$$

Reducing the augmented matrix to an equivalent row-echelon form by using elementary row operation, we get

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 65,000 \\ 6 & 8 & 9 & 4,80,000 \\ 0 & -8 & 9 & 60,000 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 - 6R_1} \left[\begin{array}{ccc|c} 1 & 1 & 1 & 65,000 \\ 0 & 2 & 3 & 90,000 \\ 0 & -8 & 9 & 60,000 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 + 4R_2} \left[\begin{array}{ccc|c} 1 & 1 & 1 & 65,000 \\ 0 & 2 & 3 & 90,000 \\ 0 & 0 & 21 & 4,20,000 \end{array} \right]$$

Writing the equivalent from the row echelon matrix we get,

$$x + y + z = 65,000 \quad \dots(1)$$

$$2y + 3z = 90,000 \quad \dots(2)$$

$$21z = 4,20,000$$

$$\Rightarrow z = \frac{4,20,000}{21} = 20,000$$

Substituting $z = 20,000$ in (2),

$$2y + 3(20,000) = 90,000$$

$$\Rightarrow 2y + 60,000 = 90,000$$

$$\Rightarrow 2y = 90,000 - 60,000$$

$$= 30,000$$

$$\Rightarrow y = \frac{30,000}{2} = 15,000$$

Substituting $y = 15,000$ and $z = 20,000$ in (1) we get,

$$x + 15,000 + 20,000 = 65,000$$

$$\Rightarrow x + 35,000 = 65,000$$

$$\Rightarrow x = 65,000 - 35,000$$

$$\Rightarrow x = 30,000$$

Thus the price of 6% bond is ₹ 30,000 the price of 8% bond is ₹ 15,000 and the price of 9% bond is ₹ 20,000.

- 4. A boy is walking along the path $y = ax^2 + bx + c$ through the points $(-6, 8)$, $(-2, -12)$ and $(3, 8)$. He wants to meet his friend at $P(7, 60)$. Will he meet his friend? (Use Gaussian elimination method.)** [Mar. - 2023]

Sol. Given $y = ax^2 + bx + c \quad \dots(1)$

$(-6, 8)$ lies on (1)

$$\Rightarrow 8 = a(-6)^2 + b(-6) + c$$

$$\Rightarrow 8 = 36a - 6b + c \quad \dots(2)$$

$(-2, -12)$ lies on (1)

$$\Rightarrow -12 = a(-2)^2 + b(-2) + c$$

$$\Rightarrow -12 = 4a - 2b + c \quad \dots(3)$$

Also $(3, 8)$ lies on (1)

$$\Rightarrow 8 = a(3)^2 + b(3) + c$$

$$\Rightarrow 8 = 9a + 3b + c \quad \dots(4)$$

Reducing the augment matrix to an equivalent row-echelon form by using elementary row operations, we get,

$$\left[\begin{array}{ccc|c} 36 & -6 & 1 & 8 \\ 4 & -2 & 1 & -12 \\ 9 & 3 & 1 & 8 \end{array} \right] \xrightarrow{\begin{array}{l} R_2 \rightarrow 9R_2 - R_1 \\ R_3 \rightarrow 4R_3 - R_1 \end{array}} \left[\begin{array}{ccc|c} 36 & -6 & 1 & 8 \\ 0 & -12 & 8 & -116 \\ 0 & 18 & 3 & 24 \end{array} \right]$$

$$\begin{array}{l} R_2 \rightarrow R_2 \div 4 \\ R_3 \rightarrow R_3 \div 3 \end{array} \rightarrow \left[\begin{array}{ccc|c} 36 & -6 & 1 & 8 \\ 0 & -3 & 2 & -29 \\ 0 & 6 & 1 & 8 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 + 2R_2} \left[\begin{array}{ccc|c} 36 & -6 & 1 & 8 \\ 0 & -3 & 2 & -29 \\ 0 & 0 & 5 & -50 \end{array} \right]$$

Writing the equivalent equation from the row echelon matrix, we get $36a - 6b + c = 8$... (1)

$$-3b + 2c = -29 \quad \dots (2)$$

$$5c = -50$$

$$\Rightarrow c = \frac{-50}{5} = -10$$

Substituting $c = -10$ in (2) we get,

$$-3b + 2(-10) = -29$$

$$\Rightarrow -3b - 20 = -29$$

$$\Rightarrow -3b = -29 + 20 \Rightarrow -3b = -9$$

$$\Rightarrow b = \frac{-9}{-3} = 3$$

Substituting $b = 3$ and $c = -10$ in (1) we get,

$$36a - 6(3) - 10 = 8$$

$$\Rightarrow 36a - 18 - 10 = 8 \Rightarrow 36a - 28 = 8$$

$$\Rightarrow 36a = 8 + 28 = 36$$

$$\Rightarrow a = \frac{36}{36} = 1$$

$$\therefore a = 1, b = 3, c = -10$$

Hence the path of the boy is

$$y = 1(x^2) + 3(x) - 10$$

$$\Rightarrow y = x^2 + 3x - 10$$

Since his friend is at P(7, 60),

$$60 = (7)^2 + 3(7) - 10$$

$$\Rightarrow 60 = 49 + 21 - 10$$

$$\Rightarrow 60 = 70 - 10 = 60$$

$$\Rightarrow 60 = 60$$

Since (7, 60) satisfies his path, he can meet his friend who is at P(7, 60)

EXERCISE 1.6

1. Test for consistency and if possible, solve the following systems of equations by rank method.

(i) $x - y + 2z = 2, 2x + y + 4z = 7,$
 $4x - y + z = 4$

(ii) $3x + y + z = 2, x - 3y + 2z = 1,$
 $7x - y + 4z = 5.$

(iii) $2x + 2y + z = 5, x - y + z = 1,$

$3x + y + 2z = 4$ [Sep. - 2020; Hy. - 2023]

(iv) $2x - y + z = 2, 6x - 3y + 3z = 6,$

$4x - 2y + 2z = 4.$ [PTA - 5; Hy - 2019]

Sol. (i) $x - y + 2z = 2, 2x + y + 4z = 7, 4x - y + z = 4$

The matrix form of the system is $AX = B$

where $A = \begin{bmatrix} 1 & -1 & 2 \\ 2 & 1 & 4 \\ 4 & -1 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ and $B = \begin{bmatrix} 2 \\ 7 \\ 4 \end{bmatrix}$

Applying elementary row operations on the augmented matrix $[A|B]$ we get,

$$[A|B] = \left[\begin{array}{ccc|c} 1 & -1 & 2 & 2 \\ 2 & 1 & 4 & 7 \\ 4 & -1 & 1 & 4 \end{array} \right] \xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 4R_1 \end{array}} \left[\begin{array}{ccc|c} 1 & -1 & 2 & 2 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & -7 & -4 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{ccc|c} 1 & -1 & 2 & 2 \\ 0 & 3 & 0 & 3 \\ 0 & 0 & -7 & -7 \end{array} \right]$$

Here $\rho(A) = 3$ and $\rho[A|B] = 3$

$\therefore \rho(A) = \rho[A|B] = 3 = \text{number of unknowns}$

Hence the system is consistent with unique solution.

Writing the equivalent equations from the row echelon matrix, we get

$$x - y + 2z = 2 \quad \dots (1)$$

$$3y = 3 \Rightarrow y = 1 \quad \dots (2)$$

$$-7z = -7 \Rightarrow z = \frac{-7}{-7} = 1 \quad \dots (3)$$

Solutions $y = 1$ and $z = 1$ in (1) we get,

$$x - 1 + 2(1) = 2 \Rightarrow x - 1 + 2 = 2$$

$$\Rightarrow x + 1 = 2 \Rightarrow x = 2 - 1 = 1$$

$$\Rightarrow x = 1$$

$$\therefore x = 1, y = 1, z = 1$$

(ii) $3x + y + z = 2, x - 3y + 2z = 1,$

$7x - y + 4z = 5.$

Then matrix form of the system is $AX = B$

where $A = \begin{bmatrix} 3 & 1 & 1 \\ 1 & -3 & 2 \\ 7 & -1 & 4 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 2 \\ 1 \\ 5 \end{bmatrix}$

Applying elementary row operations on the augmented matrix $[A|B]$ we get,

$$[A|B] = \left[\begin{array}{ccc|c} 3 & 1 & 1 & 2 \\ 1 & -3 & 2 & 1 \\ 7 & -1 & 4 & 5 \end{array} \right] \xrightarrow{R_2 \leftrightarrow R_1} \left[\begin{array}{ccc|c} 1 & -3 & 2 & 1 \\ 3 & 1 & 1 & 2 \\ 7 & -1 & 4 & 5 \end{array} \right]$$

$$\xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - 3R_1 \\ R_3 \rightarrow R_3 - 7R_1 \end{array}} \left[\begin{array}{ccc|c} 1 & -3 & 2 & 1 \\ 0 & 10 & -5 & -1 \\ 0 & 20 & -10 & -2 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - 2R_2} \left[\begin{array}{ccc|c} 1 & -3 & 2 & 1 \\ 0 & 10 & -5 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Here $\rho(A) = 2$, and $\rho[A|B] = 2$ [since there only two non-zero rows]. So, $\rho(A) = \rho[A|B] = 2 < 3$, the given system is consistent with one parameter family of solutions. So, put $z = t$, $x \in \mathbb{R}$. Writing the equivalent equations from the row echelon matrix we get,

$$x - 3y + 2z = 1 \quad \dots(1)$$

$$10y - 5z = -1 \quad \dots(2)$$

$$z = t \quad \dots(3)$$

(2) becomes $10y - 5t = -1$

$$\Rightarrow 10y = 5t - 1 \Rightarrow y = \frac{1}{10} [5t - 1]$$

$$\text{Also, from (1), } x - \frac{3}{10} [5t - 1] + 2t = 1$$

$$\Rightarrow x = \frac{3}{10} [5t - 1] - 2t + 1 = \frac{15t - 3 - 20t + 10}{10}$$

$$\Rightarrow x = \frac{1}{10} [-5t + 7]$$

Hence the solution set is

$$x = \frac{1}{10} (7 - 5t), y = \frac{1}{10} (5t - 1) \text{ and } z = t \text{ where}$$

$t \in \mathbb{R}$.

(iii) $2x + 2y + z = 5, x - y + z = 1, 3x + y + 2z = 4$

The matrix form of the given system is $AX = B$ where

$$A = \begin{bmatrix} 2 & 2 & 1 \\ 1 & -1 & 1 \\ 3 & 1 & 2 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 5 \\ 1 \\ 4 \end{bmatrix}$$

Applying elementary row operations on the augmented matrix $[A|B]$ we get,

$$[A|B] = \left[\begin{array}{ccc|c} 2 & 2 & 1 & 5 \\ 1 & -1 & 1 & 1 \\ 3 & 1 & 2 & 4 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccc|c} 1 & -1 & 1 & 1 \\ 2 & 2 & 1 & 5 \\ 3 & 1 & 2 & 4 \end{array} \right]$$

$$\xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 3R_1 \end{array}} \left[\begin{array}{ccc|c} 1 & -1 & 1 & 1 \\ 0 & 4 & -1 & 3 \\ 0 & 4 & -1 & 1 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{ccc|c} 1 & -1 & 1 & 1 \\ 0 & 4 & -1 & 3 \\ 0 & 0 & 0 & -2 \end{array} \right]$$

Here $\rho(A) = 2$ [∵ There are 2 non-Zero rows] and $\rho[A|B] = 3$ [∵ There are 3 non-zero rows]

Here, $\rho(A) \neq \rho[A|B]$

Hence, the given system is inconsistent and has no solution.

(iv) $2x - y + z = 2, 6x - 3y + 3z = 6, 4x - 2y + 2z = 4$.

The matrix form of the given system is

$AX = B$ where

$$A = \begin{bmatrix} 2 & -1 & 1 \\ 6 & -3 & 3 \\ 4 & -2 & 2 \end{bmatrix}$$

$$X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \quad B = \begin{bmatrix} 2 \\ 6 \\ 4 \end{bmatrix}$$

Applying elementary row operations on the augmented matrix $[A|B]$, we get,

$$[A|B] = \left[\begin{array}{ccc|c} 2 & -1 & 1 & 2 \\ 6 & -3 & 3 & 6 \\ 4 & -2 & 2 & 4 \end{array} \right] \xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - 3R_1 \\ R_3 \rightarrow R_3 - 2R_1 \end{array}} \left[\begin{array}{ccc|c} 2 & -1 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Here $\rho(A) = 1$ [∵ only one non zero row]

and $\rho[A|B] = 1$ [∵ only one-zero row]

∴ $\rho(A) = \rho[A|B] = 1 < 3$, the given system is consistent and has two parameter family of solutions.

So, $z = t$ and $y = s$ where $s, t \in \mathbb{R}$.

Writing the equivalent equations from the row-echelon matrix, we get

$$2x - y + z = 2 \quad \dots(1)$$

$$y = s \quad \dots(2)$$

$$z = t \quad \dots(3)$$

Substituting (2) and (3) in (1) we get,

$$2x - s + t = 2$$

$$\Rightarrow 2x = s - t + 2$$

$$\Rightarrow x = \frac{1}{2} [s - t + 2]$$

∴ Solution set is $x = \frac{1}{2} (s - t + 2), y = s, z = t$ where $s, t \in \mathbb{R}$

2. Find the value of k for which the equations
 $kx - 2y + z = 1$, $x - 2ky + z = -2$,
 $x - 2y + kz = 1$ have [Qy. - 2019 & 2023]

- (i) no solution
 (ii) unique solution
 (iii) infinitely many solution

Sol. $kx - 2y + z = 1$, $x - 2ky + z = -2$, $x - 2y + kz = 1$

The matrix form of the system is $AX = B$ where

$$A = \begin{bmatrix} k & -2 & 1 \\ 1 & -2k & 1 \\ 1 & -2 & k \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$$

Applying elementary row operation on the augmented matrix $[A|B]$ we get,

$$[A|B] = \left[\begin{array}{ccc|c} k & -2 & 1 & 1 \\ 1 & -2k & 1 & -2 \\ 1 & -2 & k & 1 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_3} \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 1 & -2k & 1 & -2 \\ k & -2 & 1 & 1 \end{array} \right]$$

$$\begin{array}{l} R_2 \rightarrow R_2 - R_1 \\ R_3 \rightarrow R_3 - kR_1 \end{array} \rightarrow \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 0 & -2k+2 & 1-k & -3 \\ 0 & -2+2k & 1-k^2 & 1-k \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 + R_2} \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 0 & -2k+2 & 1-k & -3 \\ 0 & 0 & 1-k^2+1-k & -k-2 \end{array} \right]$$

$$\begin{array}{l} \rightarrow \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 0 & -2k+2 & 1-k & -3 \\ 0 & 0 & -k^2-k+2 & -k-2 \end{array} \right] \\ \rightarrow \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 0 & -2k+2 & 1-k & -3 \\ 0 & 0 & (k+2)(1-k) & -k-2 \end{array} \right] \dots (1) \end{array}$$

Case (i) when $k = 1$

$$[A|B] \rightarrow \left[\begin{array}{ccc|c} 1 & -2 & 1 & 1 \\ 0 & 0 & 0 & -3 \\ 0 & 0 & 0 & -3 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 1 \\ 0 & 0 & 0 & -3 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Here $\rho(A) = 1$ and $\rho[A|B] = 2$

So, $\rho(A) \neq \rho[A|B] \Rightarrow$ The system has no solution.

Case (ii) When $k \neq 1$, $k \neq -2$

$$[A|B] \rightarrow \left[\begin{array}{ccc|c} 1 & -2 & k & 1 \\ 0 & -2k+2 & 1-k & -3 \\ 0 & 0 & \text{not zero} & \text{not zero} \end{array} \right]$$

$\Rightarrow \rho(A) = 3$ and $\rho[A|B] = 3$

so, $\rho(A) = \rho[A|B] = 3 =$ the number of unknowns
 Hence, the system has unique solution.

Case (iii) when $k = -2$

$$\rho[A|B] \rightarrow \left[\begin{array}{ccc|c} 1 & -2 & -2 & 1 \\ 1 & 6 & 3 & -3 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Here $\rho(A) = 2$ and $\rho[A|B] = 2$

$\therefore \rho(A) = \rho[A|B] = 2 < 3$, the number of unknowns
 so the system is consistent with infinitely many solutions.

3. Investigate the values of λ and μ the system of linear equations $2x + 3y + 5z = 9$, $7x + 3y - 5z = 8$, $2x + 3y + \lambda z = \mu$, have

- (i) no solution
 (ii) a unique solution
 (iii) an infinite number of solutions.

Sol. $2x + 3y + 5z = 9$, $7x + 3y - 5z = 8$, $2x + 3y + \lambda z = \mu$

The matrix form of the system is $AX = B$ where

$$A = \begin{bmatrix} 2 & 3 & 5 \\ 7 & 3 & -5 \\ 2 & 3 & \lambda \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 9 \\ 8 \\ \mu \end{bmatrix}$$

Applying elementary row operations on the augmented matrix $[A|B]$ we get,

$$[A|B] = \left[\begin{array}{ccc|c} 2 & 3 & 5 & 9 \\ 7 & 3 & -5 & 8 \\ 2 & 3 & \lambda & \mu \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccc|c} 7 & 3 & -5 & 8 \\ 2 & 3 & 5 & 9 \\ 2 & 3 & \lambda & \mu \end{array} \right]$$

$$\begin{array}{l} R_2 \rightarrow R_2 - \frac{2}{7}R_1 \\ R_3 \rightarrow R_3 - R_2 \end{array} \rightarrow \left[\begin{array}{ccc|c} 7 & 3 & -5 & -8 \\ 0 & \frac{15}{7} & \frac{45}{7} & \frac{45}{7} \\ 0 & 0 & \lambda - 5 & \mu - 9 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 \times 7} \left[\begin{array}{ccc|c} 7 & 3 & -5 & -8 \\ 0 & 15 & 45 & 47 \\ 0 & 0 & \lambda - 5 & \mu - 9 \end{array} \right]$$

Case (i) When $\lambda = 5$

$$[A|B] = \left[\begin{array}{ccc|c} 7 & 3 & -5 & -8 \\ 0 & 15 & 45 & 47 \\ 0 & 0 & 0 & \mu - 9 \end{array} \right]$$

Here $\rho(A) = 2$ and $\rho[A|B] = 3$

So, $\rho(A) \neq \rho[A|B]$

Hence the system is inconsistent and has no solution

Case (ii) When $\lambda \neq 5$, $\mu \neq 9$

$$[A|B] = \left[\begin{array}{ccc|c} 7 & 3 & -5 & -8 \\ 0 & 15 & 45 & 47 \\ 0 & 0 & \text{not zero} & \text{not zero} \end{array} \right]$$

Here $\rho(A) = 3$ and $\rho[A|B] = 3$

$\therefore \rho(A) = \rho[A|B] = 3 =$ number of unknowns

Hence, the system is consistent with unique solution

Case (iii) When $\lambda = 5$ and $\mu = 9$

$$[A|B] = \left[\begin{array}{ccc|c} 7 & 3 & -5 & -8 \\ 0 & 15 & 45 & 47 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Here $\rho(A) = 2$, $\rho[A|B] = 2$

∴ $\rho(A) = [A|B] = 2 < \text{number of unknowns}$

∴ The system is consistent and has infinite number of solutions.

EXERCISE 1.7

1. Solve the following system of homogeneous equations.

(i) $3x + 2y + 7z = 0, 4x - 3y - 2z = 0,$
 $5x + 9y + 23z = 0$

(ii) $2x + 3y - z = 0, x - y - 2z = 0, 3x + y + 3z = 0.$

Sol. (i) $3x + 2y + 7z = 0, 4x - 3y - 2z = 0,$
 $5x + 9y + 23z = 0$

Here the number of equations is equal to the number of unknowns.

Transforming into echelon form, the augmented matrix becomes

$$\left[\begin{array}{ccc|c} 3 & 2 & 7 & 0 \\ 4 & -3 & -2 & 0 \\ 5 & 9 & 23 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 3 & 2 & 7 & 0 \\ 0 & -17 & -34 & 0 \\ 0 & 17 & 34 & 0 \end{array} \right] \begin{array}{l} R_2 = 3R_2 - 4R_1 \\ R_3 = 3R_3 - 5R_1 \end{array}$$

$$\left[\begin{array}{ccc|c} 3 & 2 & 7 & 0 \\ 0 & -17 & -34 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \begin{array}{l} R_3 = R_3 + R_2 \end{array}$$

So, $\rho(AB) = \rho(A) = 2$ and $n = 3$.

Hence, the system has a one parameter family of solutions. Writing the equations using the echelon form, we get

$$3x + 2y + 7z = 0 \text{ and } -17y - 34z = 0$$

To solve the equations let $z = t$,

$$\text{then } -17y - 34t = 0$$

$$-17y = 34t$$

$$y = \frac{34t}{-17}$$

$$y = -2t$$

Substituting $z = t$ and $y = -2t$ in

$$3x + 2y + 7z = 0$$

$$\text{We get } 3x + 2(-2t) + 7t = 0$$

$$3x - 4t + 7t = 0$$

$$3x + 3t = 0$$

$$3x = -3t$$

$$x = -\frac{3t}{3}$$

$$x = -t,$$

So the solution is

$$x = -t, y = -2t \text{ and } z = t \text{ where } t \in \mathbb{R}$$

(ii) $2x + 3y - z = 0, x - y - 2z = 0, 3x + y + 3z = 0.$

Here the number of equations is equal to the number of unknowns. Transforming into echelon form, the augmented matrix becomes

$$\left[\begin{array}{ccc|c} 2 & 3 & -1 & 0 \\ 1 & -1 & -2 & 0 \\ 3 & 1 & 3 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & -1 & -2 & 0 \\ 2 & 3 & -1 & 0 \\ 3 & 1 & 3 & 0 \end{array} \right] \begin{array}{l} R_1 \leftrightarrow R_2 \\ R_2 = R_2 - 2R_1 \\ R_3 = R_3 - 3R_1 \end{array}$$

$$\rightarrow \left[\begin{array}{ccc|c} 1 & -1 & -2 & 0 \\ 0 & 5 & 3 & 0 \\ 0 & 0 & 33 & 0 \end{array} \right] \begin{array}{l} R_3 = 5R_2 - 4R_2 \end{array}$$

So, $\rho(AB) = \rho(A) = 3$ and $n = 3$

Hence, the system has a unique solution.

Since $x = 0, y = 0, z = 0$, is always a solution of the homogeneous system, the only solution is the trivial solution

$$x = 0, y = 0, z = 0$$

2. Determine the values of λ for which the following system of equations

$$x + y + 3z = 0, 4x + 3y + \lambda z = 0, 2x + y + 2z = 0 \text{ has}$$

(i) a unique solution

(ii) a non-trivial solution.

Sol. $x + y + 3z = 0, 4x + 3y + \lambda z = 0, 2x + y + 2z = 0$

Reducing the augmented matrix to row – echelon form we get,

$$[A|0] = \left[\begin{array}{ccc|c} 1 & 1 & 3 & 0 \\ 4 & 3 & \lambda & 0 \\ 2 & 1 & 2 & 0 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_3} \left[\begin{array}{ccc|c} 1 & 1 & 3 & 0 \\ 2 & 1 & 2 & 0 \\ 4 & 3 & \lambda & 0 \end{array} \right]$$

$$\xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 4R_1 \end{array}} \left[\begin{array}{ccc|c} 1 & 1 & 3 & 0 \\ 0 & -1 & -4 & 0 \\ 0 & -1 & \lambda - 12 & 0 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{ccc|c} 1 & 1 & 3 & 0 \\ 0 & -1 & -4 & 0 \\ 0 & 0 & \lambda - 8 & 0 \end{array} \right]$$

Case (i) when $\lambda \neq 8$

$$[A|0] = \left[\begin{array}{ccc|c} 1 & 1 & 3 & 0 \\ 0 & -1 & -4 & 0 \\ 0 & 0 & \text{not zero} & 0 \end{array} \right]$$

Here $\rho(A) = 3$, $\rho([A|0]) = 3$

$\therefore \rho(A) = \rho([A|0]) = 3 =$ the number of unknowns

$\therefore$ The given system is consistent and has unique solution.

Case (ii) when $\lambda = 8$

$$[A|0] = \left[\begin{array}{ccc|c} 1 & 1 & 3 & 0 \\ 0 & -1 & -4 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Here $\rho(A) = 2$, $\rho([A|0]) = 2$

$\therefore \rho(A) = \rho([A|0]) = 2 < 3$, the number of unknowns,

$\therefore$ The system is consistent and has non-trivial solutions.

3. By using Gaussian elimination method, balance the chemical reaction equation:



Sol. Given $\text{C}_2\text{H}_6 + \text{O}_2 \rightarrow \text{H}_2\text{O} + \text{CO}_2$

We have to find positive integers x_1, x_2, x_3 and x_4 such that



The number of carbon atoms on the LHS of (1) should be equal to the number of carbon atoms on the RHS of (1).

$$\therefore 2x_1 = 1x_4$$

$$\Rightarrow 2x_1 - x_4 = 0 \quad \dots(2)$$

Considering hydrogen atoms we get,

$$6x_1 = 2x_3 \Rightarrow 6x_1 - 2x_3 = 0$$

$$\Rightarrow 3x_1 - x_3 = 0 \quad \dots(3)$$

Also, considering oxygen atoms we get,

$$2x_2 = 1x_3 + 2x_4$$

$$\Rightarrow 2x_2 - x_3 - 2x_4 = 0 \quad \dots(4)$$

Equations (2), (3) and (4) form a homogeneous system of linear equations in 4 unknowns

$\therefore$ The augmented matrix $[A|0]$ is

$$\left[\begin{array}{cccc|c} 2 & 0 & 0 & -1 & 0 \\ 3 & 0 & -1 & 0 & 0 \\ 0 & 2 & -1 & -2 & 0 \end{array} \right]$$

By Gaussian elimination method, we get,

$$\xrightarrow{R_1 \rightarrow R_1 \div 2} \left[\begin{array}{cccc|c} 1 & 0 & 0 & -\frac{1}{2} & 0 \\ 3 & 0 & -1 & 0 & 0 \\ 0 & 2 & -1 & -2 & 0 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 - 3R_1} \left[\begin{array}{cccc|c} 1 & 0 & 0 & -\frac{1}{2} & 0 \\ 0 & 0 & -1 & \frac{3}{2} & 0 \\ 0 & 2 & -1 & -2 & 0 \end{array} \right]$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{cccc|c} 1 & 0 & 0 & -\frac{1}{2} & 0 \\ 0 & 0 & -1 & \frac{3}{2} & 0 \\ 0 & 2 & 0 & -\frac{7}{2} & 0 \end{array} \right]$$

$$\xrightarrow{R_2 \leftrightarrow R_3} \left[\begin{array}{cccc|c} 1 & 0 & 0 & -\frac{1}{2} & 0 \\ 0 & 2 & 0 & -\frac{7}{2} & 0 \\ 0 & 0 & -1 & \frac{3}{2} & 0 \end{array} \right]$$

$$\begin{array}{l} R_1 \rightarrow 2R_1 \\ R_2 \rightarrow 2R_2 \\ R_2 \rightarrow 2R_3 \end{array} \rightarrow \left[\begin{array}{cccc|c} 2 & 0 & 0 & -1 & 0 \\ 0 & 4 & 0 & -7 & 0 \\ 0 & 0 & -2 & 3 & 0 \end{array} \right]$$

Here $\rho(A) = \rho([A|B]) = 3 < 4$, the number of unknowns

$\therefore$ The system is consistent with one parameter family of solutions, so let $x_4 = t$

Writing the equations, from the row-echelon form we get (from equation (2))

$$\begin{array}{l} 2x_1 - x_4 = 0 \\ \Rightarrow 2x_1 = x_4 \\ \Rightarrow 2x_1 = t \Rightarrow x_1 = \frac{t}{2} \end{array}$$

From equation (3)

$$\begin{array}{l} \Rightarrow 3x_1 - x_3 = 0 \\ \Rightarrow x_3 = 3x_1 \Rightarrow x_3 = \frac{3t}{2} \end{array}$$

From equation (4)

$$\begin{array}{l} 2x_2 - x_3 - 2x_4 = 0 \Rightarrow 2x_2 - \frac{3t}{2} - 2t = 0 \\ \Rightarrow 2x_2 - \frac{7t}{2} = 0 \Rightarrow x_2 = \frac{7t}{4} \end{array}$$


Since x_1, x_2, x_3 and x_4 are positive integers, let us choose $t = 4$

$$\therefore x_1 = \frac{4}{2} = 2$$

$$\therefore x_2 = \frac{7(4)}{4} = 7$$

$$\therefore x_3 = \frac{3(4)}{2} = 6 \text{ and } x_4 = t = 4$$

So, the balanced equation is



EXERCISE 1.8

Choose the Correct or the most suitable answer from the given four alternatives :

1. If $|\text{adj}(\text{adj} A)| = |A|^9$, then the order of the square matrix A is [FRT -2022; Hy. 2024]

- (1) 3 (2) 4 (3) 2 (4) 5

[Ans. (2) 4]

Hint : $|\text{adj}(\text{adj} A)| = |A|^{(n-1)^2}$

$$\therefore (n-1)^2 = 9 \Rightarrow (n-1) = 3 \Rightarrow n = 4$$

2. If A is a 3×3 non-singular matrix such that $AA^T = A^T A$ and $B = A^{-1}A^T$, then $BB^T =$ [Qy. - 2024]

- (1) A (2) B (3) I_3 (4) B^T

[Ans. (3) I_3]

$$\begin{aligned} \text{Hint : } BB^T &= (A^{-1}A^T)(A^{-1}A^T)^T \\ &= (A^{-1}A^T)(A^T)^T \cdot (A^{-1})^T \\ &= (A^{-1}A^T)A(A^{-1})^T = A^{-1}(A \cdot A^T)(A^{-1})^T \\ &= (A^{-1}A) \cdot A^T(A^T)^{-1} [\because (A^{-1})^T = (A^T)^{-1}] \\ &= I \cdot I = I \quad [\because A^T \cdot (A^T)^{-1} = I] \end{aligned}$$

3. If $A = \begin{bmatrix} 3 & 5 \\ 1 & 2 \end{bmatrix}$, $B = \text{adj} A$ and $C = 3A$, then $\frac{|\text{adj} B|}{|C|} =$ [Govt. MQP-2019]

- (1) $\frac{1}{3}$ (2) $\frac{1}{9}$ (3) $\frac{1}{4}$ (4) 1

[Ans. (2) $\frac{1}{9}$]

$$\begin{aligned} \text{Hint : } \frac{|\text{adj} B|}{|C|} &= \frac{|\text{adj}(\text{adj} A)|}{|3A|} = \frac{|A|^{(n-1)^2}}{|3|^2 \cdot |A|} \\ &= \frac{|A|^2}{9 \cdot |A|} = \frac{|A|}{9} = \frac{1}{9} \end{aligned}$$

4. If $A = \begin{bmatrix} 1 & -2 \\ 1 & 4 \end{bmatrix}$, then $A =$

(1) $\begin{bmatrix} 1 & -2 \\ 1 & 4 \end{bmatrix}$

(2) $\begin{bmatrix} 1 & 2 \\ -1 & 4 \end{bmatrix}$

(3) $\begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$

(4) $\begin{bmatrix} 4 & -1 \\ 2 & 1 \end{bmatrix}$

[Ans. (3) $\begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$]

Hint :

$$\begin{aligned} AX &= B \\ \Rightarrow A &= BX^{-1} \text{ where} \\ X &= \begin{bmatrix} 1 & -2 \\ 1 & 4 \end{bmatrix}, B = \begin{bmatrix} 6 & 0 \\ 0 & 6 \end{bmatrix} \\ A &= \begin{bmatrix} 6 & 0 \\ 0 & 6 \end{bmatrix} \cdot \frac{1}{|X|} \cdot \text{adj}(X) \\ &= \begin{bmatrix} 6 & 0 \\ 0 & 6 \end{bmatrix} \cdot \frac{1}{6} \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix} = \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix} \end{aligned}$$

5. If $A = \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}$, then $9I_2 - A =$

[PTA - 2; FRT - 2022; July - 2024]

- (1) A^{-1} (2) $\frac{A^{-1}}{2}$ (3) $3A^{-1}$ (4) $2A^{-1}$

[Ans. (4) $2A^{-1}$]

$$\begin{aligned} \text{Hint : } 9I - A &= \begin{bmatrix} 9 & 0 \\ 0 & 9 \end{bmatrix} - \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 9-7 & 0-3 \\ 0-4 & 9-2 \end{bmatrix} \\ &= \begin{bmatrix} 2 & -3 \\ -4 & 7 \end{bmatrix} = \text{adj} A \end{aligned}$$

$$\text{But } A^{-1} = \frac{1}{|A|} \text{adj} A = \frac{1}{2} \text{adj} A$$

$$\Rightarrow \text{adj} A = 2A^{-1}$$

6. If $A = \begin{bmatrix} 2 & 0 \\ 1 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 4 \\ 2 & 0 \end{bmatrix}$ then $|\text{adj}(AB)| =$ [Qy. - 2023]

- (1) -40 (2) -80 (3) -60 (4) -20

[Ans. (2) -80]

$$\begin{aligned} \text{Hint : } AB &= \begin{bmatrix} 2+0 & 8+0 \\ 1+10 & 4+0 \end{bmatrix} = \begin{bmatrix} 2 & 8 \\ 11 & 4 \end{bmatrix} \\ \text{adj}(AB) &= \begin{bmatrix} 4 & -8 \\ -11 & 2 \end{bmatrix} \\ |\text{adj}(AB)| &= 8 - 88 = -80 \end{aligned}$$

7. If $P = \begin{bmatrix} 1 & x & 0 \\ 1 & 3 & 0 \\ 2 & 4 & -2 \end{bmatrix}$ is the adjoint of 3×3 matrix

A and $|A| = 4$, then x is

- (1) 15 (2) 12 (3) 14 (4) 11

Hint : $|\text{adj } A| = |A|^{n-1}$ **[Ans. (4) 11]**

$$1 \begin{vmatrix} 3 & 0 \\ 4 & -2 \end{vmatrix} - x \begin{vmatrix} 1 & 0 \\ 2 & -2 \end{vmatrix} + 0 = 4^{3-1}$$

$$\Rightarrow -6 - x(-2) = 4^2 \Rightarrow -6 + 2x = 16$$

$$\Rightarrow 2x = 22 \Rightarrow x = 11$$

8. If $A = \begin{bmatrix} 3 & 1 & -1 \\ 2 & -2 & 0 \\ 1 & 2 & -1 \end{bmatrix}$ and $A^{-1} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$

then the value of a_{23} is **[PTA - 5; Qy. - 2024]**

- (1) 0 (2) -2 (3) -3 (4) -1

[Ans. (4) -1]

Hint : $|A| = 3 \begin{vmatrix} -2 & 0 \\ 2 & -1 \end{vmatrix} - 1 \begin{vmatrix} 2 & 0 \\ 1 & -1 \end{vmatrix} - 1 \begin{vmatrix} 2 & -2 \\ 1 & 2 \end{vmatrix}$

$$= 3(2) - 1(-2) - 1(4 + 2) = 6 + 2 - 6 = 2$$

$$a_{23} = \frac{1}{|A|} \text{co-factor of } a_{32} = \frac{1}{2} \times - \begin{vmatrix} 3 & -1 \\ 2 & 0 \end{vmatrix}$$

$$= -\frac{1}{2}(0 + 2) = -\frac{2}{2} = -1$$

9. If A, B and C are invertible matrices of some order, then which one of the following is not true? **[Aug. - 2021; Mar. - 2024]**

- (1) $\text{adj } A = |A|A^{-1}$
 (2) $\text{adj } (AB) = (\text{adj } A)(\text{adj } B)$
 (3) $\det A^{-1} = (\det A)^{-1}$
 (4) $(ABC)^{-1} = C^{-1}B^{-1}A^{-1}$

[Ans. (2) $\text{adj } (AB) = (\text{adj } A)(\text{adj } B)$]

10. If $(AB)^{-1} = \begin{bmatrix} 12 & -17 \\ -19 & 27 \end{bmatrix}$ and $A^{-1} = \begin{bmatrix} 1 & -1 \\ -2 & 3 \end{bmatrix}$, then $B^{-1} =$ **[Mar. - 2020; Hy. - 2023]**

- (1) $\begin{bmatrix} 2 & -5 \\ -3 & 8 \end{bmatrix}$ (2) $\begin{bmatrix} 8 & 5 \\ 3 & 2 \end{bmatrix}$
 (3) $\begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix}$ (4) $\begin{bmatrix} 8 & -5 \\ -3 & 2 \end{bmatrix}$

[Ans. (1) $\begin{bmatrix} 2 & -5 \\ -3 & 8 \end{bmatrix}$]

Hint : Since $(AB)^{-1} = B^{-1}A^{-1}$, we get,

$$\begin{bmatrix} 12 & -17 \\ -19 & 27 \end{bmatrix} = B^{-1} \begin{bmatrix} 1 & -1 \\ -2 & 3 \end{bmatrix}$$

$$\text{Let } X = B^{-1}Y$$

$$B^{-1} = XY^{-1} = \begin{bmatrix} 12 & -17 \\ -19 & 27 \end{bmatrix} \cdot \frac{1}{|Y|} \cdot (\text{adj } Y)$$

$$= \begin{bmatrix} 12 & -17 \\ -19 & 27 \end{bmatrix} \cdot \frac{1}{1} \begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 12 & -17 \\ -19 & 27 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 36-34 & 12-17 \\ -57+54 & -19+27 \end{bmatrix} = \begin{bmatrix} 2 & -5 \\ -3 & 8 \end{bmatrix}$$

11. If $A^T A^{-1}$ is symmetric, then $A^2 =$

[Sep. - 2020; FRT-2022; June & Qy. - 2023; July - 2024]

- (1) A^{-1} (2) $(A^T)^2$
 (3) A^T (4) $(A^{-1})^2$

[Ans : (2) $(A^T)^2$]

Hint : $\Rightarrow A^T A^{-1} = (A^T A^{-1})^T$
 $= (A^{-1})^T (A^T)^T = (A^{-1})^T A$
 $\Rightarrow A^T A^{-1} = (A^{-1})^T A$
 $\Rightarrow A^T A^{-1} = (A^T)^{-1} A$

[Premultiplying by A^T and post multiplying by A on both sides, we get]

$$A^T (A^T A^{-1}) A = A^T [(A^T)^{-1} A] A$$

$$\Rightarrow (A^T)^2 (A^{-1} A) = A^T (A^T)^{-1} A^2$$

$$\Rightarrow (A^T)^2 (I) = I A^2$$

$$\Rightarrow (A^T)^2 = A^2$$

12. If A is a non-singular matrix such that

$$A^{-1} = \begin{bmatrix} 5 & 3 \\ -2 & -1 \end{bmatrix}, \text{ then } (A^T)^{-1} =$$

[PTA - 1; Hy - 2019; July - 2022; Mar. - 2024]

- (1) $\begin{bmatrix} -5 & 3 \\ 2 & 1 \end{bmatrix}$ (2) $\begin{bmatrix} 5 & 3 \\ -2 & -1 \end{bmatrix}$
 (3) $\begin{bmatrix} -1 & -3 \\ 2 & 5 \end{bmatrix}$ (4) $\begin{bmatrix} 5 & -2 \\ 3 & -1 \end{bmatrix}$

Hint : $(A^T)^{-1} = (A^{-1})^T = \begin{bmatrix} 5 & -2 \\ 3 & -1 \end{bmatrix}$ **[Ans : (4) $\begin{bmatrix} 5 & -2 \\ 3 & -1 \end{bmatrix}$]**

13. If $A = \begin{bmatrix} 3 & 4 \\ 5 & 5 \\ x & 3 \\ & 5 \end{bmatrix}$ and $A^T = A^{-1}$, then the value of x is **[July - 2022]**

- (1) $-\frac{4}{5}$ (2) $-\frac{3}{5}$ (3) $\frac{3}{5}$ (4) $\frac{4}{5}$

[Ans: (1) $-\frac{4}{5}$]

Hint : Since $A^T = A^{-1}$, $AA^T = A^T A = I$ [$\therefore$ they are orthogonal]